

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

MINIMUM PARTICULARS TO APPEAR ON THE LABEL WHERE THERE IS NO PACKAGE LEAFLET

Blister card with box containing 1, 2 or 4 pipettes

**4Fleas
protector
Spot-on solution
For Puppie & Small Dogs 0.6kg – 6kg**

**Kills flea eggs and larvae
Prevents flea multiplication**

Johnson's 4fleas protector (2%) for puppies & small sized dogs weighing between 0.6kg and 6kg. Easy spot-on administration. One vial prevents flea multiplication (*Ctenocephalides felis*) in dogs by inhibiting egg and larvae development for 3 months.



DIRECTIONS FOR USE: Do not remove vial from pack until ready for use. For external use only. Use one vial for each animal requiring treatment. Apply Johnson's 4fleas protector (2%) directly on to dog's skin at base of neck, expelling the contents by carefully squeezing bottom of vial 4 times. Care should be taken to maintain contact between vial and skin during application. A second application can be made once, 3 months after the first one, to give a further 3 months' protection.

If you consider that your dog needs repeat treatment after this period, consult your veterinary surgeon.

CONTRA-INDICATIONS AND SPECIAL WARNINGS: Do not allow dogs to swim in ponds and waterways for 3 days following treatment. Do not use on puppies less than 1 month of age, from which fleas may be removed by careful use of a flea comb. Do not use if your dog is unwell or recovering from an illness. If signs of disease persist or appear, consult a veterinary surgeon. In rare cases, transient skin irritation has been reported after use.

FURTHER INFORMATION: If there are several animals in the same household, appropriate treatment should be carried out for each dog or cat, preferably at the same time. A flea elimination strategy should be aimed at all stages of the flea life-cycle. Dogs treated with the product may be at risk of infestation by a new population of adult fleas from outside the household. Infestations occurring during the initial administration or appearing during the treatment period may be eliminated with an additional appropriate product authorised for use on dogs in order to kill adult fleas. Fleas from pets often infest pet's basket, bedding and regular resting areas, such as carpets and soft furnishings. Wash or change the dog's bedding and vacuum carpets and furnishings

regularly, then use a suitable treatment for the household that kills fleas.
The effect of bathing or wetting immediately before or after treatment has not been examined. It is likely that efficacy will be reduced if the dog is wetted or shampooed either before or up to 3 days after treatment.

OPERATOR WARNINGS: Avoid contact with skin and eyes. In case of accidental contact, rinse thoroughly with water. In case of eye irritation, seek medical attention. Wash hands after use.

DISPOSAL ADVICE: Dispose of empty packaging and any remaining product in the household refuse.

The product should not be allowed to enter watercourses as this may be dangerous for fish and other aquatic organisms.

**KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY.
FOR EXTERNAL USE ONLY.**

Contents: 1 vial containing 12mg of Pyriproxyfen in 0.6ml of solution.

AVM-GSL

Vm. 17902/4013

UK authorised veterinary medicinal product
Marketing authorisation holder: Alfamed S.A.S. 06517 Carros Cedex-France
Manufactured by Virbac S.A. 06516 Carros Cedex-France
Johnson's Veterinary Products Ltd,
Sutton Coldfield, West Midlands UK B75 7DF.
Tel: 0121 378 1684 e-mail: info@johnsons-vet.com

Approved: 10/11/2017



