

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Panacur Small Animal 2.5% oral suspension

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Panacur Small Animal 2.5% oral suspension contains 25 mg fenbendazole per ml.

3. PHARMACEUTICAL FORM

A white oral suspension of fenbendazole as a ready to use oral solution for animals

4. PACKAGE SIZE

5. TARGET SPECIES

Domestic dogs, cats, puppies and kittens

6. INDICATION(S)

For the treatment of adult dogs and cats infected with gastro-intestinal roundworms and tapeworms, including:

Toxocara canis, *Toxocara cati*, *Toxascaris leonina*, *Ancylostoma spp*, *Trichuris spp.*, *Uncinaria spp.*, *Taenia spp*

Also effective against lungworms *Oslerus (Filaroides) osleri* or protozoa *Giardia spp.* in dogs and *Aelurostrongylus abstrusus* in cats.

For the treatment of weaned and unweaned puppies and kittens infected with gastrointestinal roundworms particularly *Toxocara canis* and *Ancylostoma spp* and puppies infected with protozoa *Giardia spp*

For the treatment of pregnant dogs to reduce prenatal infections with *Toxocara canis* and the transfer of *T.canis* and *Ancylostoma caninum* to their pups via the milk.

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Routine treatment of adult dogs and cats

4 ml Panacur small animal 2.5% oral suspension per 1 kg (2.2lbs) bodyweight as a single oral dose

<Dosage table>

Panacur is easily administered mixed into feed. Panacur small animal 10% is recommended for larger breeds.

Treatment should be repeated 2 to 4 times a year as natural reinfections with worms occurs. More frequent treatment of at 6 to 8 weekly intervals is advisable for dogs in kennels.

Increased dosing for specific infections:

For the treatment of heavily infected dogs and cats infected with protozoa *Giardia*:

2 ml Panacur Small Animal 2.5% oral suspension per 1 kg bodyweight daily for 3 consecutive days.

For the control of lungworm *Oslerus (Filaroides) osleri* in dogs:

Administer 2 ml Panacur Small Animal 2.5% oral suspension per 1 kg bodyweight, daily for 7 consecutive days. A repeat course of treatment may be required in some cases.

For the control of lungworm *Aelurostrongylus abstrusus* in cats:

Administer 2 ml Panacur Small Animal 2.5% oral suspension per 1 kg bodyweight daily for 3 consecutive days.

Pregnant dogs

1 ml Panacur Small Animal 2.5% oral suspension per 1 kg (2.2lbs) bodyweight daily from day 40 of pregnancy continuously to 2 days post-whelping (approximately 25 days)

<Dosage table>

Panacur small animal 10% is recommended for larger breeds.

The puppies should be wormed at 2 and 5 weeks of age.

Puppies and kittens under six months of age

2 ml Panacur Small Animal 2.5% oral suspension per 1 kg bodyweight daily for 3 consecutive days.

Puppies should be treated at 2 weeks of age, 5 weeks of age and again before leaving the breeder's premises. Treatment may also be required at 8 and 12 weeks of age. Thereafter, frequency of treatment can be reduced unless the pups remain in kennels where reinfestation occurs more readily.

The dose should be administered to unweaned animals into the mouth directly after feeding.

For weaned animals the dose is best administered added to feed, but only add to feed directly prior to feeding.

8. WITHDRAWAL PERIOD

9. SPECIAL WARNING(S), IF NECESSARY

Assess bodyweight as accurately as possible before calculating dosage.

Direct contact with skin should be kept to a minimum. Wear impermeable rubber gloves and wash hands after use.

10. EXPIRY DATE

11. SPECIAL STORAGE CONDITIONS

Keep the container in the outer carton.

Do not freeze

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of empty packaging and any remaining product in the household refuse.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE

[Distribution category]

| |
|---------|
| NFA-VPS |
|---------|

For animal treatment only

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

KEEP OUT OF THE SIGHT AND REACH OF CHILDREN

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

MSD Animal Health UK Ltd.

Walton Manor

Walton

Milton Keynes

MK7 7AJ

16. MARKETING AUTHORISATION NUMBER

Vm 01708/4428

17. MANUFACTURER'S BATCH NUMBER

MINIMUM PARTICULARS TO APPEAR ON IMMEDIATE PACKAGING UNITS
{Bottle}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Panacur Small Animal 2.5% oral suspension

2. QUANTITY OF THE ACTIVE SUBSTANCE(S)

Panacur Small Animal 2.5% oral suspension containing 25 mg fenbendazole per ml.

3. INDICATIONS

For the treatment of adult dogs and cats infected with gastro-intestinal roundworms and tapeworms, including:

Toxocara canis, *Toxocara cati*, *Toxascaris leonina*, *Ancylostoma spp*, *Trichuris spp.*, *Uncinaria spp.*, *Taenia spp*

Also effective against lungworms *Oslerus (Filaroides) osleri* or protozoa *Giardia spp.* in dogs and *Aelurostrongylus abstrusus* in cats.

For the treatment of weaned and unweaned puppies and kittens infected with gastrointestinal roundworms particularly *Toxocara canis* and *Ancylostoma spp* and puppies infected with protozoa *Giardia spp*

For the treatment of pregnant dogs to reduce prenatal infections with *Toxocara canis* and the transfer of *T.canis* and *Ancylostoma caninum* to their pups via the milk.

4. WARNINGS Etc

Shake well before use. Direct contact with skin should be kept to a minimum. Wear impermeable rubber gloves and wash hands after use.

For oral administration only.

Do not freeze

Keep out of the reach and sight of children

Dispose of empty packaging and any remaining product in the household refuse.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

For uses and dosages see pack carton

6. BATCH NUMBER

7. EXPIRY DATE

8. THE WORDS “FOR ANIMAL TREATMENT ONLY”

NFA – VPS For animal treatment only

Vm 01708/4428

MSD Animal Health UK Ltd.

Walton Manor

Walton

Milton Keynes

MK7 7AJ

Approved: 01/04/21

A handwritten signature in blue ink, appearing to read "D. Austin", with a horizontal line extending from the end of the signature.