

ANNEX

Registration number of additive	Name and registration number of person responsible for putting additive into circulation	Additive (Trade name)	Composition, chemical formula, description	Species or category of animal	Maximum age	Minimum content		Maximum content	Other provisions	End of period of authorisation	Provisional maximum residue limits (MRLs) in the relevant foodstuffs of animal origin					
						mg of active substance/kg of complete feedingsstuff	mg of active substance/kg of complete feedingsstuff									
E 757	Eli Lilly and Company Limited	Monensin sodium (Elancoban G100, Elancoban 100, Elancogran 100, Elancoban G200, Elancoban 200)	<p>Active substance</p> <p>$C_{36}H_{61}O_{11}Na$</p> <p>sodium salt of polyether monocarboxylic acid produced by <i>Streptomyces cinnamonensis</i>, ATCC 15413 in granular form.</p> <p>Factor composition:</p> <p>Monensin A: not less than 90 %</p> <p>Monensin A + B: not less than 95 %</p> <p>Additive composition</p> <p>Granular monensin (dried fermentation product) equivalent to Monensin activity 10 % w/w</p> <p>Mineral oil 1-3 % w/w</p> <p>Limestone granular 13-23 % w/w</p> <p>Rice hulls or limestone granular qs 100 % w/w</p> <p>Granular monensin (dried fermentation product) equivalent to Monensin activity 20 % w/w</p> <p>Mineral oil 1-3 % w/w</p> <p>Rice hulls or limestone granular qs 100 % w/w</p>	Chickens for fattening	—	100	125	Use prohibited at least one day before slaughter.	Indicate in the instructions for use: 'Dangerous for equines. This feedingsstuff contains an ionophore: avoid simultaneous administration with tiamulin and monitor for possible adverse reactions when used concurrently with other medicinal substances'	30.7.2014	25 µg/kg monensin sodium of wet skin + fat					
												Chickens reared for laying	16 weeks	100	120	8 µg/kg monensin sodium of wet liver, wet kidney and wet muscle.
												Turkeys	16 weeks	60	100	

Coccidiostats and other medicinal substances