

**B. PACKAGE LEAFLET**  
**100 ml bottle**

**PACKAGE LEAFLET FOR:  
Baycox Multi 50 mg/ml oral suspension for Cattle, Pigs and Sheep**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER  
AND OF THE MANUFACTURING AUTHORISATION HOLDER  
RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Elanco Europe Ltd.  
Form 2, Bartley Way  
Bartley Wood Business Park  
Hook  
RG27 9XA  
United Kingdom

Manufacturer responsible for the batch release:

KVP Pharma + Veterinär Produkte GmbH, Projensdorfer Str. 324, 24106 Kiel,  
Germany

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Baycox Multi 50 mg/ml oral suspension for Cattle, Pigs and Sheep  
For FI, DK, IS, SE: Baycoxine vet 50 mg/ml oral suspension for Cattle, Pigs and  
Sheep  
For NO: Baycoxine vet.  
toltrazuril

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER  
INGREDIENT(S)**

**White or yellowish suspension**

**1 ml contains :**

**Active substance:**

Toltrazuril 50.0 mg

**Excipients:**

Sodium benzoate (E211) 2.1 mg

Sodium propionate (E281) 2.1 mg

**4. INDICATION(S)**

Prevention of clinical signs of coccidiosis (a parasitic disease of the intestinal tract)  
in cattle, pigs and sheep.

Cattle: For the prevention of clinical signs of coccidiosis and reduction of coccidia  
shedding in calves on farms with a confirmed history of coccidiosis caused by  
*Eimeria bovis* or *Eimeria zuernii*.

Pigs: For the prevention of clinical signs of coccidiosis in neonatal piglets (3-5 days old) on farms with a confirmed history of coccidiosis caused by *Cystoisospora suis*.

Sheep: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in lambs on farms with a confirmed history of coccidiosis caused by *Eimeria crandallis* and *Eimeria ovinoidalis*.

## 5. CONTRAINDICATIONS

Do not use in cases of known hypersensitivity to the active substance or any of the excipients.

## 6. ADVERSE REACTIONS

None known.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

## 7. TARGET SPECIES

Cattle (calves: dairy, beef suckler, bull beef), Pigs (Piglets, 3 - 5 days old), Sheep (lambs).

## 8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For oral administration.

### All Species

The ready-to-use oral suspension must be shaken for 20 seconds before use.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

Treatment during an outbreak will be of limited value for the individual animal because of damage to the small intestine having already occurred.

### Cattle

Each animal should be treated with a single oral dose of 15 mg toltrazuril/kg body weight corresponding to 3.0 ml oral suspension per 10 kg body weight.

For the treatment of a group of animals of the same breed and same or similar age, the dosing should be done according to the heaviest animal of this group.

### Pigs

Each pig to be treated on day 3-5 of life with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

Due to the small volumes required to treat individual piglets, use of dosing equipment with a dose accuracy of 0.1 ml is recommended.

### Sheep

Each animal should be treated with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

If animals are to be treated collectively rather than individually, they should be grouped according to their body weight and dosed accordingly, in order to avoid under- or over-dosing.

## **9. ADVICE ON CORRECT ADMINISTRATION**

None

## **10. WITHDRAWAL PERIODS**

### Cattle:

Meat and offal: 63 days

Milk: Not permitted for use in lactating animals producing milk for human consumption.

### Pigs:

Meat and offal: 77 days

### Sheep:

Meat and offal: 42 days

Not permitted for use in lactating sheep producing milk for human consumption.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Do not use this veterinary medicinal product after the expiry date which is stated on the label.

The expiry date refers to the last day of that month.

This veterinary medicinal product does not require any special storage conditions.

Shelf-life after first opening the container: 6 months.

## **12. SPECIAL WARNING(S)**

### Special precautions for use in animals:

As with any antiparasiticide frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance.

It is recommended to treat all animals in a pen.

Hygienic measures may reduce the risk of coccidiosis. It is therefore, recommended to improve concomitantly the hygienic conditions in the concerned facility, particularly dryness and cleanliness.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To alter the course of an established clinical coccidial infection, in individual animals already showing signs of diarrhoea, additional supportive therapy may be required.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known sensitivity to the active substance or any of the excipients should avoid contact with this product.

Avoid skin and eye contact with the product. Wash any splashes from skin or eyes immediately with water.

Do not eat, drink or smoke while using the product.

Other precautions:

The major metabolite of toltrazuril, toltrazuril sulfone (ponazuril), has been shown to be both very persistent (half-life ca. 1 year) and mobile in soil and to be toxic to plants.

For environmental reasons:

Cattle: Not to be used in veal calves.

Do not administer to dairy calves weighing more than 80 kg bodyweight or to bull beef or suckler calves weighing more than 150 kg bodyweight.

Not to be used to treat bull beef calves less than 3 months old.

For dairy calves: In order to prevent any adverse effects on plants and possible contamination of groundwater, manure from treated calves must not be spread onto land without dilution with manure from untreated cows. Manure from treated calves must be diluted with at least 3 times the weight of manure from mature cows before it can be spread onto land.

Sheep: Lambs kept throughout the whole life span indoors under an intensive rearing system must not be treated beyond the age of 6 weeks or body weight of more than 20 kg at treatment. Manure from these animals should only be applied to the same piece of land every third year.

Pigs: None.

Interaction with other medicinal products and other forms of interaction:

None known.

Incompatibilities:

In absence of compatibility studies this veterinary medicinal product must not be mixed with other veterinary medicines.

Overdose (symptoms, emergency procedures, antidotes):

A threefold overdose is well tolerated by healthy piglets and calves without signs of intolerance.

No signs of overdose have been observed in lamb safety studies with threefold overdose at a single treatment and twofold overdose treatment on 2 consecutive days.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

October 2020

**15. OTHER INFORMATION**

Pack sizes: 100, 250 and 1000 ml bottles

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

**B. PACKAGE LEAFLET**  
**250 ml bottle**

**PACKAGE LEAFLET FOR:  
Baycox Multi 50 mg/ml oral suspension for Cattle, Pigs and Sheep**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER  
AND OF THE MANUFACTURING AUTHORISATION HOLDER  
RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Elanco Europe Ltd.  
Form 2, Bartley Way  
Bartley Wood Business Park  
Hook  
RG27 9XA  
United Kingdom

Manufacturer responsible for the batch release:

KVP Pharma + Veterinär Produkte GmbH, Projensdorfer Str. 324, 24106 Kiel,  
Germany

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Baycox Multi 50 mg/ml oral suspension for Cattle, Pigs and Sheep  
For FI, DK, IS, SE: Baycoxine vet 50 mg/ml oral suspension for Cattle, Pigs and  
Sheep  
For NO: Baycoxine vet.  
toltrazuril

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER  
INGREDIENT(S)**

**White or yellowish suspension**

**1 ml contains :**

**Active substance:**

Toltrazuril	50.0 mg
-------------	---------

**Excipients:**

Sodium benzoate (E211)	2.1 mg
Sodium propionate (E281)	2.1 mg

**4. INDICATION(S)**

Prevention of clinical signs of coccidiosis (a parasitic disease of the intestinal tract)  
in cattle, pigs and sheep.

Cattle: For the prevention of clinical signs of coccidiosis and reduction of coccidia  
shedding in calves on farms with a confirmed history of coccidiosis caused by  
*Eimeria bovis* or *Eimeria zuernii*.



Pigs: For the prevention of clinical signs of coccidiosis in neonatal piglets (3-5 days old) on farms with a confirmed history of coccidiosis caused by *Cystoisospora suis*.

Sheep: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in lambs on farms with a confirmed history of coccidiosis caused by *Eimeria crandallis* and *Eimeria ovinoidalis*.

## **5. CONTRAINDICATIONS**

Do not use in cases of known hypersensitivity to the active substance or any of the excipients.

## **6. ADVERSE REACTIONS**

None known.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Cattle (calves: dairy, beef suckler, bull beef), Pigs (Piglets, 3 - 5 days old), Sheep (lambs).

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

For oral administration.

### All Species

The ready-to-use oral suspension must be shaken for 20 seconds before use.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

Treatment during an outbreak will be of limited value for the individual animal because of damage to the small intestine having already occurred.

### Cattle

Each animal should be treated with a single oral dose of 15 mg toltrazuril/kg body weight corresponding to 3.0 ml oral suspension per 10 kg body weight.

For the treatment of a group of animals of the same breed and same or similar age, the dosing should be done according to the heaviest animal of this group.

### Pigs

Each pig to be treated on day 3-5 of life with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

Due to the small volumes required to treat individual piglets, use of dosing equipment

with a dose accuracy of 0.1 ml is recommended.

### Sheep

Each animal should be treated with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

If animals are to be treated collectively rather than individually, they should be grouped according to their body weight and dosed accordingly, in order to avoid under- or over-dosing.

## **9. ADVICE ON CORRECT ADMINISTRATION**

None.

## **10. WITHDRAWAL PERIODS**

### Cattle:

Meat and offal: 63 days

Milk: Not permitted for use in lactating animals producing milk for human consumption.

### Pigs:

Meat and offal: 77 days

### Sheep:

Meat and offal: 42 days

Not permitted for use in lactating sheep producing milk for human consumption.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Do not use this veterinary medicinal product after the expiry date which is stated on the label.

The expiry date refers to the last day of that month.

This veterinary medicinal product does not require any special storage conditions.

Shelf-life after first opening the container: 6 months.

## **12. SPECIAL WARNING(S)**

### Special precautions for use in animals:

As with any antiparasiticide frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance.

It is recommended to treat all animals in a pen.

Hygienic measures may reduce the risk of coccidiosis. It is therefore, recommended to improve concomitantly the hygienic conditions in the concerned facility, particularly dryness and cleanliness.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To alter the course of an established clinical coccidial infection, in individual animals already showing signs of diarrhoea, additional supportive therapy may be required.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known sensitivity to the active substance or any of the excipients should avoid contact with this product.

Avoid skin and eye contact with the product. Wash any splashes from skin or eyes immediately with water.

Do not eat, drink or smoke while using the product.

Other precautions:

The major metabolite of toltrazuril, toltrazuril sulfone (ponazuril), has been shown to be both very persistent (half-life ca. 1 year) and mobile in soil and to be toxic to plants.

For environmental reasons:

Cattle: Not to be used in veal calves.

Do not administer to dairy calves weighing more than 80 kg bodyweight or to bull beef or suckler calves weighing more than 150 kg bodyweight.

Not to be used to treat bull beef calves less than 3 months old.

For dairy calves: In order to prevent any adverse effects on plants and possible contamination of groundwater, manure from treated calves must not be spread onto land without dilution with manure from untreated cows. Manure from treated calves must be diluted with at least 3 times the weight of manure from mature cows before it can be spread onto land.

Sheep: Lambs kept throughout the whole life span indoors under an intensive rearing system must not be treated beyond the age of 6 weeks or body weight of more than 20 kg at treatment. Manure from these animals should only be applied to the same piece of land every third year.

Pigs: None.

Interaction with other medicinal products and other forms of interaction:

None known.

Incompatibilities

In absence of compatibility studies this veterinary medicinal product must not be mixed with other veterinary medicines.

Overdose (symptoms, emergency procedures, antidotes):

A threefold overdose is well tolerated by healthy piglets and calves without signs of intolerance.

No signs of overdose have been observed in lamb safety studies with threefold overdose at a single treatment and twofold overdose treatment on 2 consecutive days.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

October 2020

**15. OTHER INFORMATION**

Pack sizes: 100, 250 and 1000 ml bottles

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

**LABELLING and Outerpack  
for 100 ml and 250 ml**

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE AND OUTER PACKAGE**

**100 ml or 250 ml Plastic Bottle**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Baycox Multi 50 mg/ml oral suspension for Cattle, Pigs and Sheep  
(For DK, FI, IS, SE Baycoxine vet 50 mg/ml oral suspension for Cattle, Pigs and Sheep  
For NO: Baycoxine vet.)

toltrazuril

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

1 ml contains 50 mg toltrazuril

**3. PHARMACEUTICAL FORM**

Oral suspension

**4. PACKAGE SIZE**

100 ml  
250 ml

**5. TARGET SPECIES**

Cattle (calves: dairy, beef suckler, bull beef), Pigs (Piglets, 3 - 5 days old), Sheep (lambs).

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use.  
Read the package leaflet before use.

**8. WITHDRAWAL PERIOD**

Withdrawal periods

Cattle:

Meat and offal: 63 days

Milk: Not permitted for use in lactating animals producing milk for human consumption.

Pigs:

Meat and offal: 77 days

Sheep:

Meat and offal: 42 days

Not permitted for use in lactating sheep producing milk for human consumption.

**9. SPECIAL WARNING(S), IF NECESSARY**

**10. EXPIRY DATE**

EXP {month/year}

Once opened, use by \_\_\_\_\_

Shelf life after first opening the container: 6 months

**11. SPECIAL STORAGE CONDITIONS**

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: Read package leaflet.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.

To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Elanco Europe Ltd.  
Form 2, Bartley Way  
Bartley Wood Business Park  
Hook  
RG27 9XA  
United Kingdom

**16. MARKETING AUTHORISATION NUMBER**

Vm 00879/4114

**17. MANUFACTURER'S BATCH NUMBER**

Batch {number}



**COMBINED LABEL-LEAFLET  
for 1000 ml bottle**

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE FOR THE  
COMBINED LABEL-LEAFLET  
1000 ml Plastic Bottle**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Baycox Multi 50 mg/ml oral suspension for Cattle, Pigs and Sheep  
For FI, DK, IS, SE: Baycoxine vet 50 mg/ml oral suspension for Cattle, Pigs and Sheep  
For NO: Baycoxine vet.

toltrazuril

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

**White or yellowish suspension**

**1 ml contains :**

**Active substance:**

Toltrazuril 50.0 mg

**Excipients:**

Sodium benzoate (E211) 2.1 mg

Sodium propionate (E281) 2.1 mg

**3. PHARMACEUTICAL FORM**

Oral suspension

**4. PACKAGE SIZE**

1000 ml

**5. TARGET SPECIES**

Cattle (calves: dairy, beef suckler, bull beef), Pigs (Piglets, 3 - 5 days old), Sheep (lambs).

**6. INDICATION(S)**

Prevention of clinical signs of coccidiosis (a parasitic disease of the intestinal tract) in cattle, pigs and sheep.

Cattle: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in calves on farms with a confirmed history of coccidiosis caused by *Eimeria bovis* or *Eimeria zuernii*.

Pigs: For the prevention of clinical signs of coccidiosis in neonatal piglets (3 – 5 days old) on farms with a confirmed history of coccidiosis caused by *Cystoisospora suis*.

Sheep: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in lambs on farms with a confirmed history of coccidiosis caused by *Eimeria crandallis* and *Eimeria ovinoidalis*.

## **7. CONTRAINDICATIONS**

Do not use in cases of known hypersensitivity to the active substance or any of the excipients.

## **8. ADVERSE REACTIONS**

None known.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

## **9. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

For oral administration.

### All Species

The ready-to-use oral suspension must be shaken for 20 seconds before use.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

Treatment during an outbreak will be of limited value for the individual animal because of damage to the small intestine having already occurred.

### Cattle

Each animal should be treated with a single oral dose of 15 mg toltrazuril/kg body weight corresponding to 3.0 ml oral suspension per 10 kg body weight.

For the treatment of a group of animals of the same breed and same or similar age, the dosing should be done according to the heaviest animal of this group.

### Pigs

Each pig to be treated on day 3-5 of life with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

Due to the small volumes required to treat individual piglets, use of dosing equipment with a dose accuracy of 0.1 ml is recommended.

Sheep

Each animal should be treated with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

If animals are to be treated collectively rather than individually, they should be grouped according to their body weight and dosed accordingly, in order to avoid under- or over-dosing.

**10. ADVICE ON CORRECT ADMINISTRATION**

None

**11. WITHDRAWAL PERIODS**

Cattle:

Meat and offal: 63 days

Milk: Not permitted for use in lactating animals producing milk for human consumption.

Pigs:

Meat and offal: 77 days

Sheep:

Meat and offal: 42 days

Not permitted for use in lactating sheep producing milk for human consumption.

**12. SPECIAL WARNING(S), IF NECESSARY**

As with any antiparasiticide frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance.

It is recommended to treat all animals in a pen.

Hygienic measures may reduce the risk of coccidiosis. It is therefore, recommended to improve concomitantly the hygienic conditions in the concerned facility, particularly dryness and cleanliness.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To alter the course of an established clinical coccidial infection, in individual animals already showing signs of diarrhoea, additional supportive therapy may be required.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known sensitivity to the active substance or any of the excipients should avoid contact with this product.

Avoid skin and eye contact with the product.

Wash any splashes from skin or eyes immediately with water.

Do not eat, drink or smoke while using the product.

Other precautions:

The major metabolite of toltrazuril, toltrazuril sulfone (ponazuril), has been shown to be both very persistent (half-life ca. 1 year) and mobile in soil and to be toxic to plants.

For environmental reasons:

Cattle: Not to be used in veal calves.

Do not administer to dairy calves weighing more than 80 kg bodyweight or to bull beef or suckler calves weighing more than 150 kg bodyweight.

Not to be used to treat bull beef calves less than 3 months old.

For dairy calves: In order to prevent any adverse effects on plants and possible contamination of groundwater, manure from treated calves must not be spread onto land without dilution with manure from untreated cows. Manure from treated calves must be diluted with at least 3 times the weight of manure from mature cows before it can be spread onto land.

Sheep: Lambs kept throughout the whole life span indoors under an intensive rearing system must not be treated beyond the age of 6 weeks or body weight of more than 20 kg at treatment. Manure from these animals should only be applied to the same piece of land every third year.

Pigs: None.

Interaction with other medicinal products and other forms of interaction:

None known.

Incompatibilities:

In absence of compatibility studies this veterinary medicinal product must not be mixed with other veterinary medicines.

Overdose (symptoms, emergency procedures, antidotes):

A threefold overdose is well tolerated by healthy piglets and calves without signs of intolerance.

No signs of overdose have been observed in lamb safety studies with threefold overdose at a single treatment and twofold overdose treatment on 2 consecutive days.

**13. EXPIRY DATE**

EXP {month/year}

Once opened, use by \_\_\_\_\_

Shelf life after first opening the container: 6 months

Do not use this veterinary medicinal product after the expiry date which is stated on the label. The expiry date refers to the last day of that month.

**14. SPECIAL STORAGE CONDITIONS / PRECAUTIONS**

The veterinary medicinal product does not require any special storage conditions.

**15. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

**16. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.  
To be supplied only on veterinary prescription.

**17. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**18. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing Authorisation Holder:  
Elanco Europe Ltd.  
Form 2, Bartley Way  
Bartley Wood Business Park  
Hook  
RG27 9XA  
United Kingdom

Manufacturing Authorisation Holder responsible for batch release:  
KVP Pharma + Veterinär Produkte GmbH  
Projensdorfer Straße 324, 24106 Kiel  
Germany

**19. MARKETING AUTHORISATION NUMBER**

Vm 00879/4114

**20. MANUFACTURER’S BATCH NUMBER**

Batch {number}

**21. DATE ON WHICH THE TEXT WAS LAST APPROVED**

October 2020

## **22. OTHER INFORMATION**

Pack sizes: 100, 250 and 1000 ml bottles

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

**B. PACKAGE LEAFLET**  
**250 ml bottle**



**PACKAGE LEAFLET FOR:  
Baycox Multi 50 mg/ml oral suspension for Cattle, Pigs and Sheep**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER  
AND OF THE MANUFACTURING AUTHORISATION HOLDER  
RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Elanco Europe Ltd.  
Form 2, Bartley Way  
Bartley Wood Business Park  
Hook  
RG27 9XA  
United Kingdom

Manufacturer responsible for the batch release:

KVP Pharma + Veterinär Produkte GmbH, Projensdorfer Str. 324, 24106 Kiel,  
Germany

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Baycox Multi 50 mg/ml oral suspension for Cattle, Pigs and Sheep  
For FI, DK, IS, SE: Baycoxine vet 50 mg/ml oral suspension for Cattle, Pigs and  
Sheep  
For NO: Baycoxine vet.  
toltrazuril

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER  
INGREDIENT(S)**

**White or yellowish suspension**

**1 ml contains :**

**Active substance:**

Toltrazuril	50.0 mg
-------------	---------

**Excipients:**

Sodium benzoate (E211)	2.1 mg
Sodium propionate (E281)	2.1 mg

**4. INDICATION(S)**

Prevention of clinical signs of coccidiosis (a parasitic disease of the intestinal tract)  
in cattle, pigs and sheep.

Cattle: For the prevention of clinical signs of coccidiosis and reduction of coccidia  
shedding in calves on farms with a confirmed history of coccidiosis caused by  
*Eimeria bovis* or *Eimeria zuernii*.

Pigs: For the prevention of clinical signs of coccidiosis in neonatal piglets (3-5 days old) on farms with a confirmed history of coccidiosis caused by *Cystoisospora suis*.

Sheep: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in lambs on farms with a confirmed history of coccidiosis caused by *Eimeria crandallis* and *Eimeria ovinoidalis*.

## 5. CONTRAINDICATIONS

Do not use in cases of known hypersensitivity to the active substance or any of the excipients.

## 6. ADVERSE REACTIONS

None known.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

## 7. TARGET SPECIES

Cattle (calves: dairy calves, beef suckler, bull beef), Pigs (Piglets, 3 - 5 days old), Sheep (lambs).

## 8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For oral administration.

### All Species

The ready-to-use oral suspension must be shaken for 20 seconds before use.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

Treatment during an outbreak will be of limited value for the individual animal because of damage to the small intestine having already occurred.

### Cattle

Each animal should be treated with a single oral dose of 15 mg toltrazuril/kg body weight corresponding to 3.0 ml oral suspension per 10 kg body weight.

For the treatment of a group of animals of the same breed and same or similar age, the dosing should be done according to the heaviest animal of this group.

### Pigs

Each pig to be treated on day 3-5 of life with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

Due to the small volumes required to treat individual piglets, use of dosing equipment

with a dose accuracy of 0.1 ml is recommended.

#### Sheep

Each animal should be treated with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

If animals are to be treated collectively rather than individually, they should be grouped according to their body weight and dosed accordingly, in order to avoid under- or over-dosing.

### **9. ADVICE ON CORRECT ADMINISTRATION**

None.

### **10. WITHDRAWAL PERIODS**

#### Cattle:

Meat and offal: 63 days

Milk: Not permitted for use in lactating animals producing milk for human consumption.

#### Pigs:

Meat and offal: 77 days

#### Sheep:

Meat and offal: 42 days

Not permitted for use in lactating sheep producing milk for human consumption.

### **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Do not use this veterinary medicinal product after the expiry date which is stated on the label.

The expiry date refers to the last day of that month.

This veterinary medicinal product does not require any special storage conditions.

Shelf-life after first opening the container: 6 months.

### **12. SPECIAL WARNING(S)**

#### Special precautions for use in animals:

As with any antiparasiticide frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance.

It is recommended to treat all animals in a pen.

Hygienic measures may reduce the risk of coccidiosis. It is therefore, recommended to improve concomitantly the hygienic conditions in the concerned facility, particularly dryness and cleanliness.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To alter the course of an established clinical coccidial infection, in individual animals already showing signs of diarrhoea, additional supportive therapy may be required.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known sensitivity to the active substance or any of the excipients should avoid contact with this product.

Avoid skin and eye contact with the product. Wash any splashes from skin or eyes immediately with water.

Do not eat, drink or smoke while using the product.

Other precautions:

The major metabolite of toltrazuril, toltrazuril sulfone (ponazuril), has been shown to be both very persistent (half-life ca. 1 year) and mobile in soil and to be toxic to plants.

For environmental reasons:

Cattle: Not to be used in veal calves.

Do not administer to dairy calves weighing more than 80 kg bodyweight or to bull beef or suckler calves weighing more than 150 kg bodyweight.

Not to be used to treat bull beef calves less than 3 months old.

For dairy calves: In order to prevent any adverse effects on plants and possible contamination of groundwater, manure from treated calves must not be spread onto land without dilution with manure from untreated cows. Manure from treated calves must be diluted with at least 3 times the weight of manure from mature cows before it can be spread onto land.

Sheep: Lambs kept throughout the whole life span indoors under an intensive rearing system must not be treated beyond the age of 6 weeks or body weight of more than 20 kg at treatment. Manure from these animals should only be applied to the same piece of land every third year.

Pigs: None.

Interaction with other medicinal products and other forms of interaction:

None known.

Incompatibilities

In absence of compatibility studies this veterinary medicinal product must not be mixed with other veterinary medicines.

Overdose (symptoms, emergency procedures, antidotes):

A threefold overdose is well tolerated by healthy piglets and calves without signs of intolerance.

No signs of overdose have been observed in lamb safety studies with threefold overdose at a single treatment and twofold overdose treatment on 2 consecutive days.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

October 2020

**15. OTHER INFORMATION**

Pack sizes: 100, 250 and 1000 ml bottles

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

**B. PACKAGE LEAFLET**  
**250 ml bottle**

**PACKAGE LEAFLET FOR:  
Baycox Multi 50 mg/ml oral suspension for Cattle, Pigs and Sheep**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER  
AND OF THE MANUFACTURING AUTHORISATION HOLDER  
RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Elanco Europe Ltd.  
Form 2, Bartley Way  
Bartley Wood Business Park  
Hook  
RG27 9XA  
United Kingdom

Manufacturer responsible for the batch release:

KVP Pharma + Veterinär Produkte GmbH, Projensdorfer Str. 324, 24106 Kiel,  
Germany

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Baycox Multi 50 mg/ml oral suspension for Cattle, Pigs and Sheep  
For FI, DK, IS, SE: Baycoxine vet 50 mg/ml oral suspension for Cattle, Pigs and  
Sheep  
For NO: Baycoxine vet.  
toltrazuril

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER  
INGREDIENT(S)**

**White or yellowish suspension**

**1 ml contains :**

**Active substance:**

Toltrazuril	50.0 mg
-------------	---------

**Excipients:**

Sodium benzoate (E211)	2.1 mg
Sodium propionate (E281)	2.1 mg

**4. INDICATION(S)**

Prevention of clinical signs of coccidiosis (a parasitic disease of the intestinal tract)  
in cattle, pigs and sheep.

Cattle: For the prevention of clinical signs of coccidiosis and reduction of coccidia  
shedding in calves on farms with a confirmed history of coccidiosis caused by  
*Eimeria bovis* or *Eimeria zuernii*.

Pigs: For the prevention of clinical signs of coccidiosis in neonatal piglets (3-5 days old) on farms with a confirmed history of coccidiosis caused by *Cystoisospora suis*.

Sheep: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in lambs on farms with a confirmed history of coccidiosis caused by *Eimeria crandallis* and *Eimeria ovinoidalis*.

## 5. CONTRAINDICATIONS

Do not use in cases of known hypersensitivity to the active substance or any of the excipients.

## 6. ADVERSE REACTIONS

None known.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

## 7. TARGET SPECIES

Cattle (calves: dairy calves, beef suckler, bull beef), Pigs (Piglets, 3 - 5 days old), Sheep (lambs).

## 8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For oral administration.

### All Species

The ready-to-use oral suspension must be shaken for 20 seconds before use.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

Treatment during an outbreak will be of limited value for the individual animal because of damage to the small intestine having already occurred.

### Cattle

Each animal should be treated with a single oral dose of 15 mg toltrazuril/kg body weight corresponding to 3.0 ml oral suspension per 10 kg body weight.

For the treatment of a group of animals of the same breed and same or similar age, the dosing should be done according to the heaviest animal of this group.

### Pigs

Each pig to be treated on day 3-5 of life with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

Due to the small volumes required to treat individual piglets, use of dosing equipment



with a dose accuracy of 0.1 ml is recommended.

#### Sheep

Each animal should be treated with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

If animals are to be treated collectively rather than individually, they should be grouped according to their body weight and dosed accordingly, in order to avoid under- or over-dosing.

### **9. ADVICE ON CORRECT ADMINISTRATION**

None.

### **10. WITHDRAWAL PERIODS**

#### Cattle:

Meat and offal: 63 days

Milk: Not permitted for use in lactating animals producing milk for human consumption.

#### Pigs:

Meat and offal: 77 days

#### Sheep:

Meat and offal: 42 days

Not permitted for use in lactating sheep producing milk for human consumption.

### **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Do not use this veterinary medicinal product after the expiry date which is stated on the label.

The expiry date refers to the last day of that month.

This veterinary medicinal product does not require any special storage conditions.

Shelf-life after first opening the container: 6 months.

### **12. SPECIAL WARNING(S)**

#### Special precautions for use in animals:

As with any antiparasiticide frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance.

It is recommended to treat all animals in a pen.

Hygienic measures may reduce the risk of coccidiosis. It is therefore, recommended to improve concomitantly the hygienic conditions in the concerned facility, particularly dryness and cleanliness.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To alter the course of an established clinical coccidial infection, in individual animals already showing signs of diarrhoea, additional supportive therapy may be required.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known sensitivity to the active substance or any of the excipients should avoid contact with this product.

Avoid skin and eye contact with the product. Wash any splashes from skin or eyes immediately with water.

Do not eat, drink or smoke while using the product.

Other precautions:

The major metabolite of toltrazuril, toltrazuril sulfone (ponazuril), has been shown to be both very persistent (half-life ca. 1 year) and mobile in soil and to be toxic to plants.

For environmental reasons:

Cattle: Not to be used in veal calves.

Do not administer to dairy calves weighing more than 80 kg bodyweight or to bull beef or suckler calves weighing more than 150 kg bodyweight.

Not to be used to treat bull beef calves less than 3 months old.

For dairy calves: In order to prevent any adverse effects on plants and possible contamination of groundwater, manure from treated calves must not be spread onto land without dilution with manure from untreated cows. Manure from treated calves must be diluted with at least 3 times the weight of manure from mature cows before it can be spread onto land.

Sheep: Lambs kept throughout the whole life span indoors under an intensive rearing system must not be treated beyond the age of 6 weeks or body weight of more than 20 kg at treatment. Manure from these animals should only be applied to the same piece of land every third year.

Pigs: None.

Interaction with other medicinal products and other forms of interaction:

None known.

Incompatibilities

In absence of compatibility studies this veterinary medicinal product must not be mixed with other veterinary medicines.

Overdose (symptoms, emergency procedures, antidotes):

A threefold overdose is well tolerated by healthy piglets and calves without signs of intolerance.

No signs of overdose have been observed in lamb safety studies with threefold overdose at a single treatment and twofold overdose treatment on 2 consecutive days.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

October 2020

**15. OTHER INFORMATION**

Pack sizes: 100, 250 and 1000 ml bottles

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

**LABELLING and Outerpack  
for 100 ml and 250 ml**

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE AND OUTER PACKAGE**

**100 ml or 250 ml Plastic Bottle**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Baycox Multi 50 mg/ml oral suspension for Cattle, Pigs and Sheep  
(For DK, FI, IS, SE Baycoxine vet 50 mg/ml oral suspension for Cattle, Pigs and Sheep  
For NO: Baycoxine vet.)

toltrazuril

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

1 ml contains 50 mg toltrazuril

**3. PHARMACEUTICAL FORM**

Oral suspension

**4. PACKAGE SIZE**

100 ml  
250 ml

**5. TARGET SPECIES**

Cattle (calves: dairy calves, beef suckler, bull beef), Pigs (Piglets, 3 - 5 days old),  
Sheep (lambs).

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use.  
Read the package leaflet before use.

**8. WITHDRAWAL PERIOD**

Withdrawal periods

Cattle:

Meat and offal: 63 days

Milk: Not permitted for use in lactating animals producing milk for human consumption.

Pigs:

Meat and offal: 77 days

Sheep:

Meat and offal: 42 days

Not permitted for use in lactating sheep producing milk for human consumption.

**9. SPECIAL WARNING(S), IF NECESSARY**

**10. EXPIRY DATE**

EXP {month/year}

Once opened, use by \_\_\_\_\_

Shelf life after first opening the container: 6 months

**11. SPECIAL STORAGE CONDITIONS**

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: Read package leaflet.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.

To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Elanco Europe Ltd.  
Form 2, Bartley Way  
Bartley Wood Business Park  
Hook  
RG27 9XA  
United Kingdom

**16. MARKETING AUTHORISATION NUMBER**

Vm 00879/4114

**17. MANUFACTURER'S BATCH NUMBER**

Batch {number}

**COMBINED LABEL-LEAFLET  
for 1000 ml bottle**



**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE FOR THE  
COMBINED LABEL-LEAFLET  
1000 ml Plastic Bottle**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Baycox Multi 50 mg/ml oral suspension for Cattle, Pigs and Sheep  
For FI, DK, IS, SE: Baycoxine vet 50 mg/ml oral suspension for Cattle, Pigs and Sheep  
For NO: Baycoxine vet.

toltrazuril

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

**White or yellowish suspension**

**1 ml contains :**

**Active substance:**

Toltrazuril 50.0 mg

**Excipients:**

Sodium benzoate (E211) 2.1 mg

Sodium propionate (E281) 2.1 mg

**3. PHARMACEUTICAL FORM**

Oral suspension

**4. PACKAGE SIZE**

1000 ml

**5. TARGET SPECIES**

Cattle (calves: dairy calves, beef suckler, bull beef), Pigs (Piglets, 3 - 5 days old),  
Sheep (lambs).

**6. INDICATION(S)**

Prevention of clinical signs of coccidiosis (a parasitic disease of the intestinal tract) in cattle, pigs and sheep.

Cattle: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in calves on farms with a confirmed history of coccidiosis caused by *Eimeria bovis* or *Eimeria zuernii*.

Pigs: For the prevention of clinical signs of coccidiosis in neonatal piglets (3 – 5 days old) on farms with a confirmed history of coccidiosis caused by *Cystoisospora suis*.

Sheep: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in lambs on farms with a confirmed history of coccidiosis caused by *Eimeria crandallis* and *Eimeria ovinoidalis*.

## **7. CONTRAINDICATIONS**

Do not use in cases of known hypersensitivity to the active substance or any of the excipients.

## **8. ADVERSE REACTIONS**

None known.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

## **9. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

For oral administration.

### All Species

The ready-to-use oral suspension must be shaken for 20 seconds before use.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

Treatment during an outbreak will be of limited value for the individual animal because of damage to the small intestine having already occurred.

### Cattle

Each animal should be treated with a single oral dose of 15 mg toltrazuril/kg body weight corresponding to 3.0 ml oral suspension per 10 kg body weight.

For the treatment of a group of animals of the same breed and same or similar age, the dosing should be done according to the heaviest animal of this group.

### Pigs

Each pig to be treated on day 3-5 of life with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

Due to the small volumes required to treat individual piglets, use of dosing equipment with a dose accuracy of 0.1 ml is recommended.

### Sheep

Each animal should be treated with a single oral dose of 20 mg toltrazuril/kg body weight corresponding to 0.4 ml oral suspension per kg body weight.

If animals are to be treated collectively rather than individually, they should be grouped according to their body weight and dosed accordingly, in order to avoid under- or over-dosing.

## 10. ADVICE ON CORRECT ADMINISTRATION

None

## 11. WITHDRAWAL PERIODS

### Cattle:

Meat and offal: 63 days

Milk: Not permitted for use in lactating animals producing milk for human consumption.

### Pigs:

Meat and offal: 77 days

### Sheep:

Meat and offal: 42 days

Not permitted for use in lactating sheep producing milk for human consumption.

## 12. SPECIAL WARNING(S), IF NECESSARY

As with any antiparasiticide frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance.

It is recommended to treat all animals in a pen.

Hygienic measures may reduce the risk of coccidiosis. It is therefore, recommended to improve concomitantly the hygienic conditions in the concerned facility, particularly dryness and cleanliness.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To alter the course of an established clinical coccidial infection, in individual animals already showing signs of diarrhoea, additional supportive therapy may be required.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known sensitivity to the active substance or any of the excipients should avoid contact with this product.

Avoid skin and eye contact with the product.

Wash any splashes from skin or eyes immediately with water.

Do not eat, drink or smoke while using the product.

### Other precautions:

The major metabolite of toltrazuril, toltrazuril sulfone (ponazuril), has been shown to be both very persistent (half-life ca. 1 year) and mobile in soil and to be toxic to plants.

For environmental reasons:

Cattle: Not to be used in veal calves.

Do not administer to dairy calves weighing more than 80 kg bodyweight or to bull beef or suckler calves weighing more than 150 kg bodyweight.

Not to be used to treat bull beef calves less than 3 months old.

For dairy calves: In order to prevent any adverse effects on plants and possible contamination of groundwater, manure from treated calves must not be spread onto land without dilution with manure from untreated cows. Manure from treated calves must be diluted with at least 3 times the weight of manure from mature cows before it can be spread onto land.

Sheep: Lambs kept throughout the whole life span indoors under an intensive rearing system must not be treated beyond the age of 6 weeks or body weight of more than 20 kg at treatment. Manure from these animals should only be applied to the same piece of land every third year.

Pigs: None.

Interaction with other medicinal products and other forms of interaction:

None known.

Incompatibilities:

In absence of compatibility studies this veterinary medicinal product must not be mixed with other veterinary medicines.

Overdose (symptoms, emergency procedures, antidotes):

A threefold overdose is well tolerated by healthy piglets and calves without signs of intolerance.

No signs of overdose have been observed in lamb safety studies with threefold overdose at a single treatment and twofold overdose treatment on 2 consecutive days.

### **13. EXPIRY DATE**

EXP {month/year}

Once opened, use by \_\_\_\_\_

Shelf life after first opening the container: 6 months

Do not use this veterinary medicinal product after the expiry date which is stated on the label. The expiry date refers to the last day of that month.

### **14. SPECIAL STORAGE CONDITIONS / PRECAUTIONS**

The veterinary medicinal product does not require any special storage conditions.

**15. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

**16. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.  
To be supplied only on veterinary prescription.

**17. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**18. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing Authorisation Holder:  
Elanco Europe Ltd.  
Form 2, Bartley Way  
Bartley Wood Business Park  
Hook  
RG27 9XA  
United Kingdom

Manufacturing Authorisation Holder responsible for batch release:  
KVP Pharma + Veterinär Produkte GmbH  
Projensdorfer Straße 324, 24106 Kiel  
Germany

**19. MARKETING AUTHORISATION NUMBER**

Vm 00879/4114

**20. MANUFACTURER'S BATCH NUMBER**

Batch {number}

**21. DATE ON WHICH THE TEXT WAS LAST APPROVED**

To be allocated

## 22. OTHER INFORMATION

Pack sizes: 100, 250 and 1000 ml bottles

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

Approved 23 October 2020

A handwritten signature in black ink, appearing to read "Hunter.", is positioned below the approval date. The signature is stylized and includes a period at the end.