

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

CLiK Extra 65 mg/ml Pour-On Suspension for Sheep.  
Dicyclanil

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each ml contains:  
Dicyclanil 65 mg

**Excipients:**

Methyl parahydroxybenzoate (E218)	1.50 mg
Propyl parahydroxybenzoate	3.00 mg
Butylated hydroxytoluene	0.50 mg
Ponceau 4R	0.05 mg

**3. PHARMACEUTICAL FORM**

Pink coloured Suspension.

**4. PACKAGE SIZE**

1 litre (filled with 0.8 litres of product)  
2.5 litres (filled with 2.2 litres of product)  
5 litres (filled with 5 litres of product)

**5. TARGET SPECIES**

Sheep

**6. INDICATION(S)**

Prevention of blowfly strike on sheep due to *Lucilia sericata* or *Wohlfahrtia magnifica*.

**7. CONTRAINDICATIONS**

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

**8. ADVERSE REACTIONS**

None known

If you notice any serious effects or other effects not mentioned on this label, please inform

your veterinary surgeon.

## 9. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF

CLiK Extra 65 mg/ml Pour-On Suspension for Sheep is applied according to the following recommendations:

Bodyweight (kg)	Dose Volume (ml)
10-20	20
21-30	24
31-50	30
>50	36

(Guide dose volumes correspond to 0.6–2 ml (39–130 mg dicyclanil) per kg bodyweight.)

To ensure administration of a correct dose, body weight should be determined as accurately as possible; accuracy of the dosing device should be checked.


If animals are to be treated collectively rather than individually, they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- or overdosing.

Recommended for treatment in sheep with any wool lengths; including off-shears.

The product must be applied with a manual or automatic dosing gun, fitted with a spray nozzle, which guarantees the correct spreading of the product on the fleece.

Best results will be achieved by holding the gun approximately 45 cm from the sheep during application. Apply as a fan spray along the spine of the animal in bands at least 10 cm wide from the

middle of the shoulders and in an arc around the crutch and tail. Half the dose should be applied along the spine with the remainder over the tail and crutch area.



CLiK Extra should be applied before, or at the start of, predicted fly activity.

The product will protect against fly strike for 19 weeks. It is good agricultural practice to check animals regularly for fly strike.

## **10. ADVICE ON CORRECT ADMINISTRATION**

Please refer to section “Special Warnings” for further advice on administration.

## **11. WITHDRAWAL PERIOD**

Meat and offal: 40 days

Not authorised for use in animals producing milk for human consumption.

## **12. SPECIAL WARNING(S), IF NECESSARY**

For external use only

### Special warnings for target animals

CLiK Extra is best applied before an anticipated blowfly challenge, or when a blowfly challenge is identified on or in the vicinity of the farm. Established strikes may require a separate treatment with a knockdown insecticide.

It is recommended that animals with dirty back-ends are dagged or crutched prior to application. If dagging or crutching is undertaken in the weeks following application, these animals should be re- treated, otherwise protection could be lost.

Do not apply during heavy rainfall, or when such conditions are expected. The resulting wash out may reduce the protection period.

Shake the container well before

use.

Do not dilute with water.

### User Warnings

- Redness and irritation may develop after skin or eye contact with the product.
- Contact with skin and eyes should be avoided.
- Personal protective equipment consisting of synthetic rubber gloves and PVC trousers should be worn when handling the product.
- In case of skin contact remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water.
- In case of eye contact wash immediately with clean water.
- Always wash hands and exposed skin with soap and water after work.
- Do not eat, drink or smoke whilst using the product.
- Residues remain on the fleece for some time after treatment, therefore, it is good agricultural practice to minimise handling of sheep after treatment. If you need to handle sheep within 3 months after treatment, wear synthetic rubber gloves and long trousers or coveralls. If sheep are wet wear waterproof trousers.
- Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

The safety of the product has not been established during pregnancy or lactation. Use only

accordingly to the benefit/risk assessment by the responsible veterinarian. Laboratory studies have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects.

#### Other precautions

Treated sheep must be kept away from watercourses for at least one hour after treatment. There is a serious risk to aquatic life if this advice is not followed.

Dicyclanil has the potential to cause harmful effects on aquatic invertebrates and dung fly larvae.

Following use of this product, levels of dicyclanil potentially harmful to dung fly larvae have been shown to be excreted in faeces for approximately 4 weeks. Faeces from treated animals may temporarily reduce the abundance of dung fly larvae which may impact on dung degradation.

### **13. EXPIRY DATE**

<EXP {month/year}>

Shelf-life after first opening the immediate packaging: 12 months.

When the container is opened for the first time, using the in-use shelf life which is specified on this label, the date on which any product remaining in the container should be discarded should be determined. This discard date should be written in the space provided \_\_\_\_\_

### **14. SPECIAL STORAGE CONDITIONS/PRECAUTIONS**

Store in the original container.

Keep the container tightly closed, away from food, drink and animal feedstuffs. Protect from direct sunlight.

Protect from frost

### **15. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Harmful to aquatic and dung fauna.

Do not contaminate ponds or other waterways with product or empty containers.

### **16. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE,**

For animal treatment only

**17. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children

**18. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder

Elanco Europe Ltd  
Form 2, Bartley Way  
Bartley Wood Business Park  
Hook  
RG27 9XA  
United Kingdom

Manufacturer responsible for batch release

Elanco France S.A.S  
26 Rue de la Chapelle  
68330 Huningue  
France

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

**19. MARKETING AUTHORISATION NUMBER**

Vm 00879/4003

**20. MANUFACTURER'S BATCH NUMBER**

<Batch> <Lot> <BN> {number}

**21. DATE ON WHICH TEXT WAS LAST APPROVED**

December 2020

**22. OTHER INFORMATION**

1 L, 2.5 L and 5 L white opaque polyethylene back pack containers filled with 0.8 L, 2.2 L and 5 L, respectively, and closed with a polypropylene screw cap.  
Not all pack sizes may be marketed.

## **B. PACKAGE LEAFLET**

Inclusion of a package leaflet in the packaging of CLiK Extra 65 mg/ml Pour-On Suspension for Sheep is not intended as all information is contained on the immediate packaging.

Approved: 26/04/21

A handwritten signature in black ink, appearing to read "D. Austin", with a horizontal line extending to the right.