

LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Cardboard box

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Porcilis Ery+Parvo+Lepto suspension for injection for pigs

2. STATEMENT OF ACTIVE SUBSTANCES

Inactivated *Erysipelothrix rhusiopathiae*, Porcine parvovirus and *Leptospira*

3. PHARMACEUTICAL FORM

Suspension for injection.

4. PACKAGE SIZE

20 ml (10 doses)
10x 20 ml (10 doses)
50 ml (25 doses)
10x 50 ml (25 doses)
100 ml (50 doses)
250 ml (125 doses)

5. TARGET SPECIES

Pig for reproduction

6. INDICATION(S)

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Intramuscular use in the neck.
Read the package leaflet before use.

8. WITHDRAWAL PERIOD(S)

Withdrawal period: zero days.

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

10. EXPIRY DATE

EXP {month/year}

Once broached use within 10 hours.

11. SPECIAL STORAGE CONDITIONS

Store in a refrigerator.

Do not freeze.

Protect from light.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: read package leaflet.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only. To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

MSD Animal Health UK Ltd.
Walton Manor, Walton
Milton Keynes
MK7 7AJ

16. MARKETING AUTHORISATION NUMBER(S)

Vm 01708/4633

17. MANUFACTURER'S BATCH NUMBER

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

PET vials (20 ml)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Porcilis Ery+Parvo+Lepto for pigs

2. QUANTITY OF THE ACTIVE SUBSTANCE(S)

Inactivated *Erysipelothrix rhusiopathiae*, Porcine parvovirus and *Leptospira*

3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

20 ml (10 doses)

4. ROUTE(S) OF ADMINISTRATION

i.m.

5. WITHDRAWAL PERIOD(S)

Withdrawal period: zero days.

6. BATCH NUMBER

Lot {number}

7. EXPIRY DATE

EXP {month/year}

Once broached use within 10 hours.

8. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

PARTICULARS TO APPEAR ON IMMEDIATE PACKAGING UNITS

PET vials (50, 100 and 250 ml)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Porcilis Ery+Parvo+Lepto suspension for injection for pigs

2. STATEMENT OF ACTIVE SUBSTANCES

Inactivated *Erysipelothrix rhusiopathiae*, Porcine parvovirus and *Leptospira*

3. PHARMACEUTICAL FORM

Suspension for injection.

4. PACKAGE SIZE

50 ml (25 doses)
100 ml (50 doses)
250 ml (125 doses)

5. TARGET SPECIES

Pig for reproduction

6. INDICATION(S)

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Intramuscular injection.
Read the package leaflet before use.

8. WITHDRAWAL PERIOD(S)

Withdrawal period: zero days.

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

10. EXPIRY DATE

EXP {month/year}
Once broached use within 10 hours.

11. SPECIAL STORAGE CONDITIONS

Store in a refrigerator.
Do not freeze.
Protect from light.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: Read package leaflet.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only. To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

MSD Animal Health UK Ltd.
Walton Manor, Walton
Milton Keynes
MK7 7AJ

16. MARKETING AUTHORISATION NUMBER(S)

Vm 01708/4633

17. MANUFACTURER'S BATCH NUMBER

Lot {number}

B. PACKAGE LEAFLET

PACKAGE LEAFLET:
Porcilis Ery+Parvo+Lepto suspension for injection for pigs

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

MSD Animal Health UK Ltd.
Walton Manor, Walton
Milton Keynes
Buckinghamshire
MK7 7AJ

Manufacturer responsible for batch release:

Intervet International B.V.
Wim de Körverstraat 35
5831 AN Boxmeer
The Netherlands

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Porcilis Ery+Parvo+Lepto suspension for injection for pigs

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each dose of 2 ml contains:

Active substances:

Inactivated strains of:

Erysipelothrix rhusiopathiae, serotype 2 (strain M2) ≥ 1 ppd¹
Porcine parvovirus (strain 014) ≥ 130
U²

Leptospira interrogans serogroup Canicola serovar Portland-Vere (strain Ca-12-000) ≥ 2816 U²

L. eptospira interrogans serogroup Icterohaemorrhagiae serovar Copenhageni (strain Ic-02-001) ≥ 210
U²

Leptospira interrogans serogroup Australis serovar Bratislava (strain As-05-073) ≥ 1310 U²

Leptospira kirschneri serogroup Grippotyphosa serovar Dadas (strain Gr-01-005) ≥ 648 U²

Leptospira interrogans serogroup Pomona serovar Pomona (strain Po-01-000) ≥ 166 U²

Leptospira santarosai serogroup Tarassovi serovar Gatuni (strain S1148/02) ≥ 276 U²

Adjuvant:

dl- α -tocopheryl acetate 150
mg

¹ Pig protective dose as compared to a reference preparation known to be protective in pigs.

² As determined in the *in vitro* antigenic mass ELISA potency test.

Suspension for injection.

Homogenous white to nearly white suspension after shaking.

4. INDICATION(S)

For the active immunization of pigs:

- to reduce clinical signs (skin lesions and fever) of swine erysipelas caused by *Erysipelothrix rhusiopathiae*, serotype 1 and serotype 2
- to reduce transplacental infection, viral load and foetal mortality caused by Porcine parvovirus.
- to reduce clinical signs (increase of body temperature and reduction in feed intake or activity), infection and bacterial excretion caused by *L. interrogans* serogroup Canicola serovar Canicola
- to reduce clinical signs (increase of body temperature and reduction in feed intake or activity), severity of infection and foetal mortality caused by *L. interrogans* serogroup Pomona serovar Pomona.
- to reduce infection caused by *L. interrogans* serogroup Icterohaemorrhagiae serovars Copenhageni and Icterohaemorrhagiae, *L. interrogans* serogroup Australis serovar Bratislava, *L. kirschneri* serogroup Grippotyphosa serovars Grippotyphosa and Bananal/Liangguang, *L. weilii* serogroup Tarassovi serovar Vughia and *L. borgpetersenii* serogroup Tarassovi serovar Tarassovi

Onset of Immunity:

E. rhusiopathiae: 3 weeks

Porcine parvovirus: 10 weeks

Leptospira serogroups: 2 weeks

Duration of Immunity:

E. rhusiopathiae: 6 months

Porcine parvovirus: 12 months

Leptospira serogroup Australis: 6 months

Leptospira serogroups Canicola, Icterohaemorrhagiae, Grippotyphosa, Pomona and Tarassovi: 12 months

5. CONTRAINDICATIONS

None.

6. ADVERSE REACTIONS

An increase in body temperature may very commonly occur up until two days after vaccination. The observed mean increase was 0.5°C (in individual cases the maximum increase was 1.5°C). Transient local reactions, mostly consisting of red, mild to hard, non-painful swellings are a very common observation. In general, local reactions may have a diameter of ≤ 5 cm, in very rare cases local

reactions in individual animals can be up to 20 cm in diameter. All local reactions disappear completely within approximately 2 weeks after vaccination. In individual animals intermediate systemic reactions, such as vomiting, redness, rapid breathing and twitching, may rarely be observed, which resolve in a few minutes. In individual animals transient reductions in feed intake or activity may uncommonly occur. Feed intake and activity are completely restored within a week.

In post marketing experience:

A hypersensitivity reaction may occur in very rare cases.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

7. TARGET SPECIES

Pig for reproduction

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For intramuscular use. Administer a single dose of 2 ml in the neck region.

Basic vaccination scheme: Pigs which have not yet been vaccinated shall be given a primary injection 6 to 8 weeks before the expected date of insemination and a booster injection 4 weeks later.

Revaccination: A single revaccination with the veterinary medicinal product should be given once a year. Six months post each vaccination with the veterinary medicinal product, a single revaccination with an *Erysipelotrix rhusiopathiae* containing product should be given to maintain immunity against *Erysipelotrix rhusiopathiae*. In case of known infection pressure with *L. interrogans* serogroup Australis, a single revaccination with the veterinary medicinal product should be given every six months, as it is unknown if or for how long the duration of immunity for this serogroup persists beyond six months.

9. ADVICE ON CORRECT ADMINISTRATION

Before use allow the vaccine to reach room temperature.

Shake well before use.

Avoid introduction of contamination by multiple broaching.

10. WITHDRAWAL PERIOD(S)

Zero days.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Store in a refrigerator (2°C – 8°C).

Do not freeze.

Protect from light.

Do not use this veterinary medicinal product after the expiry date which is stated on the label.

Shelf life after first opening the container: 10 hours.

12. SPECIAL WARNING(S)

Special warnings for each target species:

Vaccinate healthy animals only.

Special precautions for use in animals:

Not applicable.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental self-injection, seek medical advice immediately and show the package insert or label to the physician.

Pregnancy and lactation:

Can be used during pregnancy and lactation.

Interaction with other medicinal products and other forms of interactions:

No information is available on the safety and efficacy of this vaccine with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Overdose (symptoms, emergency procedures, antidotes):

No adverse reactions other than those mentioned in section 6 were observed after the administration of a double dose of vaccine.

Incompatibilities:

Do not mix with any other veterinary medicinal product.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

June 2021

15. OTHER INFORMATION

Pack size:

Cardboard box with 1 vial of 20 ml.

Cardboard box with 10 vials of 20 ml.

Cardboard box with 1 vial of 50 ml.

Cardboard box with 10 vials of 50 ml.

Cardboard box with 1 vial of 100 ml.

Cardboard box with 1 vial of 250 ml.

Not all pack sizes may be marketed.

Approved 25 August 2021

