

**MINIMUM PARTICULARS TO APPEAR ON THE LABEL WHERE THERE IS NO  
PACKAGE LEAFLET, E.g. Concertina Labels. {NATURE/TYPE}**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND  
OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR  
BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Alfasan Nederland B.V.  
Kuipersweg 9  
3449 JA Woerden  
The Netherlands

Manufacturer responsible for batch release:

Dales Pharmaceuticals Ltd  
Snaygill Industrial Estate  
Keighley Road  
Skipton  
North Yorkshire  
BD23 2RW  
United Kingdom

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Glucose 40% w/v solution for injection

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS**

1 ml contains:  
Active substances:  
Glucose 40.0% w/v

**4. PHARMACEUTICAL FORM**

Solution for injection.

**5. PACKAGE SIZE**

400 ml

**6. INDICATION**

For the treatment of primary and secondary hypoglycaemia, acetoanaemia and as a  
source of energy in cattle.

As an aid in the treatment of pregnancy toxemia in sheep.

## **7. CONTRAINDICATIONS**

Do not give by subcutaneous injection.

## **8. ADVERSE REACTIONS**

Rapid intravenous injection of dextrose will cause the renal threshold for glucose to be exceeded, with consequent clinical signs of stress and loss of glucose in the urine. For best results intravenous infusions should be given slowly, preferably at a rate not exceeding 0.5 ml/kg/hour.

If you notice any serious effects or other effects not mentioned on this label, please inform your veterinary surgeon.

## **9. TARGET SPECIES**

Cattle and sheep.

## **10. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Cattle: 1 ml/kg, by slow intravenous injection. In some cases, a higher dose rate may be required but should only be given on veterinary advice.  
Treatment may be repeated as necessary.

Sheep: 1 ml/kg to be given by slow intravenous injection. Treatment may be repeated after 6 hours if necessary.

The solution should be warmed to body temperature before use.  
Strict aseptic precautions should be observed in preparation of the site of injection.  
Do not exceed the recommended dosage.

## **11. WITHDRAWAL PERIOD**

Cattle and sheep: Meat: Zero days; Milk: Zero hours

## **12. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C.

Protect from light.

This product does not contain an antimicrobial preservative. Any remaining solution in the bottle following withdrawal of the required dose should be discarded.

## **13. SPECIAL WARNINGS**

Special warnings for each target species:

Intravenous injection should be given slowly, preferably not exceeding 0.5 ml/kg/hour.

Do not administer by subcutaneous route.

User warnings:

Care should be taken to avoid accidental self-injection.  
Wash hands after use.

Interaction with other medicinal products and other forms of interaction:

None known (supplementary treatment can be provided as appropriate).

#### **14. EXPIRY DATE**

EXP: month/year

#### **15. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS**

Dispose of empty containers in the farm refuse. Used containers should not be recycled.

#### **16. DATE ON WHICH THE LABEL WAS LAST APPROVED**

XX/XX/XX

#### **17. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only. To be supplied only on a veterinary prescription.

POM- VPS
-------------

Prescription Only Medicine – Veterinarian, Pharmacist, Suitably Qualified Person

#### **18. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

#### **19. MARKETING AUTHORISATION NUMBER**

Vm 36408/4013

#### **20. MANUFACTURER’S BATCH NUMBER**

Lot: {number}

## **21. OTHER INFORMATION**

UK authorised veterinary medicinal product.

Local representative:

Dechra Veterinary Products Limited, Sansaw Business Park, Hadnall, Shrewsbury, Shropshire, SY4 4AS, United Kingdom.