

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

1 kg, 5 kg or 25 kg Bag label

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Denagard 10% w/w Premix for Medicated Feed for Pigs, Chickens, Turkeys and Rabbits

2. STATEMENT OF ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS

100 mg/g tiamulin hydrogen fumarate

Other ingredients:

Gelatin

Calcium carbonate

Soya-bean oil, refined

3. PHARMACEUTICAL FORM

Premix for medicated feeding stuff

4. PACKAGE SIZE

<1 kg> <5 kg> <25 kg>

5. TARGET SPECIES

Pigs

Chickens (broiler, replacement pullet, layer/breeder)

Turkeys (poult (grower) and breeder)

Rabbits

6. INDICATION(S)

Pigs

For the treatment and metaphylaxis, when the disease is present in the group, of swine dysentery caused by *Brachyspira hyodysenteriae* susceptible to tiamulin. The presence of the disease in the group must be established before the product is used.

For the treatment of colitis caused by *Brachyspira pilosicoli*.

For the treatment of ileitis caused by *Lawsonia intracellularis*.

For the treatment of enzootic pneumonia caused by *Mycoplasma hyopneumoniae*.

Chickens

For the treatment and prevention of chronic respiratory disease (CRD) and airsacculitis caused by *Mycoplasma gallisepticum* and *Mycoplasma synoviae*.

Turkeys

For the treatment and prevention of infectious sinusitis and airsacculitis caused by *Mycoplasma gallisepticum*, *Mycoplasma meleagridis* and *Mycoplasma synoviae*.

Rabbits

For the treatment and prevention of epizootic rabbit enterocolitis (ERE).

See Methods and Route(s) of Administration for further details about indications.

7. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Calculations to achieve the correct dose rate and achieve the correct inclusion rate should be based on:

Inclusion rate (ppm) = dose rate (mg/kg bodyweight) x bodyweight (kg) / daily feed intake (kg).

To ensure a correct dosage, bodyweight should be determined as accurately as possible to avoid underdosing.

The intake of medicated feed depends on the clinical condition of the animals. In order to obtain the correct dosage the concentration of tiamulin hydrogen fumarate has to be adjusted accordingly.

Pigs

Treatment and metaphylaxis of swine dysentery caused by *B. hyodysenteriae*, treatment of porcine colonic spirochaetosis (colitis) caused by *B. pilosicoli*

Dosage: 5 - 10 mg tiamulin hydrogen fumarate/kg bodyweight daily administered for 7 to 10 consecutive days. The dosage will normally be achieved by an inclusion level of 100 – 200 ppm tiamulin hydrogen fumarate in the finished feed provided that feed intake is unaffected.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	1.0 – 2.0 kg

Treatment of porcine proliferative enteropathy (ileitis) caused by *L. intracellularis*

Dosage: 7.5 mg tiamulin hydrogen fumarate/kg bodyweight daily administered for 10 to 14 consecutive days. The dosage will normally be achieved by an inclusion level of 150 ppm tiamulin hydrogen fumarate in the finished feed providing that feed intake is unaffected.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	1.5 kg

Treatment of enzootic pneumonia caused by *M. hyopneumoniae*

Dosage: 5.0 – 10.0 mg tiamulin hydrogen fumarate/kg bodyweight daily administered for 7 to 10 consecutive days. The dosage will normally be achieved by an inclusion level of 100 - 200 ppm tiamulin hydrogen fumarate in the finished feed provided that feed intake is unaffected.

Secondary infection by organisms such as *Pasteurella multocida* and *Actinobacillus pleuropneumoniae* may complicate enzootic pneumonia and require specific medication.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	1.0 – 2.0 kg

Chickens (broiler, replacement pullet, laying and breeding hens)

Treatment and prevention of chronic respiratory disease (CRD) caused by *M. gallisepticum* and airsacculitis and infectious synovitis caused by and *M. synoviae*.

Dosage – Treatment and prevention: 25 mg tiamulin hydrogen fumarate/kg bodyweight daily administered for the period of 3 to 5 consecutive days. This is normally achieved by an inclusion level of 250 - 500 ppm tiamulin hydrogen fumarate in finished feed provided that feed intake is unaffected. Inclusion levels in the higher range will in most cases be needed to avoid underdosing. In fast growing birds, e.g. broiler chickens during the first 2 to 4 weeks of life, inclusion levels in the lower range may be sufficient.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	2.5 - 5.0 kg

Turkeys (young poults, breeding turkeys)

Treatment and prevention of infectious sinusitis and airsacculitis caused by *M. gallisepticum*, *M. synoviae* and *M. meleagridis*.

Dosage – Treatment and prevention: 40 mg tiamulin hydrogen fumarate/kg bodyweight daily administered for the period of 3 to 5 consecutive days. This is normally achieved by an inclusion level of 250-500 ppm tiamulin hydrogen fumarate in finished feed provided that feed intake is unaffected. Inclusion levels in the higher range will in most cases be needed to avoid underdosing. In fast growing birds, e.g. poults during the first 2 to 4 weeks of life, inclusion levels in the lower range may be sufficient.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	2.5 – 5.0 kg

Preventive treatment with tiamulin should only be initiated after confirmed infection with *M. gallisepticum*, *M. synoviae* or *M. meleagridis* and then as an aid in the prevention strategy to reduce the clinical signs and mortality from respiratory disease in flocks, where infection in ovum is likely because the disease is known to exist in the parent generation. The prevention strategy should include efforts to eliminate the infection from the parent generation.

Rabbits

Treatment of Epizootic Rabbit Enterocolitis (ERE) and prevention of ERE in farms with clinical signs of ERE in the previous fattening cycle as part of a programme including measures aiming to eradicate or control the infection in the farm.

Dosage: 3 mg tiamulin hydrogen fumarate /kg bodyweight daily. The dosage will normally be achieved by an inclusion level of 40 ppm tiamulin hydrogen fumarate in

the finished feed provided that feed intake is unaffected. Treatment should be administered until 2 to 3 days after clinical signs has resolved. Prevention should be administered during 3 to 4 weeks from the first week after weaning.

Amount of THF (mg/g) per premix formulation	Amount of premix formulation per one tonne of feed
100.0	0.4 kg

8. WITHDRAWAL PERIOD(S)

Pigs

Meat and offal:

Treatment (at 5 to 10 mg/kg bodyweight): Meat and offal: 6 days

Chickens

Meat and offal: 1 day

Eggs: 0 days

Turkeys

Meat and offal: 4 days

Rabbits

Meat and offal: 0 days

9. SPECIAL WARNING(S), IF NECESSARY

Contraindications

Animals should not receive products containing ionophores (monensin, narasin or salinomycin) during or for at least 7 days before or after treatment with tiamulin. Severe growth depression or death may result.

Warnings for target species

If there is reduced feed intake, the inclusion levels in feed may need to be increased to achieve target dosage. Acute cases and severely diseased animals with reduced feed intake should be treated with a product of suitable formulation such as an injectable or water solution.

Special precautions for use in animals

It is sound clinical practice to base treatment on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of target bacteria.

Adverse reactions

On rare occasions erythema or mild oedema of the skin may occur in pigs following the use of tiamulin.

Use during pregnancy, lactation or lay

Can be used in pigs during pregnancy and lactation.

Can be used in laying and breeding chickens and turkeys.

Can be used in rabbits during pregnancy and lactation.

Interactions

Tiamulin has been shown to interact with ionophores such as monensin, salinomycin and narasin and may result in signs indistinguishable from an ionophore toxicosis. Animals should not receive products containing monensin, salinomycin or narasin during or at least 7 days before or after treatment with tiamulin. Severe growth depression, ataxia, paralysis or death may result.

If signs of an interaction do occur, administration of contaminated feed should be stopped immediately. The feed should be removed and replaced with fresh feed not containing the anticoccidials monensin, salinomycin or narasin.

Overdose

Pigs: Single oral doses of 100 mg/kg bodyweight in pigs caused hyperpnoea and abdominal discomfort. At 150 mg/kg, no CNS effects were noted except for sedation. At 55 mg/kg given for 14 days a transient salivation and slight gastric irritation occurred. A minimum lethal dose has not been established in pigs.

Chickens and turkeys: The LD₅ for chickens is 1290 mg/kg and turkeys 840 mg/kg bodyweight.

The clinical signs of acute toxicity in chickens are - vocalization, clonic cramps and lateral recumbency. In turkeys signs of acute toxicity include clonic cramps, lateral or dorsal recumbency, salivation and ptosis.

If signs of intoxication do occur promptly remove the medicated feed, replace with fresh unmedicated feed and apply supportive, symptomatic therapy.

Operator warnings

When mixing the veterinary medicinal product and handling the medicated feed, direct contact with eyes, skin and mucous membranes should be avoided. Personal protective equipment should be worn when mixing the veterinary medicinal product or handling the medicated feed: overalls, impervious gloves and either a disposable half-mask respirator conforming to European Standard EN 149 or a non-disposable respirator conforming to European Standard EN 140, with a filter to European Standard EN 143. Wash contaminated skin.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or label to the physician.

People with known hypersensitivity to tiamulin should administer the product with caution.

READ FULL LABEL BEFORE USE

10. EXPIRY DATE

EXP: <mmm yyyy>

Shelf life after first opening the container: 12 weeks

Shelf life after incorporation into meal or pelleted feed: 8 weeks.

11. SPECIAL STORAGE CONDITIONS

Do not store above 25°C. Protect from direct sunlight.

Do not use this veterinary medicinal product after the expiry date which is stated on the label. The expiry date refers to the last day of that month.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of waste material in accordance with local requirements.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only.

Consideration should be given to official guidance on the incorporation of medicated premixes in final feeds.

POM-V

To be supplied only on a veterinary prescription.

Prohibition of sale, supply and/or use:

The product can be incorporated into pelleted feed at temperatures of 65°C and maximums of 80°C.

For incorporation into dry feed at the registered mill:

A manufacturer who is approved to incorporate directly at any concentration, veterinary medicinal products or premixtures containing such products must be responsible for mixing when incorporation is less than 2 kg per tonne for final feed.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the reach and sight of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

Elanco Europe Ltd.
Form 2, Bartley Way
Bartley Wood Business Park
Hook
RG27 9XA
United Kingdom

Manufacturers responsible for batch release:

Elanco France
26, rue de la Chapelle

68330 Huningue
France

Andres Pinaluba, S.A.
C/Prudenci Bertrana
5 y 10, Pol. Ind. Agro-Reus
Reus,
43206 Tarragona
Spain

16. MARKETING AUTHORISATION NUMBER(S)

Vm 00879/4051

17. MANUFACTURER'S BATCH NUMBER

Lot: <number>

OTHER INFORMATION

Date of last revision of the label:
Available in 1 kg, 5 kg and 25 kg sacks.
Not all pack sizes may be marketed

Approved: 26/03/21

