

## **PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE {250/500 ml label}**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Fly & Lice Spot On Insecticide 1% w/v Cutaneous Solution

### **2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Contains: 1.0% w/v Deltamethrin

### **3. PHARMACEUTICAL FORM**

Cutaneous Solution

### **4. PACKAGE SIZE**

250 ml

500 ml

### **5. TARGET SPECIES**

Cattle, sheep and lambs

### **6. INDICATIONS**

Ready-to-use insecticide for cattle, sheep and lambs.

**On Cattle:** As a ready-to-use topical application for the control of both sucking and biting lice on all ages of cattle including dairy cattle producing milk for human consumption. As an aid in the control of biting and nuisance flies on cattle.

**On Sheep:** For control of ticks, lice and keds and established blowfly strike.

**On Lambs:** For the control of ticks and lice.

### **7. METHOD AND ROUTE OF ADMINISTRATION**

Important: Read before use

**Dose:** Cattle 10 ml / Sheep: Ticks, lice and keds – 5 ml, Established blowfly strike 5 ml; Lambs (under 10 kg or 1 month old): Ticks and lice 2.5 ml.

**Administration:** Remove the screw cap from the dispensing chamber. Remove the foil. Squeeze the bottle to fill the chamber to the required dose. Release the pressure and pour dose on one spot on the mid-line of the back at the shoulders. For established blowfly strike apply directly to the infested area.

## 8. WITHDRAWAL PERIOD

Cattle (meat) - 17 days, Cattle (milk) - zero hours, Sheep (meat) - 35 days. Do not use in sheep producing milk for human consumption.

## 9. SPECIAL WARNINGS, IF NECESSARY

This product has been granted an exemption from the Poisonous Substances in Agriculture Regulations 1984 by the Health and Safety Executive.

### **Protection of operators:**

Wear protective clothing including waterproof apron and boots and impervious gloves when either applying the product or handling recently treated animals. Remove heavily contaminated clothing immediately and wash before use.

Wash splashes from skin immediately with soap and plenty of water. Wash hands and exposed skin after handling this product and before meals. In case of contact with eyes, rinse immediately with plenty of clean, running water and seek medical advice. In case of accidental ingestion, wash out mouth immediately with plenty of water and seek medical advice. Do not smoke, drink or eat while handling the product.

This product contains deltamethrin which may produce tingling, itchiness and blotchy redness on exposed skin. If you feel unwell after working with this product, consult your doctor and show this label.

**Advice to medical practitioners:** Advice on clinical management is available from the National Poisons Information Service.

**Side-effects:** Minor signs of discomfort have been noted in some cattle during the 48 hours after treatment; this is of no long term detriment to the animal.

For external use only.

## 10. EXPIRY DATE

Exp:

## 11. SPECIAL STORAGE CONDITIONS

Do not store above 25°C. Keep bottles in outer carton.

Store in tightly closed original container away from food, drink and animal feeding stuffs.

Do not freeze.

**12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dangerous to fish and other aquatic life. Do not contaminate ponds, waterways or ditches with the product or used container. Dispose of any unused product or empty containers in accordance with guidance from your local waste regulation authority.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

POM-VPS

To be supplied only on veterinary prescription.

For animal treatment only.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of reach and sight of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited  
1st Floor, Birchwood Building  
Springfield Drive  
Leatherhead  
Surrey  
KT22 7LP

**16. MARKETING AUTHORISATION NUMBER**

Vm 42058/5198

**17. MANUFACTURER’S BATCH NUMBER**

Lot:

## **PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE {1 & 2.5 L label}**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Fly & Lice Spot On Insecticide 1% w/v Cutaneous Solution

### **2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Contains: 1.0% w/v Deltamethrin

### **3. PHARMACEUTICAL FORM**

Cutaneous Solution

### **4. PACKAGE SIZE**

1 L

2.5 L

### **5. TARGET SPECIES**

Cattle, sheep and lambs

### **6. INDICATIONS**

Ready-to-use insecticide for cattle, sheep and lambs.

**On Cattle:** As a ready-to-use topical application for the control of both sucking and biting lice on all ages of cattle including dairy cattle producing milk for human consumption. As an aid in the control of biting and nuisance flies on cattle.

**On Sheep:** For control of ticks, lice and keds and established blowfly strike.

**On Lambs:** For the control of ticks and lice.

### **7. METHOD AND ROUTE OF ADMINISTRATION**

Important: Read before use

**Dose:** Cattle: 10 ml / Sheep: Ticks, lice and keds – 5 ml, Established blowfly strike – 5 ml / Lambs (under 10 kg or 1 month old): Ticks and lice – 2.5 ml.

**Administration:** Apply a single dose with the special SPOT ON APPLICATOR GUN on one spot on the mid-line of the back at the shoulders. For blowfly strike on sheep see application directions below.

**Lice on cattle:** One application will generally eradicate all lice. Complete clearance of all lice may take 4-5 weeks during which time lice hatch from the eggs and are

killed. A very few lice may survive on a small minority of animals. Therefore it may be necessary on rare occasions to re-treat an animal 6 to 8 weeks later.

**Flies on cattle:** To control biting and non-biting flies, frequency of treatment will depend on numbers and species of flies present. Where horn flies predominate good control can be expected for 4-8 weeks. Treatment for fly control should not be repeated within four weeks.

**Ticks on sheep:** Application to the mid-point of the shoulders will provide useful control of ticks attaching to animals of all ages, for up to six weeks after treatment.

**Keds and lice on sheep:** Application to the mid-point of the shoulders of sheep in short or long fleece will reduce the incidence of a biting louse or ked infestation over a 4-6 week period after treatment.

**Established blowfly strike on sheep:** Apply directly to the maggot infested area as soon as fly strike is seen. One application will ensure blowfly larvae are killed in a short time. In the case of more advanced strike lesions, clipping out of stained wool before treatment is advisable.

**Lice and ticks on lambs (under 10 kg or 1 month old):** Application to the midpoint of the shoulders of lambs will provide useful control of ticks for up to 6 weeks after treatment, and will reduce the incidence of biting lice over a 4-6 week period after treatment.

**Note:** When using the SPOT ON Applicator, application to the skin can be achieved by inserting the T-bar nozzle into the fleece in line with the back of the animal and using the T-bar to separate the fleece and penetrate down to skin level before squeezing the applicator handle.

## **8. WITHDRAWAL PERIOD**

Cattle (meat) - 17 days, Cattle (milk) - zero hours, Sheep (meat) - 35 days. Do not use in sheep producing milk for human consumption.

## **9. SPECIAL WARNINGS, IF NECESSARY**

This product has been granted an exemption from the Poisonous Substances in Agriculture Regulations 1984 by the Health and Safety Executive.

### **Protection of operators:**

Wear protective clothing including waterproof apron and boots and impervious gloves when either applying the product or handling recently treated animals. Remove heavily contaminated clothing immediately and wash before use.

Wash splashes from skin immediately with soap and plenty of water. Wash hands and exposed skin after handling this product and before meals. In case of contact

with eyes, rinse immediately with plenty of clean, running water and seek medical advice. In case of accidental ingestion, wash out mouth immediately with plenty of water and seek medical advice. Do not smoke, drink or eat while handling the product. This product contains deltamethrin which may produce tingling, itchiness and blotchy redness on exposed skin. If you feel unwell after working with this product, consult your doctor and show this label.

**Advice to medical practitioners:** Advice on clinical management is available from the National Poisons Information Service.

**Side-effects:** Minor signs of discomfort have been noted in some cattle during the 48 hours after treatment; this is of no long term detriment to the animal.

For external use only.

## **10. EXPIRY DATE**

Exp:

## **11. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C. Keep bottles in outer carton.

Store in tightly closed original container away from food, drink and animal feeding stuffs.

Do not freeze.

## **12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dangerous to fish and other aquatic life. Do not contaminate ponds, waterways or ditches with the product or used container. Dispose of any unused product or empty containers in accordance with guidance from your local waste regulation authority.

## **13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

POM-VPS

To be supplied only on veterinary prescription.

For animal treatment only.

## **14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of reach and sight of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited  
1st Floor, Birchwood Building  
Springfield Drive  
Leatherhead  
Surrey  
KT22 7LP

**16. MARKETING AUTHORISATION NUMBER**

Vm 42058/5198

**17. MANUFACTURER'S BATCH NUMBER**

Lot:

## **PARTICULARS TO APPEAR ON THE OUTER PACKAGE {250/500 ml carton}**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Fly & Lice Spot On Insecticide 1% w/v Cutaneous Solution

### **2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Contains: 1.0% w/v Deltamethrin

### **3. PHARMACEUTICAL FORM**

Cutaneous Solution

### **4. PACKAGE SIZE**

250 ml

500 ml

### **5. TARGET SPECIES**

Cattle, sheep and lambs

### **6. INDICATIONS**

Ready-to-use insecticide for cattle, sheep and lambs.

**On Cattle:** As a ready-to-use topical application for the control of both sucking and biting lice on all ages of cattle including dairy cattle producing milk for human consumption. As an aid in the control of biting and nuisance flies on cattle.

**On Sheep:** For control of ticks, lice and keds and established blowfly strike.

**On Lambs:** For the control of ticks and lice.

### **7. METHOD AND ROUTE OF ADMINISTRATION**

Important: Read before use

**Dose:** Cattle: 10 ml / Sheep: 5 ml / Lambs (under 10 kg or 1 month old): 2.5 ml.

**Administration:** Apply a single dose on one spot on the mid-line of the back at the shoulders. For blowfly strike on sheep see application directions below.

**Lice on cattle:** One application will generally eradicate all lice. Complete clearance of all lice may take 4-5 weeks during which time lice hatch from the eggs and are killed. A very few lice may survive on a small minority of animals. Therefore it may be necessary on rare occasions to re-treat an animal 6 to 8 weeks later.

**Flies on cattle:** To control biting and non-biting flies, frequency of treatment will depend on numbers and species of flies present. Where horn flies predominate good control can be expected for 4-8 weeks. Treatment for fly control should not be repeated within four weeks.

**Ticks on sheep:** Application to the mid-point of the shoulders will provide useful control of ticks attaching to animals of all ages, for up to six weeks after treatment.

**Keds and lice on sheep:** Application to the mid-point of the shoulders of sheep in short or long fleece will reduce the incidence of a biting louse or ked infestation over a 4-6 week period after treatment.

**Established blowfly strike on sheep:** Apply directly to the maggot infested area as soon as fly strike is seen. One application will ensure blowfly larvae are killed in a short time. In the case of more advanced strike lesions, clipping out of stained wool before treatment is advisable.

**Lice and ticks on lambs (under 10 kg or 1 month old):** Application to the midpoint of the shoulders of lambs will provide useful control of ticks for up to 6 weeks after treatment, and will reduce the incidence of biting lice over a 4-6 week period after treatment.

## **8. WITHDRAWAL PERIOD**

Cattle (meat) - 17 days, Cattle (milk) - zero hours, Sheep (meat) - 35 days. Do not use in sheep producing milk for human consumption.

## **9. SPECIAL WARNINGS, IF NECESSARY**

This product has been granted an exemption from the Poisonous Substances in Agriculture Regulations 1984 by the Health and Safety Executive.

### **Protection of operators:**

Wear protective clothing including waterproof apron and boots and impervious gloves when either applying the product or handling recently treated animals. Remove heavily contaminated clothing immediately and wash before use.

Wash splashes from skin immediately with soap and plenty of water. Wash hands and exposed skin after handling this product and before meals. In case of contact with eyes, rinse immediately with plenty of clean, running water and seek medical advice. In case of accidental ingestion, wash out mouth immediately with plenty of water and seek medical advice. Do not smoke, drink or eat while handling the product. This product contains deltamethrin which may produce tingling, itchiness and blotchy redness on exposed skin. If you feel unwell after working with this product, consult your doctor and show this label.

**Advice to medical practitioners:** Advice on clinical management is available from the National Poisons Information Service.

**Side-effects:** Minor signs of discomfort have been noted in some cattle during the 48 hours after treatment; this is of no long term detriment to the animal.

Application site reactions, including squamosis and pruritus have been very rarely seen in cattle during the 48 hours after treatment.

The frequency of adverse reactions is defined using the following convention:

Very rare (less than 1 animal in 10,000 animals treated, including isolated reports)

For external use only.

## **10. EXPIRY DATE**

Exp:

## **11. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C. Keep bottles in outer carton.

Store in tightly closed original container away from food, drink and animal feeding stuffs.

Do not freeze.

## **12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dangerous to fish and other aquatic life. Do not contaminate ponds, waterways or ditches with the product or used container. Dispose of any unused product or empty containers in accordance with guidance from your local waste regulation authority.

## **13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

POM-VPS

To be supplied only on veterinary prescription.

For animal treatment only.

## **14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of reach and sight of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited  
1st Floor, Birchwood Building  
Springfield Drive  
Leatherhead  
Surrey  
KT22 7LP

**16. MARKETING AUTHORISATION NUMBER**

Vm 42058/5198

**17. MANUFACTURER'S BATCH NUMBER**

Lot:

## **PARTICULARS TO APPEAR ON THE OUTER PACKAGE {1 & 2.5 L carton}**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Fly & Lice Spot On Insecticide 1% w/v Cutaneous Solution

### **2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Contains: 1.0% w/v Deltamethrin

### **3. PHARMACEUTICAL FORM**

Cutaneous Solution

### **4. PACKAGE SIZE**

1 L

2 x 2.5 L

### **5. TARGET SPECIES**

Cattle, sheep and lambs

### **6. INDICATIONS**

Ready-to-use insecticide for cattle, sheep and lambs.

**On Cattle:** As a ready-to-use topical application for the control of both sucking and biting lice on all ages of cattle including dairy cattle producing milk for human consumption. As an aid in the control of biting and nuisance flies on cattle.

**On Sheep:** For control of ticks, lice and keds and established blowfly strike.

**On Lambs:** For the control of ticks and lice.

### **7. METHOD AND ROUTE OF ADMINISTRATION**

Important: Read before use

**Dose:** Cattle: 10 ml / Sheep: 5 ml / Lambs (under 10 kg or 1 month old): 2.5 ml.

**Administration:** Apply a single dose on one spot on the mid-line of the back at the shoulders. For blowfly strike on sheep see application directions below.

**Lice on cattle:** One application will generally eradicate all lice. Complete clearance of all lice may take 4-5 weeks during which time lice hatch from the eggs and are killed. A very few lice may survive on a small minority of animals. Therefore it may be necessary on rare occasions to re-treat an animal 6 to 8 weeks later.

**Flies on cattle:** To control biting and non-biting flies, frequency of treatment will depend on numbers and species of flies present. Where horn flies predominate good control can be expected for 4-8 weeks. Treatment for fly control should not be repeated within four weeks.

**Ticks on sheep:** Application to the mid-point of the shoulders will provide useful control of ticks attaching to animals of all ages, for up to six weeks after treatment.

**Keds and lice on sheep:** Application to the mid-point of the shoulders of sheep in short or long fleece will reduce the incidence of a biting louse or ked infestation over a 4-6 week period after treatment.

**Established blowfly strike on sheep:** Apply directly to the maggot infested area as soon as fly strike is seen. One application will ensure blowfly larvae are killed in a short time. In the case of more advanced strike lesions, clipping out of stained wool before treatment is advisable.

**Lice and ticks on lambs (under 10 kg or 1 month old):** Application to the midpoint of the shoulders of lambs will provide useful control of ticks for up to 6 weeks after treatment, and will reduce the incidence of biting lice over a 4-6 week period after treatment.

**Note:** When using the SPOT ON Applicator, application to the skin can be achieved by inserting the T-bar nozzle into the fleece in line with the back of the animal and using the T-bar to separate the fleece and penetrate down to skin level before squeezing the applicator handle.

How to use 1 / 2.5 litre pack:

1. Remove the bottle from the pack.
2. Pass the end of the tape provided through one of the holes at the bottom of the bottle and tie. Pass the other end through the other hole and draw through a convenient length for hanging on the operator. Tie in position
3. With bottle upright remove the plain screw cap on the bottle and replace it by the nozzle cap provided.
4. Attach the tube from the Fly & Lice SPOT ON Applicator firmly to the nozzle of the bottle.
5. Hang the bottle in the inverted position on the operator.
6. Carefully prime the applicator, otherwise inaccurate dosing may result.

Part used bottles (packs) may be kept with the nozzle cap replaced by the screw cap.

## 8. WITHDRAWAL PERIOD

Cattle (meat) - 17 days, Cattle (milk) - zero hours, Sheep (meat) - 35 days. Do not use in sheep producing milk for human consumption.

## 9. SPECIAL WARNINGS, IF NECESSARY

This product has been granted an exemption from the Poisonous Substances in Agriculture Regulations 1984 by the Health and Safety Executive.

### **Protection of operators:**

Wear protective clothing including waterproof apron and boots and impervious gloves when either applying the product or handling recently treated animals. Remove heavily contaminated clothing immediately and wash before use.

Wash splashes from skin immediately with soap and plenty of water. Wash hands and exposed skin after handling this product and before meals. In case of contact with eyes, rinse immediately with plenty of clean, running water and seek medical advice. In case of accidental ingestion, wash out mouth immediately with plenty of water and seek medical advice. Do not smoke, drink or eat while handling the product. This product contains deltamethrin which may produce tingling, itchiness and blotchy redness on exposed skin. If you feel unwell after working with this product, consult your doctor and show this label.

**Advice to medical practitioners:** Advice on clinical management is available from the National Poisons Information Service.

**Side-effects:** Minor signs of discomfort have been noted in some cattle during the 48 hours after treatment; this is of no long term detriment to the animal.

For external use only.

## 10. EXPIRY DATE

Exp:

## 11. SPECIAL STORAGE CONDITIONS

Do not store above 25°C. Keep bottles in outer carton.

Store in tightly closed original container away from food, drink and animal feeding stuffs.

Do not freeze.

**12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dangerous to fish and other aquatic life. Do not contaminate ponds, waterways or ditches with the product or used container. Dispose of any unused product or empty containers in accordance with guidance from your local waste regulation authority.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

POM-VPS

To be supplied only on veterinary prescription.

For animal treatment only.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of reach and sight of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited  
1st Floor, Birchwood Building  
Springfield Drive  
Leatherhead  
Surrey  
KT22 7LP

**16. MARKETING AUTHORISATION NUMBER**

Vm 42058/5198

**17. MANUFACTURER’S BATCH NUMBER**

Lot:

*Gavin Hall*

Approved: 22 May 2025