PARTICULARS TO APPEAR ON THE OUTER PACKAGE

CARTON

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Droncit Spot-on 20 mg solution
Tapewormer

2. STATEMENT OF ACTIVE SUBSTANCES

Each 0.5ml pipette contains:
praziquantel 20mg

Excipients
N-methylpyrrolidone 497.8 mg

3. PACKAGE SIZE

4 x 0.5 ml pipettes.

4. TARGET SPECIES

Cats.

5. INDICATIONS

IMPORTANT: USE CORRECTLY TO SAFEGUARD YOUR PET AND YOUR HEALTH. ALWAYS READ THE PACKAGE LEAFLET CAREFULLY BEFORE USE.

Use: For the treatment of tapeworms in cats.

6. ROUTES OF ADMINISTRATION

Spot- on.

Dosage: The minimum dose rate is 8 mg/kg bodyweight, equivalent to 1 tube per 2.5 kg bodyweight.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {mm/yyyy}

9. SPECIAL STORAGE PRECAUTIONS

ALWAYS READ THE PACKAGE LEAFLET CAREFULLY BEFORE USE.

FOR EXTERNAL USE ONLY.

Do NOT administer by mouth.
Do not use this veterinary medicinal product on cats weighing less than 1 kg.
Do not allow recently treated animals to groom each other.
Use during pregnancy and lactation – read the package leaflet.
The safety of the veterinary medicinal product has not been established in cats during pregnancy, lactation or in animals intended for breeding.

Do not use after expiry date.

The solvent in the veterinary medicinal product may stain certain materials including leather, fabrics, plastics and finished surfaces. Allow the application site to dry before permitting contact with such materials.

Keep the container and the leaflet in the outer carton.

10. THE WORDS "READ THE PACKAGE LEAFLET BEFORE USE"

Read the package leaflet before use.
User safety
For full user safety warnings, read the package leaflet.
Pregnant women and women suspected of being pregnant should not administer the veterinary medicinal product.
Do not stroke or groom animals until the area of application is dry (typically around 1 hour after application).
Wash hands thoroughly after use.

11. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

12. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Vetoquinol UK Limited

14. MARKETING AUTHORISATION NUMBERS

Vm 08007/5026

15. BATCH NUMBER

Lot {number}
Vetoquinol logo

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS: UNIT DOSE TUBE LABEL

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Droncit Spot-on

2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

0.5 ml contains:

Praziquantel 20 mg

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

Vetoquinol logo

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS : BLISTER FOIL

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Droncit Spot-on 20 mg Solution

2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

0.5 ml contains:

Praziquantel 20 mg

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

Vetoquinol logo

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Droncit Spot-on 20 mg solution
Tapewormer

For the treatment of tapeworms of cats. To safeguard the health of yourself and your pet, please keep this leaflet and read it before you apply the product to your pet.
Please read this leaflet carefully before you apply the veterinary medicinal product to your pet. It provides a summary of information about this veterinary medicinal product. Please ask your veterinary surgeon if you have any questions. This leaflet applies only to this veterinary medicinal product.

2. Composition

What is Droncit-Spot-on?

This veterinary medicinal product is called Droncit Spot-on and each pack comprises 4 plastic tubes. The active ingredient is praziquantel which kills your pet's tapeworms. Each tube of product contains 0.5 ml of liquid containing 20 mg of praziquantel. The solution also contains inactive ingredients which are needed to allow the drug to be absorbed across the skin and into the cat's body where it can be effective. Each tube contains 497.8 mg of N-methylpyrrolidone.

3. Target species

Cats.

4. Indications for use

What is Droncit Spot-on used for?

Droncit Spot-on is used for the treatment of tapeworms. It is effective against adult and immature forms of the tapeworms known as *Dipylidium caninum* and tapeworms of the *Taenia* species. The product has been specifically designed for administration to cats. This veterinary medicinal product must only be used in cats weighing more than 1 kg bodyweight.

5. Contraindications

Do not use this veterinary medicinal product on animals weighing less than 1 kg.

6. Special warnings

Special warnings

Do not allow recently treated animals to groom each other.

Special precautions for safe use in the target species:

Care should be taken to avoid the contents of the tube coming into contact with the eyes or mouth of the user or recipient animal.

For external use only.

The safety of the veterinary medicinal product has not been established in cats during pregnancy, lactation or in animals intended for breeding. Laboratory studies in rabbits and rats with the excipient N-methyl pyrrolidone have shown evidence of foetotoxic effects. Use only according to the benefit-risk assessment by the responsible veterinarian.

User safety – IMPORTANT: ALWAYS READ THIS SECTION CAREFULLY

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Laboratory studies in rabbits and rats with the excipient N-methyl pyrrolidone have shown evidence of foetotoxic effects. The veterinary medicinal product should not be administered by pregnant women and women suspected of being pregnant. Personal protective equipment consisting of gloves should be worn when handling the veterinary medicinal product by women of childbearing age.

This veterinary medicinal product can be an irritant to the skin and eyes. Care should be taken to prevent contact of the solution with the skin or eyes. In case of accidental spillage onto skin, wash off any skin contamination with soap and water immediately. If accidental contact occurs with the eyes, flush the affected eyes thoroughly with clean fresh water. In case of accidental contact with the eyes or skin, seek medical advice and if irritation persists show the package leaflet or the label to the physician.

Do not stroke or groom animals until area of application is dry (typically around 1 hour after application).

Wash hand thoroughly after use.

Do not eat, drink or smoke during application.

Store away from food, drink or animal feeding stuffs.

If signs of disease persist or appear, consult a veterinary surgeon.

For Animal Treatment Only.

Pregnancy and lactation

The safety of the veterinary medicinal product has not been established in cats during pregnancy, lactation or in animals intended for breeding. Laboratory studies in rabbits and rats with the excipient N-methyl pyrrolidone have shown evidence of foetotoxic effects. Use only according to the benefit-risk assessment by the responsible veterinarian.

Other precautions:

The solvent in the veterinary medicinal product may stain certain materials including leather, fabrics, plastics and finished surfaces. Allow the application site to dry before permitting contact with such materials.

7. Adverse events

Cats:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Application site reaction[1] (e.g. dandruff[1], exudation[1,2], hair loss[1], reddening[1], itching[1], scab[1]) Hypersalivation[3]

1 Transient
2 Mild
3 Due to the bitter taste of the product if the cat licks the application site immediately after treatment. This is not a sign of intoxication and disappears after a short time without treatment.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system at:

Website: https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine

e-mail: adverse.events@vmd.gov.uk

8. Dosage for each species, routes and method of administration

When to administer Droncit Spot-on?

Droncit Spot-on should be administered at monthly intervals or when signs of tapeworm infestation appear. The number of tubes to be used on your cat depends on how much it weighs.

For cats weighing from 1 to 2.5 kg: use 1 tube of Droncit Spot-on.

For cats weighing over 2.5 kg up to 5 kg use 2 tubes of Droncit Spot-on.
For cats weighing more than 5 kg: use 3 tubes of Droncit Spot-on.

9. Advice on correct administration

How to administer Droncit Spot-on?

Warning: Do NOT Administer Droncit Spot-on by Mouth.

Weigh your cat to see how many tubes of Droncit Spot-on he/she will require. Underdosing could result in ineffective use and may favour resistance development.

Take the required number of tubes from the blister pack. Hold one tube in an upright position with the cap at the top. Twist and pull off the cap. Reverse the cap and replace it onto the tube.

Twist reversed cap, this will break the seal across the top of the tube.

On the back of the animal's neck, immediately behind the head, part the coat until the skin is visible.

Place the tip of the opened tube on the skin and squeeze tube firmly several times to empty the entire contents directly onto the skin. When it is necessary to use more than one tube, care must be taken to prevent the solution running off the skin of the animal. In this situation, it is advised that the applications should be made slowly to allow time for absorption of the solution. Only begin application from the second or third tube when the contents of the previous tube has been completely absorbed. It is particularly important to apply the dose to an area where it cannot be licked off. Application to the back of the neck at the base of the skull will minimise the opportunity for the cat to lick the veterinary medicinal product.

10. Withdrawal periods

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.

Store this veterinary medicinal product away from food, drink and animal feeding stuffs.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton after Exp. The expiry date refers to the last day of that month.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.
After you have applied the product, replace the cap on the tube and dispose of empty packaging and any remaining product in the household refuse.

13. Classification of veterinary medicinal products

Veterinary medicinal product not subject to prescription.

14. MARKETING AUTHORISATION NUMBER AND PACK SIZES

Vm 08007/5026.

White polypropylene pipettes with caps in aluminium blisters.
Blister packs in a cardboard box containing 4 unit dose pipettes (0.5 ml in each pipette).

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the ‘Product Information Database’ on www.gov.uk.

16. Contact details

Marketing authorisation and contact details to report suspected adverse reactions:

Vetoquinol UK Limited,
Steadings Barn,
Pury Hill Business Park, Nr. Alderton,
Towcester,
Northamptonshire,
NN12 7LS

Manufacturer responsible for batch release:

VETOQUINOL BIOWET Sp. z o.o.Żwirowa, 14066-400, Gorzów, Wlkp., Poland

KVP Pharma + Veterinär Produkte GmbH,
Projensdorfer Str. 324,
24106 Kiel,
Germany

17. Other information

Further information
Tapeworm infestation is certain to re-occur unless control of intermediate hosts such as fleas, mice etc. is undertaken. Flea infestations can be controlled by the regular use of effective flea control remedies. Mice control - if cats roam and hunt, they will come into contact with, and possibly consume, mice. As the mouse is one of the hosts of a tapeworm known as *Taenia taeniaeformis* re-infestation of the cat with this tapeworm will be impossible to prevent.

AVM-GSL

Vetoquinol logo

Gavin Hall

Approved: 12 June 2026