Reviewed: August 2017 AN: 00873/2017

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE {25 kg label}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Lincocin® Premix 44 g/kg Premix for Medicated Feed

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Each kilogram contains 44 g Lincomycin (as Lincomycin hydrochloride) in a soyabean millfeed carrier

3. PHARMACEUTICAL FORM

Premix for Medicated Feed

4. PACKAGE SIZE

25 kg

5. TARGET SPECIES

Pigs

6. INDICATION(S)

For the control and treatment of swine dysentery and mycoplasmas pneumonia.

7. METHOD AND ROUTE OF ADMINISTRATION

Control of swine dysentery: Feed 44 grams lincomycin per tonne (1000 kg) of complete feed as the sole ration throughout the period of risk.

Treatment of swine dysentery: Feed 110 grams of lincomycin per tonne (1000 kg) of complete feed as the sole ration for three weeks or until clinical signs of disease (watery, mucoid or bloody stools) disappear.

Treatment and control of swine mycoplasmal pneumonia: Feed 220 grams of lincomycin per tonne (1000 kg) of complete feed as the sole ration for three weeks or until clinical signs of disease disappear.

Mixing directions

For the control of swine dysentery: to make complete feed containing 44 grams of lincomycin, add 1 kg of Lincocin Premix per tonne (1000 kg).

For the treatment of swine dysentery: to make complete feed containing 110 grams of lincomycin, add 2.5 kg of Lincocin Premix per tonne (1000 kg).

Reviewed: August 2017 AN: 00873/2017

For the treatment and control of swine mycoplasmal pneumonia: to make complete feed containing 220 grams of lincomycin, add 5 kg of Lincocin Premix per tonne (1000 kg).

To ensure thorough dispersion of the product it should first be mixed with a suitable quantity of feed before incorporation into the final mix.

For inclusion rates less than 2 kg/tonne: Category A manufacturers only. Incorporation into pig feeding stuffs must follow the product licence directions given above or be in accordance with a veterinary written direction.

8. WITHDRAWAL PERIOD

Animals must not be slaughtered for human consumption during treatment.

Withdrawal periods for the different dosing regimes are as shown below.

Swine dysentery (44 g or 110 g lincomycin/tonne feed):

24 hours

Swine mycoplasmal pneumonia (220 g lincomycin/tonne feed):

3 days

9. SPECIAL WARNING(S), IF NECESSARY

Occasionally pigs fed lincomycin may, within the first two days after the onset of treatment, develop soft stools and/or mild swelling of the anus. On rare occasions, some pigs may show skin irritation and mild irritable behaviour. These conditions are usually self-correcting within 5-8 days without discontinuing the lincomycin treatment.

Do not use in animals with known hypersensitivity to lincomycin. Clinical antagonism may exist between lincomycin and erythromycin and concurrent use is therefore not recommended. Do not use in animals other than the target species.

User Warnings:

Persons known to be hypersensitive (allergic) to lincomycin, spectinomycin or soybean millfeed should not handle this product.

Care should be taken not to inhale any dust. The wearing of approved dust masks (either disposable half-mask respirator conforming to European Standard EN 149 or a non-disposable respirator to European Standard EN 140 with a filter to EN 143) and safety glasses is recommended during the handling and mixing of this product. Skin contact should be avoided. Wash hands and any exposed skin with soap and water immediate after use.

10. EXPIRY DATE

Exp:

11. SPECIAL STORAGE CONDITIONS

Do not store above 25°C. Store in a dry place.

Reviewed: August 2017 AN: 00873/2017

The product will remain stable in the finished feed/medicated supplement for 3 months.

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of any unused product, empty containers and unused medicated feed in accordance with guidance from your local waste regulation authority.

13. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE

POM-V

To be supplied only on veterinary prescription.

For animal treatment only.

14. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"

Keep out of reach and sight of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Zoetis UK Limited 5th Floor, 6 St. Andrew Street London EC4A 3AE

16. MARKETING AUTHORISATION NUMBER

Vm 42058/4077

17. MANUFACTURER'S BATCH NUMBER

Lot:

Approved: 09/08/2017