

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE (1000 ml, 500 ml, 200 ml and 50 ml cartons)**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Ivomec Super Injection for Cattle  
(ivermectin and clorsulon)

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Sterile solution for injection containing 1% w/v ivermectin and 10% w/v clorsulon.

**3. PHARMACEUTICAL FORM**

Solution for injection.

**4. PACKAGE SIZE**

1000 ml  
500 ml  
200 ml  
50 ml

**5. TARGET SPECIES**

Cattle.

**6. INDICATIONS**

For the treatment and control\* of gastro-intestinal roundworms, lungworms, adult liver fluke, eyeworms, warbles, mites\* and lice\* of beef and non-lactating dairy cattle.

\*for details see back of carton and package leaflet.

- IVOMEK Super kills adult and immature gastro-intestinal roundworms, including inhibited *Ostertagia ostertagi*.
- IVOMEK Super kills adult and immature lungworms (*Dictyocaulus viviparus*).
- IVOMEK Super controls re-infection with *Haemonchus placei* and *Cooperia* spp. and *Trichostrongylus axei* acquired up to 14 days after treatment; *Ostertagia ostertagi* and *Oesophagostomum radiatum* acquired up to 21 days after treatment and *Dictyocaulus viviparus* acquired up to 28 days after treatment.
- IVOMEK Super kills adult liver fluke (*Fasciola hepatica*).
- IVOMEK Super kills adult eyeworms (*Thelazia* spp.).
- IVOMEK Super kills warble larvae (*Hypoderma bovis* and *H. lineatum*).
- IVOMEK Super kills mange mites (*Sarcoptes scabiei* var. *bovis*, *Psoroptes bovis*).
- IVOMEK Super kills sucking lice (*Linognathus vituli*, *Haematopinus eurysternus* and *Solenopotes capillatus*).

- IVOMEK Super may also be used as an aid in the control of biting lice (*Damalinia bovis*) and the mange mite (*Chorioptes bovis*), but complete elimination may not occur.

IVOMEK Super contains ivermectin and clorsulon. The timing of treatment should be based on epidemiological factors and should be customised for each individual farm. A dosing program should be established by a qualified professional person.

## 7. METHOD AND ROUTE OF ADMINISTRATION

### DOSAGE AND ADMINISTRATION

**Dosage:** 1 ml per 50 kg bodyweight (based on a recommended dosage level of 200 mcg ivermectin and 2 mg clorsulon per kg bodyweight).

For example:

Bodyweight (kg)	Dose Volume (ml)	Doses per Pack			
		1000 ml	500 ml	200 ml	50 ml
Up to 50	1	1000	500	200	50
51-100	2	500	250	100	25
101-150	3	332	166	66	16
151-200	4	250	125	50	12
201-250	5	200	100	40	10
251-300	6	166	83	33	8

Over 300 kg bodyweight, give 1 ml per 50 kg bodyweight.

**Administration:** Inject SUBCUTANEOUSLY ONLY behind the shoulder.

Divide doses greater than 10 ml between two injection sites. A sterile 17 gauge 1/2-inch needle is recommended. When the temperature of the product is below 5°C, difficulty in administration may be encountered due to increased viscosity. Warming the product and injection equipment to about 15°C will greatly increase the ease with which the product can be injected. Different injection sites should be used for other parenteral (injectable) products.

## 8. WITHDRAWAL PERIODS

Cattle (meat) - 66 days.

Do not use in cattle producing milk for human consumption, or in non-lactating dairy cows including pregnant heifers within 60 days of calving.

## 9. SPECIAL WARNINGS, IF NECESSARY

### PRECAUTIONS

Do not use this product intramuscularly or intravenously.

IVOMEK Super is a low-volume product registered for use in cattle. Do not use in other species as severe adverse reactions, including fatalities in dogs, may occur.

Different injection sites should be used for other parenteral products.

Do not smoke, eat or drink while handling the product.

Wash hands after use.

Take care to avoid self injection: the product may cause local irritation and/or pain at the injection site.

This product does not contain an antimicrobial preservative.

Swab the septum before removing each dose.

Use a dry, sterile needle and syringe. When using the 200 ml, 500 ml and 1000 ml pack sizes, use only automatic syringe equipment.

The expiry date shall be not more than 60 months from the date of manufacture.

#### **NOTE TO USER**

Transitory discomfort has been observed in some cattle following subcutaneous administration.

Soft-tissue swelling at the injection site has also been observed. These reactions have disappeared without treatment. Divide doses greater than 10 ml between two injection sites to reduce occasional discomfort or site reaction.

#### **10. EXPIRY DATE**

Expiry Date:

#### **11. SPECIAL STORAGE CONDITIONS**

Protect from light.

Following withdrawal of the first dose, use the product within 6 months.

Discard unused material.

#### **12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

##### **CONTAINER DISPOSAL**

##### **EXTREMELY DANGEROUS TO FISH AND AQUATIC LIFE.**

Do not contaminate surface waters or ditches with product or used containers.

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

#### **13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE [Distribution category]**

For animal treatment only.

POM-VPS

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim Animal Health UK Ltd, Bracknell, RG12 8YS, UK

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 08327/4167

**17. MANUFACTURER’S BATCH NUMBER**

Batch No.:

**18. ADDITIONAL INFORMATION**

IMPORTANT: READ PACKAGE LEAFLET BEFORE USE.

To be supplied only on veterinary prescription.

If broken, do not accept.

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE (1000 ml, 500 ml,  
200 ml and 50 ml labels)**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Ivomec Super Injection for Cattle  
(ivermectin and clorsulon)

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Sterile solution for injection containing 1% w/v ivermectin and 10% w/v clorsulon.

**3. PHARMACEUTICAL FORM**

Solution for injection.

**4. PACKAGE SIZE**

1000 ml  
500 ml  
200 ml  
50 ml

**5. TARGET SPECIES**

Cattle.

**6. INDICATIONS**

For the treatment and control\* of gastro-intestinal roundworms, lungworms, adult liver fluke, eyeworms, warbles, mites\* and lice\* of beef and non-lactating dairy cattle.

\*IMPORTANT: READ PACKAGE LEAFLET BEFORE USE.

**7. METHOD AND ROUTE OF ADMINISTRATION**

**DOSAGE AND ADMINISTRATION**

**Dosage:** 1 ml per 50 kg bodyweight (based on a recommended dosage level of 200 mcg ivermectin and 2 mg clorsulon per kg bodyweight).

**Administration:** Inject SUBCUTANEOUSLY ONLY behind the shoulder. Divide doses greater than 10 ml between two injection sites. Different injection sites should be used for other parenteral products.

## **8. WITHDRAWAL PERIODS**

Cattle (meat) - 66 days.

Do not use in cattle producing milk for human consumption

Do not use in non-lactating dairy cows including pregnant heifers within 60 days of calving.

## **9. SPECIAL WARNINGS, IF NECESSARY**

Read package leaflet before use.

### **PRECAUTIONS**

Do not use this product intramuscularly or intravenously.

IVOMEC Super is a low-volume product registered for use in cattle. Do not use in other species as severe adverse reactions, including fatalities in dogs, may occur.

Do not smoke, eat or drink while handling the product.

Wash hands after use.

Take care to avoid self injection: the product may cause local irritation and/or pain at the injection site.

This product does not contain an antimicrobial preservative.

Swab the septum before removing each dose.

Use a dry, sterile needle and syringe. When using the 200 ml, 500 ml and 1000 ml pack sizes, use only automatic syringe equipment.

## **10. EXPIRY DATE**

Expiry Date:

## **11. SPECIAL STORAGE CONDITIONS**

Protect from light.

Following withdrawal of the first dose, use the product within 6 months.

Discard unused material.

## **12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

### **CONTAINER DISPOSAL**

#### **EXTREMELY DANGEROUS TO FISH AND AQUATIC LIFE.**

Do not contaminate surface waters or ditches with product or used containers.

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE [Distribution category]**

For animal treatment only.

POM-VPS

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim Animal Health UK Ltd, Bracknell, RG12 8YS, UK

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 08327/4167

**17. MANUFACTURER’S BATCH NUMBER**

Batch No.:

**18. ADDITIONAL INFORMATION**

To be supplied only on veterinary prescription.

**PACKAGE LEAFLET FOR: Ivomec Super Injection for Cattle (ivermectin and clorsulon)**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing Authorisation Holder:  
Boehringer Ingelheim Animal Health UK Ltd, Bracknell, RG12 8YS, UK

Batch Release site:  
Boehringer Ingelheim Animal Health France SCS  
4 Chemin du Calquet  
31000 Toulouse  
France

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Ivomec Super Injection for Cattle  
(ivermectin and clorsulon)

**3. STATEMENT OF THE ACTIVE SUBSTANCES AND OTHER INGREDIENTS**

Sterile solution for injection containing 1% w/v ivermectin and 10% w/v clorsulon.

**4. INDICATIONS**

For the treatment and control\* of gastro-intestinal roundworms, lungworms, adult liver fluke, eyeworms, warbles, mites\*\* and lice\*\* of beef and non-lactating dairy cattle.

IVOMEC Super Injection for Cattle is indicated for the effective treatment and control of the following harmful species of gastro-intestinal roundworms, lungworms, liver fluke, eyeworms, warbles, mites and lice.

**Gastro-intestinal roundworms** (Adults and fourth-stage larvae)  
*Ostertagia ostertagi* (including inhibited larval stages), *O. lyrata*, *Haemonchus placei*, *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Cooperia oncophora*, *C. punctata*, *C. pectinata*, *Bunostomum phlebotomum*, *Oesophagostomum radiatum*, *Strongyloides papillosus* (adult), *Nematodirus helvetianus* (adult), *N. spathiger* (adult), and *Trichuris* spp. (adult)

**Lungworms** (Adult and fourth-stage larvae)

*Dictyocaulus viviparus*

**Liver fluke** (Adult)

*Fasciola hepatica*

**Eyeworms (Adult)**

*Thelazia* spp.

**Warbles (parasitic stages)**

*Hypoderma bovis*, *H. lineatum*

**\*\*Mange Mites**

*Psoroptes bovis*, *Sarcoptes scabiei* var. *bovis*

**\*\*Sucking Lice**

*Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*

IVOMEC Super may also be used as an aid in the control of the biting louse *Damalinia bovis* and the mange mite *Chorioptes bovis*, but complete elimination may not occur.

**PROLONGED ACTIVITY**

When cattle have to graze on pasture contaminated with infective larvae of cattle nematodes, treatment with IVOMEC Super at the recommended dose rate controls re-infection with *Haemonchus placei*, *Cooperia* spp. and *Trichostrongylus axei* acquired up to 14 days after treatment; *Ostertagia* spp. and *Oesophagostomum radiatum* acquired up to 21 days after treatment and *Dictyocaulus viviparus* acquired up to 28 days after treatment.

**5. CONTRAINDICATIONS**

Do not use this product intramuscularly or intravenously.

IVOMEC Super is a low-volume product registered for use in cattle. Do not use in other species as severe adverse reactions, including fatalities in dogs, may occur.

**6. ADVERSE REACTIONS**

**NOTE TO USER**

Transitory discomfort has been observed in some cattle following subcutaneous administration. Soft-tissue swelling at the injection site has also been observed. These reactions have disappeared without treatment. Divide doses greater than 10 ml between two injection sites to reduce occasional discomfort or site reaction.

**7. TARGET SPECIES**

Cattle.

## 8. DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION DOSAGE

IVOMEK Super should only be given by subcutaneous injection at the recommended dosage level of 200 mcg ivermectin and 2 mg clorsulon per kg of bodyweight. Each ml contains 10 mg of ivermectin and 100 mg of clorsulon, sufficient to treat 50 kg of bodyweight. Use the following dosage schedule:

Bodyweight kg	Dose Volume ml	Bodyweight kg	Dose Volume ml
Up to 50	1.0	301-350	7.0
51-100	2.0	351-400	8.0
101-150	3.0	401-450	9.0
151-200	4.0	451-500	10.0
201-250	5.0	501-550	11.0
251-300	6.0	551-600	12.0


Divide doses greater than 10 ml between two injection sites.

To ensure administration of a correct dose, bodyweight should be determined as accurately as possible; accuracy of the dosing device should be checked.

### ADMINISTRATION

IVOMEK Super is to be given SUBCUTANEOUSLY ONLY.

Inject under the loose skin behind the shoulder (see illustration).



Use of a 17 gauge, 1/2 inch (15-20 mm) needle is suggested. Injection of animals with wet or dirty hides is not recommended.


If using a hypodermic syringe, use a separate sterile needle to withdraw IVOMEK Super from the pack. When the temperature of the product is below 5°C, difficulty in administration may be encountered due to increased viscosity. Warming the product and injection equipment to about 15°C will greatly increase the ease with which the product can be injected. Different injection sites should be used for other parenteral products.

## 9. ADVICE ON CORRECT ADMINISTRATION

### INSTRUCTIONS FOR USE WITH AUTOMATIC INJECTION EQUIPMENT

- Disinfect all needles and syringes before using by boiling in clean water for 15-20 minutes.

- Boiled needles should be stored in an antiseptic solution before use and changed frequently when injecting cattle.
- The use of a sterile draw-off assembly is recommended with this product. Handle carefully to avoid contamination.
- Connect the plastic tube firmly to the dosing syringe. Use a stepped adaptor if needed.
- Remove cap from bottle and disinfect rubber stopper with alcohol or other suitable chemical disinfectant. Hold bottle upright and fully insert draw-off needle into centre of rubber stopper (see illustration A). Secure firmly with screw-on cap attached to tube.
- Hang bottle comfortably in inverted position from neck (see illustration B), shoulder or belt.
- Gently prime injector. Equipment is now ready for use.
- After use, remove draw-off assembly and flush out entire apparatus with clean water before storing.
- Store partly-used bottle in carton to protect from light. Do not re-use empty bottles.
- If the connecting tube is used a second time, it should also be boiled for 15-20 minutes before use along with the injecting syringe and needles.



## 10. WITHDRAWAL PERIODS

Cattle must not be slaughtered for human consumption during treatment.

Cattle (meat) - 66 days.

Do not use in cattle producing milk for human consumption, or in non-lactating dairy cows including pregnant heifers within 60 days of calving.

## 11. SPECIAL STORAGE PRECAUTIONS

Protect from light.

Following withdrawal of the first dose, use product within 6 months.

Discard unused material.

## **12. SPECIAL WARNINGS**

### **PRECAUTIONS**

Do not smoke, eat or drink while handling the product.

Wash hands after use.

Take care to avoid self injection: the product may cause local irritation and/or pain at the injection site.

For animal treatment only.

Keep out of the reach of children.

### **Resistance Management**

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.
- Underdosing, which may be due to underestimation of bodyweight, misadministration of the product, or lack of calibration of the dosing device.

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used.

Resistance to macrocyclic lactones (which includes ivermectin) has been reported in *Cooperia* spp. in cattle within the EU. Therefore, the use of this product should be based on local (regional, farm) epidemiological information about susceptibility of nematodes and recommendations on how to limit further selection for resistance to anthelmintics.

This product does not contain an antimicrobial preservative.

Swab the septum before removing each dose.

Use a dry sterile needle and syringe.

When using the 200 ml, 500 ml and 1000 ml pack sizes, use only automatic syringe equipment.

## **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

### **CONTAINER DISPOSAL**

#### **EXTREMELY DANGEROUS TO FISH AND AQUATIC LIFE.**

Do not contaminate surface waters or ditches with product or used container.

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

## **14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

## 15. OTHER INFORMATION

Further information is available on request.

POM-VPS

Vm 08327/4167

To be supplied only on veterinary prescription.

### INTRODUCTION

IVOMEK Super Injection for Cattle is an injectable parasiticide.

One low-volume dose effectively kills internal and external parasites that impair the health and productivity of cattle.

IVOMEK Super Injection contains ivermectin and the flukicide, clorsulon. Its convenience, broad-spectrum efficacy, and wide safety margin make it an ideal product for parasite control of cattle.

### PRODUCT DESCRIPTION

IVOMEK Super Injection is a sterile, non-aqueous solution containing 1% w/v ivermectin and 10% w/v clorsulon for parenteral administration.

At the rate of 1 ml per 50 kg bodyweight by subcutaneous injection, this formulation delivers the recommended dosage level of 200 mcg ivermectin and 2 mg clorsulon per kg bodyweight.

### MODE OF ACTION

Ivermectin paralyzes and ultimately kills parasitic nematodes, arachnids and insects, including warbles, by its effect on the nervous systems of these parasites. At therapeutic doses, ivermectin has no adverse effect on cattle since it does not readily penetrate their central nervous systems. Ivermectin belongs to the avermectin class of anthelmintic endectocides. The mode of action exhibited by the avermectins is unique to this class of antiparasitic agents. Clorsulon works by interrupting the metabolic activity of liver fluke and inhibiting enzymes that are essential for their energy production.

### PRODUCT ADVANTAGES

#### Low-volume Injection

IVOMEK Super is highly effective against internal and external parasites at a dose volume of 1 ml per 50 kg bodyweight. It can be administered quickly and easily.

#### Broad Spectrum

IVOMEK Super provides broad-spectrum efficacy against roundworms (including inhibited *Ostertagia*), lungworms, adult liver fluke, eyeworms, warbles, mange mites and lice.

#### Prolonged Activity

IVOMEK Super remains highly effective against newly ingested larvae of important parasites grazed from the pasture for an extended period after treatment. Re-infection with lungworm is controlled for up to 28 days after treatment and important

roundworms (*Ostertagia* spp. and *Cooperia* spp.) for up to 21 days and 14 days respectively.

IVOMEC Super contains ivermectin and clorsulon. The timing of treatment should be based on epidemiological factors and should be customised for each individual farm. A dosing program should be established by a qualified professional person.

### **SAFETY**

Studies have demonstrated a wide safety margin and the recommended use level had no adverse effect on breeding performance.

Approved 06 October 2020

A handwritten signature in black ink, appearing to read "J. Hunter.", is positioned below the approval date. The signature is stylized and includes a period at the end.