

**ANNEX B**  
**LABELLING AND PACKAGE LEAFLET**

## **I. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**Cardboard box single packs and multipacks**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Spizobactin 750,000 IU / 125 mg chewable tablets for dogs  
Spiramycin / Metronidazole



**2. STATEMENT OF ACTIVE SUBSTANCES**

One tablet contains:  
Spiramycin 750,000 IU  
Metronidazole 125 mg

**3. PHARMACEUTICAL FORM**

Chewable tablets

**4. PACKAGE SIZE**

10 tablets  
20 tablets  
30 tablets  
10 x 10 tablets

**5. TARGET SPECIES**

Dogs

**6. INDICATION(S)**

-

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

For oral use  
Read the package leaflet before use.

**8. WITHDRAWAL PERIODS**

**9. SPECIAL WARNINGS, IF NECESSARY**

User warning: Metronidazole may cause severe adverse reactions. **IMPERVIOUS GLOVES SHOULD BE WORN.** See package leaflet for full user warnings.

**10. EXPIRY DATE**

EXP:

Shelf life of divided tablets: 3 days.

**11. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED VETERINARY MEDICINAL PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

**13. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.

To be supplied only on veterinary prescription.

**14. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Le Vet Beheer B.V.  
Wilgenweg 7  
3421 TV Oudewater  
The Netherlands

**16. MARKETING AUTHORISATION NUMBER**

Vm 41821/4051

**17. MANUFACTURER'S BATCH NUMBER**

Lot.

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

Aluminium-PVC/PE/PVDC **blisters**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Spizobactin 750,000 IU / 125 mg tablets  
Spiramycin / Metronidazole



**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Le Vet. Beheer B.V.

**3. EXPIRY DATE**

EXP:

**4. BATCH NUMBER**

Lot:

**5. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only

## **II. PACKAGE LEAFLET**

## PACKAGE LEAFLET

Spizobactin chewable tablets for dogs

### 1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

Name: Le Vet Beheer B.V.  
Address: Wilgenweg 7  
3421 TV Oudewater  
The Netherlands

Manufacturer responsible for batch release:

Name: LelyPharma B.V.  
Address: Zuiveringweg 42  
8243 PZ Lelystad  
The Netherlands

### 2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Spizobactin 750,000 IU / 125 mg chewable tablets for dogs

### 3. STATEMENT OF THE ACTIVE SUBSTANCES AND OTHER INGREDIENTS

1 tablet Spizobactin 750,000 IU / 125 mg contains:

**Active substances:**

Spiramycin	750,000 IU
Metronidazole	125 mg

Light brown with brown spots, round and convex flavored chewable tablet with a cross-shaped break line on one side.

Tablets can be divided into 2 or 4 equal parts.

### 4. INDICATIONS

For the adjunct treatment of mechanical or surgical periodontal therapy in the treatment of multi-bacterial infections of periodontal and related (peri)oral conditions e.g.

Stomatitis (inflammation of the oral mucosa),

Gingivitis (inflammation of the gingiva),

Glossitis (inflammation of the tongue),

Periodontitis (periodontal inflammation)

Tonsillitis (inflammation of the tonsils),

Dental fistula and other fistulas wounds in the oral cavity,

Cheilitis (inflammation of the mucosa of the lips),

and sinusitis (inflammation of the sinuses) -

in dogs caused by microorganisms susceptible to spiramycin / metronidazole, such as Gram-positive bacteria and anaerobes. See also section 12 (special warnings).

## **5. CONTRAINDICATIONS**

Do not use in case of hepatic disorders.

Do not use in case of hypersensitivity to spiramycin, metronidazole or to any of the excipients.

Do not use concomitantly with bactericidal antibiotics

## **6. ADVERSE REACTIONS**

Vomiting has rarely been observed in dogs.

In rare cases hypersensitivity can occur. In cases of hypersensitivity reactions the treatment should be stopped.

In very rare cases disorders of spermatogenesis may occur.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated )
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Dogs



## **8. DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION**

For oral administration.

75 000 IU spiramycin + 12.5 mg metronidazole per kg body weight, in more severe cases 100 000 IU spiramycin + 16.7 mg metronidazole per kg body weight, administered daily for 6 - 10 days depending on the severity of the disease. In severe cases one can start with the higher dose and go back in the course of treatment on to the lower dose.

The daily dose may be given once daily or divided equally for twice daily administration.

The treatment should always be continued for 1-2 days after resolution of symptoms in order to prevent relapses.

To ensure administration of the correct dosage bodyweight should be determined as accurately as possible to avoid underdosing. The following table is intended as a guide to dispensing the product at approximately the standard dose rate of 75 000 IU spiramycin + 12.5 mg metronidazole per kg body weight.




Bodyweight	Spizobactin 750,000 IU / 125 mg for dogs	Spizobactin 1,500,000 IU / 250 mg for dogs	Spizobactin-L 3,000,000 IU / 500 mg for dogs
2.5 kg	◐		
5.0 kg	◑	◐	
7.5 kg	◑		
10 kg	⊕	◑	◐
12.5 kg	⊕ ◐		
15 kg	⊕ ◑	◑	
17.5 kg	⊕ ◑		
20 kg	⊕ ⊕	⊕	◑
25 kg		⊕ ◐	
30 kg		⊕ ◑	◑
35 kg		⊕ ◑	
40 kg		⊕ ⊕	⊕
50 kg			⊕ ◐
60 kg			⊕ ◑
70 kg			⊕ ◑
80 kg			⊕ ⊕

◐ = 1/4 Tablet    ◑ = 1/2 Tablet    ◑ = 3/4 Tablet    ⊕ = 1 Tablet

### 9. ADVICE ON CORRECT ADMINISTRATION

The tablets are to be administered either deep in the mouth (on the base of the tongue) or given with a small amount of food containing the tablet, to ensure all the tablet is consumed.

Tablets can be divided into 2 or 4 equal parts to ensure accurate dosing. Place the tablet on a flat surface, with its scored side facing up and the convex (rounded) side facing the surface.



Halves: press down with your thumbs on both sides of the tablet.

Quarters: press down with your thumb in the middle of the tablet

### 10. WITHDRAWAL PERIODS

Not applicable

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Shelf life of divided tablets after first opening the immediate packaging: 3 days.

Do not store above 30°C.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

## **12. SPECIAL WARNINGS**

In many cases of endodontic/periodontal disease the primary treatment is non-medicinal and does not require antimicrobial medication.

Antimicrobial treatment of periodontal disease should be accompanied or preceded by endodontic therapy and/or professional dental cleaning especially if the disease is advanced. Dog owners are encouraged to routinely brush their dog's teeth to remove plaque to prevent or to control periodontal disease

### **Special precautions for use in animals**

The combination of spiramycin and metronidazole should not be used as first-line empirical treatment. Whenever possible, metronidazole and spiramycin should only be used based on susceptibility testing.

Official, national and regional antimicrobial policies should be taken into account when the veterinary medicinal product is used.

Treatment with this medicinal product must normally not exceed the specified treatment period (6 to 10 days). This time limit may only be exceeded in rare cases with particular strict indications. Repetition of the treatment is also allowed only when having strict indications. Limiting the duration of treatment is necessary because damage to the germ cells cannot be excluded with the use of metronidazole, and because in long-term studies with high doses, an increase of certain tumours was seen in rodents.

### **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

Metronidazole has confirmed mutagenic and genotoxic properties in laboratory animals as well as in humans. Metronidazole is a confirmed carcinogen in laboratory animals and has possible carcinogenic effects in humans. However, there is inadequate evidence in humans for the carcinogenicity of metronidazole.

Spiramycin may in rare cases induce hypersensitivity reactions, e.g. contact dermatitis.

Direct contact with the skin or mucous membranes of the user should be avoided because of the risk of sensitization. Do not handle the product if you are known to be hypersensitive to the active substances or to any of the excipients. **IMPERVIOUS GLOVES SHOULD BE WORN** during administration of the product to avoid skin contact with the product.

To avoid accidental ingestion, particularly by a child, unused part-tablets should be returned to the open blister space and inserted back into the carton.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Wash hands thoroughly after handling the tablets.

### **Use during pregnancy and lactation**

Spiramycin has not been found to be teratogenic or embryo- or foetotoxic. Studies in laboratory animals have shown inconsistent results with regard to teratogenic/embryotoxic effects of metronidazole. Therefore, use of this product during pregnancy is not recommended. Metronidazole and spiramycin are excreted in milk and use during lactation is therefore not recommended.

### **Interactions with other medicinal products and other forms of interaction**

Macrolides, such as e.g. spiramycin act antagonistic to penicillins and cephalosporins.

The product should not be used concurrently with other antibiotics of the macrolide group.

Metronidazole may have an inhibitory effect on the degradation of other drugs in the liver, such as phenytoin, cyclosporine and warfarin.

Phenobarbital may increase hepatic metabolism of metronidazole resulting in decreased serum concentration of metronidazole.

### **Overdose (symptoms, emergency procedures, antidotes)**

Adverse events are more likely to occur at doses and treatment durations in excess of the recommended treatment regimen. If neurological signs occur, treatment should be discontinued and the patient should be treated symptomatically.

### **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED VETERINARY MEDICINAL PRODUCTS OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements..

### **14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

### **15. OTHER INFORMATION**


Aluminium - PVC/PE/PVDC blister

Cardboard box of 1, 2 or 3 blisters of 10 tablets

Cardboard box containing 10 separate cardboard boxes, each containing 1 blister of 10 tablets.

Not all pack sizes may be marketed.

Divisible tablet



**Approved: 12/09/2017**

