

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE {Folding carton}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Synulox Lactating Cow Intramammary suspension

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each 3 g intramammary syringe contains: Amoxicillin (as amoxicillin trihydrate)

200 mg

Clavulanic acid (as potassium clavulanate) 50 mg

Prednisolone 10 mg

**3. PACKAGE SIZE**

3 syringes

12 syringes

24 syringes

300 syringes

**4. TARGET SPECIES**

Cattle (lactating cows)

**5. INDICATION(S)**

**6. ROUTES OF ADMINISTRATION**

Intramammary use.

**7. WITHDRAWAL PERIODS**

Withdrawal period:

Meat and offal: 7 days.

Milk: 84 hours, i.e. 7 milking times with 2 times a day milking, or 11 milking times with 3 times a day milking.

**8. EXPIRY DATE**

Exp. {mm/yyyy}

**9. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C. Store in a dry place.

Keep the syringe in the outer carton.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited  
1st Floor, Birchwood Building  
Springfield Drive  
Leatherhead  
Surrey  
KT22 7LP

**14. MARKETING AUTHORISATION NUMBERS**

Vm 42058/5105

**15. BATCH NUMBER**

Lot {number}

**16. SPECIAL WARNING(S), IF NECESSARY**

User warnings:

Penicillins and cephalosporins may occasionally cause severe allergic reactions.  
See package leaflet for user warnings.

**17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

**18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF  
APPLICABLE**

**POM-V**

To be supplied only on veterinary prescription.

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING**  
**UNITS {Syringe label}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Synulox Lactating Cow

**2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCE(S)**

Each syringe contains:

Amoxicillin	200 mg
Clavulanic acid	50 mg
Prednisolone	10 mg

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/yyyy}

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

3 g

**6. ROUTE(S) OF ADMINISTRATION**

For intramammary use.

**7. WITHDRAWAL PERIOD**

Meat and offal: 7 days. Milk: 84 hours.

**8. THE WORDS "FOR ANIMAL TREATMENT ONLY"**

For animal treatment only.

Keep out of the sight and reach of children.

Keep the syringe stored in the outer carton.

POM-V

To be supplied only on veterinary prescription.

Vm 42058/5105

Zoetis UK Limited

1st Floor, Birchwood Building

Springfield Drive

Leatherhead

Surrey  
KT22 7LP

**PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Synulox Lactating Cow Intramammary suspension

**2. COMPOSITION**

Each 3 g intramammary syringe contains:

**Active ingredients:**

Amoxicillin (as amoxicillin trihydrate): 200 mg

Clavulanic acid (as potassium clavulanate): 50 mg

Prednisolone: 10 mg

Pale cream/buff coloured oily intramammary suspension.

**3. TARGET SPECIES**

Cattle (lactating cows).

**4. INDICATIONS FOR USE**

For use in clinical cases of mastitis including cases associated with infections with the following pathogens:

Staphylococci (including  $\beta$ -lactamase producing strains)

Streptococci (including *S. agalactiae*, *S. dysgalactiae* and *S. uberis*)

Escherichia coli (including  $\beta$ -lactamase producing strains)

**5. CONTRAINDICATIONS**

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in animals which are known to be hypersensitive to  $\beta$ -lactamase antibiotics.

**6. SPECIAL WARNING(S)**

Special warnings:

Do not use in cases associated with *Pseudomonas*.

Special precautions for safe use in the target species:

Swab teat end with appropriate disinfectant before treatment.

**Recommendations for prudent use**

The product should be used for treatment of clinical mastitis only.

Use of the product should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria and take into account official and local antimicrobial policies.

The use of the product should preferably be based on susceptibility tests.

Avoid use of the product in herds where no  $\beta$ -lactamase producing *Staphylococci* strains have been isolated. Veterinarians should strive to use narrow spectrum antibiotics if possible.

Inappropriate use of the product may increase the prevalence of bacteria resistant to  $\beta$ -lactam antibiotics and may decrease the effectiveness of treatment with  $\beta$ -lactam antibiotics, due to the potential for cross-resistance.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact.

Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa.

Allergic reactions to these substances may occasionally be serious.

Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations.

Handle this product with great care to avoid exposure, taking all recommended precautions.

If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing, are more serious symptoms and require urgent medical attention.

Wash hands after use.

#### Pregnancy and lactation:

No special precautions.

#### Interaction with other medicinal products and other forms of interaction:

None known.

## **7. ADVERSE EVENTS**

None known.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system *{national system details}*.

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

The contents of one syringe should be infused into each affected quarter via the teat canal, immediately after milking, at 12 hour intervals for three consecutive milkings. In cases of infections caused by *Staphylococcus aureus*, a longer course of antibacterial therapy may be required. Therefore overall treatment length must be at the veterinarian's discretion but should be long enough to ensure complete resolution of intramammary infection.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Before the infusion is made, the teat end should be cleaned and disinfected.

## **10. WITHDRAWAL PERIOD(S)**

Meat and offal: 7 days.

Milk: 84 hours, i.e. 7 milking times with 2 times a day milking or 11 milking times with 3 times a day milking.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Do not store above 25°C.

Store in a dry place.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp. The expiry date refers to the last day of that month.

## **12. SPECIAL PRECAUTIONS FOR DISPOSAL**

Medicines should not be disposed of via wastewater or household waste.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements. Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help protect the environment.

## **13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.

## **14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES**

Vm 42058/5105



Pack sizes:

Carton containing 3, 12, 24 or 300 intramammary syringes. Not all pack sizes may be marketed.

## **15. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST REVISED**

April 2023

Detailed information on this veterinary medicinal product is available in the Union Product Database.

## **16. CONTACT DETAILS**


### **Marketing authorisation holder:**

Zoetis UK Limited  
1st Floor, Birchwood Building  
Springfield Drive  
Leatherhead  
Surrey  
KT22 7LP

### **Manufacturer responsible for batch release:**

Haupt Pharma Latina S.r.l.  
Strada Statale 156 Dei Monti Lepini Km 47600  
Latina  
04100  
Italy

## **17. OTHER INFORMATION**

A handwritten signature in black ink, consisting of several loops and a final horizontal stroke.

Approved: 27 February 2024