Reviewed: January 2018

MINIMUM PARTICULARS TO APPEAR ON THE LABEL WHERE THERE IS NO PACKAGE LEAFLET, E.g. Concertina Labels. {NATURE/TYPE}

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Bob Martin (UK) Ltd,

Wemberham Lane, Yatton,

North Somerset, BS49 4BS. UK.

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Dog Flea Collar with dimpylate 15% w/w

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS

15% of dimpylate. This collar contains 3.6 g of dimpylate

4. PHARMACEUTICAL FORM

Collar

5. PACKAGE SIZE

6. INDICATION(S)

For the control of fleas and ticks on dogs over 3 months of age. Provides up to 4 months protection.

7. CONTRAINDICATIONS

DO NOT allow animals to chew collar. DO NOT use any other insecticide while the collar is being worn or within 7 days of removal. DO NOT use on puppies under 3 months of age or on nursing bitches. Remove collar immediately should any signs of skin irritation or adverse reaction occur. REMOVE COLLAR FROM DOG IMMEDIATELY SHOULD ANY SIGNS OF DROWSINESS, LISTLESSNESS OR RESPIRATORY DIFFICULTIES DEVELOP SOON AFTER FIXING OF COLLAR.

8. ADVERSE REACTIONS

9. TARGET SPECIES

Dogs

10. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

RECOMMENDED DOSE AND DOSING SCHEDULE: Remove collar from sachet and fasten around dog's neck, adjusting until a comfortable fit is achieved. In the fully grown dog correct fitting will allow two fingers to be inserted comfortably between the collar and neck. Loosen the collar as the dog grows bigger. Any excess length of collar extending beyond the buckle should be cut off and disposed of in the domestic refuse. This collar does not need to be removed when the animal is being bathed as insecticidal activity remains unaffected by water. Leave in place until the collar's effective life is exhausted. Replace every 4 months if re-infestation with fleas or ticks is a risk.

11. ADVICE ON CORRECT ADMINISTRATION

This product is a VETERINARY MEDICINE. Please read the information provided carefully before fitting the collar. This sachet should be retained until the collar is thrown away

12. WITHDRAWAL PERIOD

N/A

13. SPECIAL STORAGE PRECAUTIONS

Do not remove collar from pack until ready to use. Do not store above 25°C. Protect from direct sunlight.

14. SPECIAL WARNING(S)

OPERATOR WARNINGS: Do not remove collar from pack until ready to use. DO NOT ALLOW CHILDREN TO PLAY WITH THE COLLAR. Do not allow children to suck or chew the collar. Do not smoke, eat or drink while handling collar. Do not sleep with pets wearing collars. DIMPYLATE / DIAZINON is an organophosphorus compound: DO NOT handle if under medical advice not to work with ANTICHOLINESTERASES. Avoid excessive handling of the collar. Cut off any excess collar with scissors and dispose of safely. IF YOU FEEL UNWELL AFTER HANDLING THE COLLAR, SEEK MEDICAL ADVICE. WASH HANDS THOROUGHLY AFTER HANDLING THE COLLAR.

15. EXPIRY DATE

16. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

EXTREMELY DANGEROUS to fish and aquatic life. Do not contaminate ponds, waterways or ditches with the collar or empty packaging. Dispose of empty packaging, used and unused collar with the household refuse.

17. DATE ON WHICH THE LABEL WAS LAST APPROVED

Reviewed: January 2018

18. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

[Distribution category]

AVM-GSL

For animal treatment only

19. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"

Keep out of the reach of children

20. MARKETING AUTHORISATION NUMBER(S)

Vm 00715/4112

21. MANUFACTURER'S BATCH NUMBER

22. OTHER INFORMATION

Approved: 04/01/2018