

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**Cardboard for 2 x 10 tablets, 5 x 10 tablets, 10 x 10 tablets, 50 x 10 tablets and 100 x 10 tablets**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Ronaxan 20 mg tablets for dogs and cats  
doxycycline hyclate

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each tablet contains:  
Doxycycline (as doxycycline hyclate) .....20 mg

**3. PHARMACEUTICAL FORM**

Tablets

**4. PACKAGE SIZE**

2 x 10 tablets  
5 x 10 tablets  
10 x 10 tablets  
50 x 10 tablets  
100 x 10 tablets

**5. TARGET SPECIES**

Dogs and cats

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use.  
Read the package leaflet before use.

**8. WITHDRAWAL PERIOD(S)**

**9. SPECIAL WARNING(S), IF NECESSARY**

**10. EXPIRY DATE**

EXP {month/year}

**11. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C.  
Keep the blister in the outer carton.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim Animal Health UK Ltd  
Ellesfield Avenue  
Bracknell  
Berkshire  
RG12 8YS  
United Kingdom.

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 08327/4069

**17. MANUFACTURER'S BATCH NUMBER**

Lot {number}


**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**Blister of 10 tablets**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Ronaxan 20 mg tablets for dogs and cats  
doxycycline hyclate

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**



**3. EXPIRY DATE**

EXP {month/year}

**4. BATCH NUMBER**

Lot {number}

**5. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

## **B. PACKAGE LEAFLET**

**PACKAGE LEAFLET:  
Ronaxan 20 mg tablets for dogs and cats**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER  
AND OF THE MANUFACTURING AUTHORISATION HOLDER  
RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:  
Boehringer Ingelheim Animal Health UK Ltd  
Ellesfield Avenue  
Bracknell  
Berkshire  
RG12 8YS  
United Kingdom

Manufacturer responsible for batch release:  
Boehringer Ingelheim Animal Health France SCS  
4 Chemin du Calquet  
31000 Toulouse  
France

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Ronaxan 20 mg tablets for dogs and cats  
doxycycline hyclate

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER  
INGREDIENTS**

Each tablet contains:

**Active substance:**  
Doxycycline (as doxycycline hyclate) .....20 mg

Light yellow to yellow, biconvex, round, scored tablets.  
The tablets can be divided into two equal parts.

**4. INDICATION(S)**

Dogs

For the treatment of respiratory tract infections including rhinitis, tonsillitis and bronchopneumonia caused by *Bordetella bronchiseptica* and *Pasteurella* spp. susceptible to doxycycline.

For the treatment of canine ehrlichiosis (a disease transmitted by ticks) caused by *Ehrlichia canis*.



## Cats

For the treatment of respiratory tract infections including rhinitis, tonsillitis and bronchopneumonia caused by *Bordetella bronchiseptica* and *Pasteurella* spp. susceptible to doxycycline.

## **5. CONTRAINDICATIONS**

Do not use in cases of hypersensitivity to the active substance or any of the excipients.

Do not use in animals with renal or hepatic insufficiency.

Do not use in animals with diseases associated with vomiting or dysphagia (difficulty to swallow) (see also section "Adverse reactions").

Do not use in animals with known photosensitivity (see also section "Adverse reactions").

Do not use in puppies and kittens before completion of teeth enamel formation.

## **6. ADVERSE REACTIONS**

Gastrointestinal adverse reactions including vomiting, nausea (signs the animal may be sick), salivation (drooling), oesophagitis (irritation of the oesophagus) and diarrhoea have been reported very rarely in spontaneous reports.

Photosensitivity and photodermatitis (irritation of the skin) can occur following tetracycline therapy, after exposure to intense sunlight or ultraviolet light. (See also section "Contraindications").

Use of tetracycline during the period of tooth development may lead to tooth discolouration.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Dogs and cats.

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

For oral use.

The dosage is 10 mg doxycycline per kg bodyweight per day corresponding to one tablet per 2 kg bodyweight. The dosage can be divided into two daily administrations. The duration of treatment might be adapted depending on the clinical response, after benefit/risk assessment by the veterinarian.

Disease	Dosage regimen	Duration of treatment
Respiratory tract infection	10 mg/kg per day	5-10 days
Canine ehrlichiosis	10 mg/kg per day	28 days

## 9. ADVICE ON CORRECT ADMINISTRATION

To ensure a correct dosage, bodyweight of the animals should be determined as accurately as possible to avoid overdosing or underdosing. In order to adjust the dosage, the tablets can be divided into two equal parts. Tablets should be administered with food in order to avoid vomiting.

## 10. WITHDRAWAL PERIOD(S)

Not applicable.

## 11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Do not store above 25°C.

Keep the blister in the outer carton.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

## 12. SPECIAL WARNING(S)

### Special warnings for each target species

#### For the veterinarian

Ehrlichia canis infection: treatment should be initiated at the onset of clinical signs. Complete eradication of the pathogen is not always achieved, but treatment for 28 days generally leads to a resolution of the clinical signs and a reduction of the bacterial load. A longer duration of treatment, based on a benefit/risk assessment by the responsible veterinarian, may be required particularly in severe or chronic ehrlichiosis. All treated patients should be regularly monitored, even after clinical cure.

### Special precautions for use in animals

Tablets should be administered with food in order to avoid vomiting and to reduce the likelihood of oesophageal irritation.

The product should be administered with caution to young animals, since tetracyclines as a class may cause permanent discolouration of the teeth, when administered during tooth development. However, human literature indicates that doxycycline is less likely than other tetracyclines to cause these abnormalities, due to its reduced ability to chelate calcium.

#### For the veterinarian

Use of the veterinary medicinal product should be based on identification and susceptibility testing of the target pathogens. If this is not possible, therapy should be based on epidemiological information and knowledge of susceptibility of the target pathogens at local/regional level.

Use of the veterinary medicinal product deviating from the instructions given in the leaflet may increase the prevalence of bacteria resistant to doxycycline and may decrease the effectiveness of treatment with other tetracyclines, due to the potential for cross resistance.

Use of the veterinary medicinal product should be in accordance with official, national and regional antimicrobial policies.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals

People with known hypersensitivity to doxycycline or other tetracyclines should avoid contact with the veterinary medicinal product and personal protective equipment consisting of gloves should be worn when handling the veterinary medicinal product. In case of skin irritation, seek medical advice immediately and show the package leaflet or the label to the physician.

Accidental ingestion, especially by children, may cause adverse reactions such as emesis. To avoid accidental ingestion, blisters should be inserted back into the outer packaging and kept in a safe place. In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

#### Pregnancy and lactation

Laboratory studies in rats and rabbits have not produced any evidence of teratogenic or embryotoxic effects (malformations or deformities of the embryo) of doxycycline. However, as there is no information available in the target species, use is not recommended during pregnancy.

Use only according to the benefit/risk assessment by the responsible veterinarian.

#### Interaction with other medicinal products and other forms of interaction

Doxycycline should not be used concurrently with other antibiotics especially bactericidal drugs such as the  $\beta$ -lactams (for example penicillin, ampicillin). Cross-resistance to tetracyclines may occur.

The half-life of doxycycline is reduced by concurrent administration of barbiturates (some types of sedatives or tranquilisers), phenytoin and carbamazepine (two types of anti-epileptic medications).

Dosage adjustments may be necessary in subjects under anticoagulant therapy (blood thinners), as tetracyclines depress the plasma activity of prothrombin. Simultaneous administration of oral absorbents, antacids (protectants for the stomach) and preparations including multivalent cations should be avoided as they reduce doxycycline availability.

Overdose (symptoms, emergency procedures, antidotes)

Vomiting may occur in dogs with 5 times the recommended dose. Increased levels of ALT, GGT, ALP and total bilirubin were reported in dogs at 5-fold overdose.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

January 2022

**15. OTHER INFORMATION**

Pack sizes:

2 x 10 tablets, 5 x 10 tablets, 10 x 10 tablets, 50 x 10 tablets or 100 x 10 tablets.  
Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

Approved 27 January 2022

A handwritten signature in black ink, appearing to read "Hunter.", is positioned below the approval date.