

**PARTICULARS TO APPEAR ON <THE OUTER PACKAGE> <AND> <THE IMMEDIATE PACKAGE> {NATURE/TYPE}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Appertex 2.5 mg Tablets

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Clazuril 2.5 mg

**3. PHARMACEUTICAL FORM**

Tablets

**4. PACKAGE SIZE**

**5. TARGET SPECIES**

Homing and show pigeons

**6. INDICATION(S)**

For the oral treatment and control of Coccidiosis caused by *Eimeria labbeana* & *Eimeria columbarum* in homing and show pigeons

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

One tablet per pigeon regardless of weight. To be placed into mouth.

If signs of disease persist or appear, consult a veterinary surgeon.

**8. WITHDRAWAL PERIOD**

Not for use in pigeons intended for human consumption.

**9. SPECIAL WARNING(S), IF NECESSARY**

Do not administer with drugs that may cause vomiting. Birds should be treated when mated and again 3-4 weeks later to prevent infection of nestlings. Birds should be treated following transportation to shows or races where they may be exposed to contaminated cages, baskets etc. Wash hands after administration of the tablets to birds and after cleaning the waste from treated birds.

If signs of disease persist or appear, consult a veterinary surgeon.

For use in pigeons only.

**10. EXPIRY DATE**

## **11. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C.

Store in a dry place.

Keep the container in the outer carton.

## **12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

See pack leaflet

## **13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE [Distribution category]**

For animal treatment only

|             |
|-------------|
| AVM-<br>GSL |
|-------------|

## **14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

KEEP OUT OF THE REACH AND SIGHT OF CHILDREN

## **15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Harkers Limited  
Unit 2, Cavendish Road  
Bury St Edmunds  
Suffolk  
IP33 3TE

## **16. MARKETING AUTHORISATION NUMBER(S)**

**Vm 11245/4000**

## **17. MANUFACTURER’S BATCH NUMBER**

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**  
**{NATURE/TYPE}**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Appertex 2.5 mg Tablets

Clazuril 2.5 mg

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Harkers

**3. EXPIRY DATE**

**4. BATCH NUMBER**

**5. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only

Keep container in outer carton

**PACKAGE LEAFLET FOR:**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Harkers Limited  
Unit 2, Cavendish Road  
Bury St Edmunds  
Suffolk  
IP33 3TE

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Appertex 2.5mg Tablets

**3. STATEMENT OF THE ACTIVE SUBSTANCE (S) AND OTHER INGREDIENTS**

Presentation: round white tablets containing 2.5 mg clazuril

**4. INDICATION(S)**

For the oral treatment and control of Coccidiosis caused by *Eimeria labbeana* & *Eimeria columbarum* in homing and show pigeons

**5. CONTRAINDICATIONS**

Do not administer with drugs that may cause vomiting.

**6. ADVERSE REACTIONS**

**7. TARGET SPECIES**

Homing and show pigeons

**8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

One tablet per pigeon regardless of weight. Administer into mouth. All birds in the same loft should be treated simultaneously to prevent re-infection.

**9. ADVICE ON CORRECT ADMINISTRATION**

**10. WITHDRAWAL PERIOD(S)**

Not for use in pigeons intended for human consumption.

## **11. SPECIAL STORAGE PRECAUTIONS**

Do not store above 25°C.

Store in a dry place.

Keep the container in the outer carton.

Keep out of the reach and sight of children

## **12. SPECIAL WARNING(S)**

Birds should be treated when mated and again 3-4 weeks later to prevent infection of nestlings. Birds should be treated following transportation to shows or races where they may be exposed to contaminated cages, baskets etc. Wash hands after administration of the tablets to birds and after cleaning the waste from treated birds.

For Animal Treatment Only

## **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Dispose of empty packaging and any remaining product in the household refuse. Excreta from treated birds should not be spread onto land used for growing crops.

## **14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

## **15. OTHER INFORMATION**

AVM - GSL

*Vm 11245/4000*

Appertex can safely be administered to pigeons during the breeding season, it has no adverse effects on the condition of the nestlings. In order to reduce the risk of re-infection following treatment, good management and hygiene practices are essential, in particular the regular cleaning of the premises to remove pigeon excreta.

If signs of disease persist or appear, consult a veterinary surgeon.

**Approved: 08/01/2018**

