

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Box with 5 vials containing 4 ml & Box with 5 vials containing 20 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Paracox-5 suspension for oral suspension

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Per dose of 0.004 ml:

<i>Eimeria acervulina</i> HP	500 – 650 oocysts
<i>Eimeria maxima</i> CP	200 – 260 oocysts
<i>Eimeria maxima</i> MFP	100 – 130 oocysts
<i>Eimeria mitis</i> HP	1000 – 1300 oocysts
<i>Eimeria tenella</i> HP	500 – 650 oocysts

3. PACKAGE SIZE

5 x 4 ml (1 000 doses)
5 x 20 ml (5 000 doses)

4. TARGET SPECIES

Chickens.

5. INDICATION(S)

6. ROUTES OF ADMINISTRATION

Oral use.
Solvent can be used when sprayed on chickens.

7. WITHDRAWAL PERIODS

Withdrawal period: Zero days.

+

8. EXPIRY DATE

Exp.: {dd/mm/yyyy}
Once broached/diluted use immediately.

9. SPECIAL STORAGE PRECAUTIONS

Store and transport refrigerated.
Do not freeze.
Protect from light.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

MSD Animal Health UK Ltd.

14. MARKETING AUTHORISATION NUMBERS

Vm 01708/5101

15. BATCH NUMBER

Lot {number}

16. SPECIAL WARNING(S), IF NECESSARY

17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: Read package leaflet.

18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

POM-VPS

Veterinary medicinal product subject to prescription

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

Label vials 4 ml & 20 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Paracox-5



2. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

Per dose of 0.004 ml:

<i>E. acervulina</i> HP	500 – 650 oocysts
<i>E. maxima</i> CP	200 – 260 oocysts
<i>E. maxima</i> MFP	100 – 130 oocysts
<i>E. mitis</i> HP	1000 – 1300 oocysts
<i>E. tenella</i> HP	500 – 650 oocysts

4 ml (1 000 doses)

20 ml (5 000 doses)

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp.: {dd/mm/yyyy}

Once broached/diluted use immediately.

5. ROUTE(S) OF ADMINISTRATION

Oral use.

6. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

PARTICULARS TO APPEAR ON THE IMMEDIATE LABEL OF THE SOLVENT-

Label Bottles 100 ml & 500 ml

1. NAME OF THE SOLVENT

Solvent for spray-on-chickens

Bottle of 100 ml

Bottle of 500 ml

2. CONTENT BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

100 ml

500 ml

3. ROUTES OF ADMINISTRATION

Read the Paracox-5/-8 package leaflet before use.

4. STORAGE CONDITIONS

Store between 2 – 25 °C.

5. BATCH NUMBER

Lot {number}

6. EXPIRY DATE

Exp.: {mm/yyyy}

7. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

8. NAME OF THE MARKETING AUTHORISATION HOLDER

MSD Animal Health UK Ltd.

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Paracox-5- suspension for oral suspension for chickens

2. COMPOSITION

Each 0.004 ml dose of vaccine contains:

Active substances:

Sporulated oocysts derived from precocious lines of coccidia:

Active substances:

<i>Eimeria acervulina</i> , strain HP, live	500 – 650 oocysts*
<i>Eimeria maxima</i> , strain CP, live	200 – 260 oocysts*
<i>Eimeria maxima</i> , strain MFP, live	100 – 130 oocysts*
<i>Eimeria mitis</i> , strain HP, live	1000 – 1300 oocysts*
<i>Eimeria tenella</i> , strain HP, live	500 – 650 oocysts*

* According to the *in vitro* counting procedure of the manufacturer at the time of blending and at release.

Suspension: milky suspension after mixing.

Solvent for spray-on-chickens: semi-opaque, red, viscous solution.

3. TARGET SPECIES

Chickens.

4. INDICATIONS FOR USE

Spray-on-feed, spray-on-chicken without solvent or in drinking water

For the active immunisation of chickens to reduce infection and clinical signs of coccidiosis caused by *Eimeria acervulina*, *E. maxima*, *E. mitis* and *E. tenella*.

Onset of immunity: 14 days after vaccination.

Duration of immunity: 40 days after vaccination.

Spray-on-chickens with solvent

For the active immunisation of chickens against coccidiosis caused by *Eimeria acervulina*, *E. maxima*, *E. mitis* and *E. tenella*:

- to reduce oocyst excretion for *E. acervulina*, *E. maxima*, and *E. tenella*.
- to reduce loss in weight gain for *E. acervulina*, *E. mitis* and *E. tenella*.

Onset of immunity: 21 days after vaccination.

Duration of immunity: 10 weeks after vaccination.

5. CONTRAINDICATIONS

None.

6. SPECIAL WARNINGS

Special warnings:

Vaccinate healthy animals only.

Special precautions for safe use in the target species:

Do not administer to stressed chicks, e.g. chilled, not feeding or drinking.

For administration by spray-on-chickens a red food colouring agent should be added to the diluted vaccine, or the vaccine should be diluted using “Solvent for spray-on-chickens”. For the spray-on-chicken method of administration a significant reduction in efficacy may be observed if diluted in tap water only.

Chickens should be strictly floor reared on litter. The vaccine contains live coccidia and is dependent upon replication of the vaccinal lines within the host for development of protection.

It is common to find oocysts in the gastrointestinal tract of vaccinated birds from 1 – 3 weeks or more after vaccination. These oocysts are most likely to be vaccinal oocysts which recycle in the birds via the litter. Recycling ensures satisfactory flock protection against all the pathogenic species of *Eimeria* contained in the vaccine.

Measures should be taken to ensure that the bulk diluted vaccine is resuspended at intervals during administration.

Since the protection against coccidial infection following the vaccine administration is enhanced by natural challenge, it should be noted that access to any therapeutic agents having anti-coccidial activity at any time following vaccination may reduce the duration of effective protection. This is important throughout the life of the chicken.

To reduce the chance of coccidial field challenge before the onset of immunity, litter should be removed and chicken housing should be thoroughly cleaned between rearing cycles.

Ensure that all vaccination equipment is thoroughly cleaned before use.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Personal protective equipment consisting of well-fitting masks and eye protection should be worn when spraying the vaccine.

Laying birds:

Do not use in birds in lay.

Interaction with other medicinal products and other forms of interactions:

Do not administer anticoccidial agents including sulphonamides and antibacterial agents before or after vaccination with the veterinary medicinal product .

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Overdose:

Severe overdose (5-fold or more) may lead to a temporary reduction in daily live-weight gain.

Major incompatibilities:

Do not mix with any other veterinary medicinal product except the solvent recommended for use for spray administration.

7. ADVERSE EVENTS

Chickens:

Common (1 to 10 animals / 100 animals treated):	Intestinal lesion ¹ .
-------------------------------------------------------	----------------------------------

¹ Mild intestinal lesions of e.g. *E. Acervulina* and *E. tenella* (lesion scores of +1 or +2 using the numerical ranking system of Johnson and Reid, 1970), 3 to 4 weeks after vaccination. Lesions of this severity will not affect the performance of immune chickens.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in

the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system:

E-mail: adverse.events@vmd.gov.uk

Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

8. DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION

Dosage

A single dose of vaccine is administered to chickens from one day of age via spray on feed, via spray on chickens, or at 3 days old via drinking water. The nominal dose is 0.004 ml per chicken. Each 4 ml container will provide sufficient vaccine for 1 000 chickens and each 20 ml container will provide sufficient vaccine for 5 000 chickens.

Administration

Chickens should be floor reared on litter.

On feed

Sufficient starter feed for the first 24 - 48 hours should be laid out on paper or plastic along the floor of the poultry house. Shake the container vigorously for 30 seconds before use, to ensure re-suspension of the oocysts. Dilute the vaccine in water at the rate of approximately 5000 doses in up to 3 litres of water and spray evenly over the surface of the feed using a coarse spray. Ensure a controlled, even coverage of the total surface area of the feed available to the chicks.

Agitate the applicator reservoir regularly throughout application to avoid settling out of oocysts. Once the vaccine has been diluted for use it should be sprayed onto feed and birds should be placed with access to feed within two hours. When the treated allocation of feed has been consumed, routine feeding may continue.

In drinking water

Place chicks in the house at day-old and encourage them to become accustomed to the nipple drinker system. When the chicks are 3 days old the lighting system is turned off for about 7 hours. Raise all drinking lines out of reach of the chicks for about two hours before administration of the vaccine. At the same time the lighting is switched on. Drain each drinking line completely.

Dilute the vaccine to a concentration of 1 dose/2 – 4 ml in cold tap water. Calculate the average number of birds per drinking line and calculate the volume of diluted vaccine needed per drinking line at a rate 2 – 4 ml per bird.

Fill each line with diluted vaccine and lower to allow the birds access to the nipples. An initial charge (about 1 litre) of an indicator (e.g. milk) can be used to show when the line has been filled to the end and can be closed, without wasting

vaccine. As the birds drink, keep each line full via its reservoir until all the diluted vaccine prepared for that line has been added. Normal water supply then follows.

By spray-on-chickens

For administration by spray on chickens, red food colouring agent Cochineal E120 should be added to the diluted vaccine, or the vaccine should be diluted using the recommended solvent "Solvent for spray-on chickens" The solvent contains red colouring agent and xanthan gum, both of which are included for better uptake.

a) Solvent for spray-on-chickens

Vaccine should be delivered using a dose volume of between 0.21 and 0.28 ml diluted vaccine per bird using a coarse spray. Determine the delivery capacity of the spray device in terms of the volume delivered per 100 birds. Multiply this volume by 50 to give the total volume of diluted vaccine required for 5000 doses (or by 10 for 1000 doses). I.e. for the preparation of 5000 doses diluted vaccine, a total of $0.21 \times 5\,000 = 1\,050$ ml diluted vaccine is needed and is divided over the vaccine, solvent and water as below:

1. 20 ml vaccine (1 vial)
2. 500 ml Solvent (1 bottle)
3. Fill up to 1 050 ml with tap water

Water used for vaccine dilution should be fresh, cool and free of pollution. Take a clean container for vaccine preparation, add the solvent to the container, and add the calculated amount of water to the container and mix solvent and water to a uniform solution. Shake the 5 000 dose (or 1 000 dose) vial of Paracox-5 vigorously for 30 seconds to ensure re-suspension of the oocysts. Add the entire contents of the vial into the container with solvent and water and mix thoroughly. Add the diluted vaccine to the applicator reservoir and spray evenly over the birds using a coarse spray. Ensure a controlled, even coverage of the total internal surface area of the box containing the chickens. Leave the birds in the box for at least 30 minutes in a well-lighted area to allow time for the birds to preen.

b) Red food colouring agent (E120)

Vaccine should be delivered using a dose volume of between 0.21 and 0.28 ml diluted vaccine per bird using coarse spray. Determine the delivery capacity of the spray device in terms of the volume delivered per 100 birds. Multiply this volume by 50 to give the total volume of diluted vaccine required for 5 000 doses (or by 10 for 1 000 doses) and add this volume of water to a suitable container (normally between 1.0 and 1.5 litres for 5 000 doses or 200 and 300 ml for 1,000 doses). Uptake of the vaccine by the birds, and therefore the efficacy of the vaccine, is improved if a red food colouring agent is added to the diluted vaccine before administration by spray. Add sufficient red food colouring agent (cochineal E120) to the water to give a concentration of 0.1% w/v.

Shake one 5000 dose (or 1 000 dose) vial of the vaccine vigorously for 30 seconds to ensure resuspension of the oocysts. Add the entire contents of the

vial to the solvent and mix thoroughly. Add the diluted vaccine to the applicator reservoir and operate the cabinet to spray evenly over the birds using a coarse spray.

Ensure a controlled, even coverage of the total internal surface area of the box containing the chicks. Agitate the applicator reservoir regularly throughout application to avoid settling out of oocysts. Leave the birds in the box for at least 30 minutes in a well-lit area to allow time for the birds to preen.

9. ADVICE ON CORRECT ADMINISTRATION

On feed

Ensure that all available feed is treated and that the total number of doses dispensed matches the number of birds in the house. Do not administer the vaccine via an automatic feeding machine or place treated feed directly under heating lamps.

In drinking water

It is recommended that before using the vaccine in a facility for the first time, precautions are taken to check that the procedure ensures the drinking lines have been properly primed with the vaccine, as shown by the appearance of the indicator from nipples at the end the line, before the chicks are allowed to start drinking.

By spray-on-chickens

For administration by spray on chickens a red food colouring agent should be added to the diluted vaccine or the vaccine should be diluted using “Solvent for spray-on chickens”, supplied by the manufacturer. For the spray-on-chicken method of administration a significant reduction in efficacy may be observed if diluted in tap water only. The purity of Cochineal E120 should be in compliance with Commission Directive 95/45/EC.

10. WITHDRAWAL PERIODS

Zero days.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Vaccine

Store and transport refrigerated (2 °C – 8 °C).

Do not freeze.

Protect from light.

Shelf life after dilution according to directions: use immediately.

Solvent for spray-on-chickens

Store between 2 °C – 25 °C.

Do not use this veterinary medicinal product after the expiry date, which is stated on the label/ box after Exp.

12. SPECIAL PRECAUTIONS FOR DISPOSAL

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 01708/5101

Pack sizes:

Vaccine

Box with 5 vials containing 4 ml (1 000 doses)

Box with 5 vials containing 20 ml (5 000 doses)

Solvent

100 ml container for 1000 doses

500 ml container for 5000 doses

The appropriate volume of solvent is supplied together with the vaccine (100 ml solvent for 1 000 doses, 500 ml for 5 000 doses).

Not all pack sizes may be marketed.

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. CONTACT DETAILS

Marketing authorisation holder and contact details to report suspected adverse reactions:

MSD Animal Health UK Ltd.
Walton Manor, Walton
Milton Keynes
MK7 7AJ
Tel.: +44 (0)1908 685685

Manufacturer responsible for batch release¹:

- (i) MSD Animal Health UK Ltd.
Walton Manor, Walton
Milton Keynes
MK7 7AJ
United Kingdom

- (ii) Merck Sharp & Dohme Animal Health S.L.
Poligono Industrial El Montalvo I
C/Zepelin 6, Parcela 38,
37008 Carbajosa de La Sagrada (Salamanca)
Spain

¹ The printed package leaflet will state the name and address of the manufacturer responsible for the release of the concerned batch only.

17. OTHER INFORMATION

In any animal population there may be a small number of individuals which fail to respond fully to vaccination. Successful vaccination depends upon correct storage and administration of the vaccine together with the animal's ability to respond. This can be influenced by such factors as genetic constitution, intercurrent infection, age, nutritional status, concurrent drug therapy and stress.

POM-VPS Veterinary medicinal product subject to prescription

For animal treatment only.

Approved 11 December 2024
Gavin Hall