

ANNEX III
LABELLING AND PACKAGE LEAFLET

1. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Carton of 1, 2, 3, 4, 6, 9, 12, 21 and 42 pipette(s) – 2.5 ml pipettes

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Moxiclear 250 mg + 62.5 mg spot-on solution for large dogs
Imidacloprid, moxidectin

2. STATEMENT OF ACTIVE SUBSTANCES

Each 2.5 ml pipette contains:

Active substances:

| | |
|--------------|---------|
| Imidacloprid | 250 mg |
| Moxidectin | 62.5 mg |

3. PHARMACEUTICAL FORM

Spot-On Solution.

4. PACKAGE SIZE

1 x 2.5 ml
2 x 2.5 ml
3 x 2.5 ml
4 x 2.5 ml
6 x 2.5 ml
9 x 2.5 ml
12 x 2.5 ml
21 x 2.5 ml
42 x 2.5 ml

5. TARGET SPECIES

For large dogs weighing between 10 kg and 25 kg.

6. INDICATION(S)

Indication to be included only for medicinal products not subject to medical prescription.

For dogs suffering from, or at risk from, mixed parasitic infections:

- For the treatment and prevention of flea infestation (*Ctenocephalides felis*),
 - the treatment of biting lice (*Trichodectes canis*),
 - the treatment of ear mite infestation (*Otodectes cynotis*), sarcoptic mange (caused by *Sarcoptes scabiei* var. *canis*),
 - the prevention of heartworm disease (L3 and L4 larvae of *Dirofilaria immitis*),
 - the treatment of circulating microfilariae (*Dirofilaria immitis*),
 - the treatment of cutaneous dirofilariosis (adult stages of *Dirofilaria repens*),
 - the prevention of cutaneous dirofilariosis (L3 larvae of *Dirofilaria repens*),
 - the reduction of circulating microfilariae (*Dirofilaria repens*),
 - the prevention of angiostrongylosis (L4 larvae and immature adults of *Angiostrongylus vasorum*),
 - the treatment of *Angiostrongylus vasorum* and *Crenosoma vulpis*,
 - the prevention of spirocercosis (*Spirocerca lupi*),
 - the treatment of infections with gastrointestinal nematodes (L4 larvae, immature adults and adults of *Toxocara canis*, *Ancylostoma caninum* and *Uncinaria stenocephala*, adults of *Toxascaris leonina* and *Trichuris vulpis*).
- The product can be used as part of a treatment strategy for flea allergy dermatitis (FAD).

7. METHOD AND ROUTE(S) OF ADMINISTRATION

For external use only.

Spot-on use. Do not remove the pipette from the sachet until ready to use.

Read the package leaflet before use.

8. WITHDRAWAL PERIOD

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

10. EXPIRY DATE

EXP {month/year}

11. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

Store in the original package in order to protect from light and moisture.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: Read the package leaflet

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable

For animal treatment only.
To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

(EU)
Norbrook Laboratories (Ireland) Limited
Rossmore Industrial Estate
Monaghan
Ireland

(UK)
Norbrook Laboratories Ltd
Station Works
Newry
Co. Down
BT35 6JP

16. MARKETING AUTHORISATION NUMBER

Vm 02000/4439

17. MANUFACTURER’S BATCH NUMBER

BN



Flea



Flea larvae



Whipworm



Hookworm



Roundworm



Angiostrongylus



Crenosoma



Heartworm



Otodectes



Sarcoptes



Louse



Microfilaria



Spirocerca



Skinworm

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

Sachet for 2.5 ml pipette

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Moxiclear 250 mg + 62.5 mg spot-on solution for large dogs
Imidacloprid, moxidectin

2. QUANTITY OF ACTIVE SUBSTANCE(S)

One 2.5 ml pipette contains:
Imidacloprid 250 mg
Moxidectin 62.5 mg

3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

2.5 ml

4. ROUTE(S) OF ADMINISTRATION

For external use only.
Spot-on use
<Pictogram of a spot-on pipette>

5. WITHDRAWAL PERIOD

6. BATCH NUMBER

Lot

7. EXPIRY DATE

EXP {mm/yyyy}

8. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.
<Pictogram of a dog>

9. MARKETING AUTHORISATION HOLDER

Norbrook Laboratories Limited



MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

2.5 ml pipette

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Moxiclear

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Norbrook

3. EXPIRY DATE

EXP {mm/yyyy}

4. BATCH NUMBER

XXXX XXX

5. THE WORDS "FOR ANIMAL TREATMENT ONLY"

<Pictogram of a dog>

6. PHARMACEUTICAL FORM

<Pictogram of a spot-on pipette>

7. VOLUME

2.5 ml

8. TARGET ANIMAL WEIGHT RANGE

>10 ≤ 25 kg

9. QUANTITY OF ACTIVE SUBSTANCE(S)

250mg/62.5mg



PACKAGE LEAFLET

PACKAGE LEAFLET:

Moxiclear 250 mg + 62.5 mg spot-on solution for large dogs

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder

(EU)

Norbrook Laboratories (Ireland) Limited
Rossmore Industrial Estate
Monaghan
Ireland

(UK)

Norbrook Laboratories Limited,
Station Works,
Newry,
Co. Down,
BT35 6JP,
United Kingdom

Manufacturer responsible for batch release:

Norbrook Manufacturing Ltd.
Rossmore Industrial Estate
Monaghan
Ireland

Norbrook Laboratories Limited,
Station Works,
Newry,
Co. Down,
BT35 6JP,
United Kingdom

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Moxiclear 250 mg + 62.5 mg spot-on solution for large dogs
Imidacloprid, moxidectin

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS

Each 2.5 ml pipette contains:

Active substances:

| | |
|--------------|---------|
| Imidacloprid | 250 mg |
| Moxidectin | 62.5 mg |

Excipients:

| | |
|-----------------------------|-----------|
| Butylhydroxytoluene (E 321) | 2.5 mg |
| Benzyl Alcohol | to 2.5 ml |

A clear colourless to yellow solution with light opalescence.

4. INDICATIONS

For dogs suffering from, or at risk from, mixed parasitic infections:

- For the treatment and prevention of flea infestation (*Ctenocephalides felis*),
 - the treatment of biting lice (*Trichodectes canis*),
 - the treatment of ear mite infestation (*Otodectes cynotis*), sarcoptic mange (caused by *Sarcoptes scabiei* var. *canis*), ,
 - the prevention of heartworm disease (L3 and L4 larvae of *Dirofilaria immitis*),
 - the treatment of circulating microfilariae (*Dirofilaria immitis*),
 - the treatment of cutaneous dirofilariosis (skinworm) (adult stages of *Dirofilaria repens*)
 - the prevention of cutaneous dirofilariosis (skinworm) (L3 larvae of *Dirofilaria repens*),
 - the reduction of circulating microfilariae (*Dirofilaria repens*),
 - the prevention of angiostrongylosis (L4 larvae and immature adults of *Angiostrongylus vasorum*),
 - the treatment of *Angiostrongylus vasorum* and *Crenosoma vulpis*,
 - the prevention of spirocercosis (*Spirocerca lupi*),
 - the treatment of infections with gastrointestinal nematodes (L4 larvae, immature adults and adults of *Toxocara canis* (round worm), *Ancylostoma caninum* (hookworm) and *Uncinaria stenocephala* (hookworm), adults of *Toxascaris leonina* (roundworm) and *Trichuris vulpis* (whipworm)).
- The product can be used as part of a treatment strategy for flea allergy dermatitis (FAD).

5. CONTRAINDICATIONS

Do not use in puppies under 7 weeks of age.

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

Do not use in dogs classified as Class 4 for heartworm disease as the safety of the product has not been evaluated in this animal group.

For cats, the corresponding “Moxiclear for cat” product, which contains 100 mg/ml imidacloprid and 10 mg/ml moxidectin, must be used.

For ferrets: Do not use Moxiclear for dogs. Only “Moxiclear for small cats and ferrets” (0.4 ml) must be used.

Do not use on canaries.

6. ADVERSE REACTIONS

Use of the product may result in transient pruritus in dogs. On rare occasions greasy hair, erythema and vomiting can occur. These signs disappear without further treatment. The product may, in rare cases, cause local hypersensitivity reactions. If the animal licks the application site after treatment, neurological signs (most of which

are transient) such as ataxia, generalized tremors, ocular signs (dilated pupils, little pupillary reflex, nystagmus), abnormal respiration, salivation and vomiting may be observed in very rare cases.

The product tastes bitter. Salivation may occasionally occur if the animal licks the application site immediately after treatment. This is not a sign of intoxication and disappears within some minutes without treatment. Correct application will minimise licking of the application sites.

The product may in very rare cases cause at the application site a sensation resulting in transient behavioural changes such as lethargy, agitation, and inappetence.

A field study has shown that in heartworm positive dogs with microfilaraemia there is a risk of severe respiratory signs (coughing, tachypnea and dyspnea) that may require prompt veterinary treatment. In the study these reactions were common (seen in 2 of 106 treated dogs). Gastrointestinal signs (vomiting, diarrhoea, inappetence) and lethargy are also common adverse reactions following treatment in such dogs.

In case of accidental oral uptake, symptomatic treatment should be administered. There is no known specific antidote. The use of activated charcoal may be beneficial.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

7. TARGET SPECIES

Dogs.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Spot-on use.

For external use only.

Dosage schedule:

The recommended minimum doses are 10 mg/kg bodyweight imidacloprid and 2.5 mg/kg bodyweight moxidectin, equivalent to 0.1 ml/kg bodyweight of product.

The treatment schedule should be based on individual veterinary diagnosis and on the local epidemiological situation.

| Weight of dog | Pipette size to be used | Volume | Imidacloprid [mg/kg bw] | Moxidectin [mg/kg bw] |
|---------------|---|--------|-------------------------|-----------------------|
| ≤ 4 kg | Moxiclear for small dogs | 0.4 ml | minimum of 10 | minimum of 2.5 |
| > 4 ≤ 10 kg | Moxiclear for medium dogs | 1.0 ml | 10–25 | 2.5–6.25 |
| > 10 ≤ 25 kg | Moxiclear for large dogs | 2.5 ml | 10–25 | 2.5–6.25 |
| > 25 ≤ 40 kg | Moxiclear for very-large dogs | 4.0 ml | 10–16 | 2.5–4 |
| > 40 kg | the appropriate combination of pipettes | | | |

Flea treatment and prevention (*Ctenocephalides felis*)

One treatment prevents future flea infestation for 4 weeks. Existing pupae in the environment may emerge for 6 weeks or longer after treatment is initiated, depending upon climatic conditions. Therefore, it may be necessary to combine the product treatment with environmental treatments aimed at breaking the flea life cycle in the surroundings. This can result in a more rapid reduction in the household flea population. The product should be administered at monthly intervals when used as part of a treatment strategy for flea allergy dermatitis.

Treatment of biting lice (*Trichodectes canis*)

A single dose should be administered. A further veterinary examination 30 days after treatment is recommended as some animals may require a second treatment.

Treatment of ear mite infestation (*Otodectes cynotis*)

A single dose of the product should be administered. Loose debris should be gently removed from the external ear canal at each treatment. A further veterinary examination 30 days after treatment is recommended as some animals may require a second treatment. Do not apply directly to the ear canal.

Treatment of sarcoptic mange (caused by *Sarcoptes scabiei* var. *canis*)

A single dose should be administered twice 4 weeks apart.

Prevention of heartworm disease (*D. immitis*) and cutaneous dirofilariosis (skin worm)(*D. repens*)

Dogs in areas endemic for heartworm, or those which have travelled to endemic areas, may be infected with adult heartworms. Therefore prior to treatment with the product, the advice provided in the section “SPECIAL WARNING(S)” should be considered.

For prevention of heartworm disease and cutaneous dirofilariosis, the product must be applied at regular monthly intervals during the time of the year when mosquitoes

(the intermediate hosts which carry and transmit *D. immitis* and *D. repens* larvae) are present. The product may be administered throughout the year or at least 1 month before the first expected exposure to mosquitoes. Treatment should continue at regular monthly intervals until 1 month after the last exposure to mosquitoes. To establish a treatment routine, it is recommended that the same day or date be used each month. When replacing another heartworm preventative product in a heartworm

prevention programme, the first treatment with the product must be given within 1 month of the last dose of the former medication.

In non-endemic areas there should be no risk of dogs having heartworm. Therefore they can be treated without special precautions.

Treatment of microfilariae (*D. immitis*)

The product should be administered monthly for two consecutive months.

Treatment of cutaneous dirofilariosis (skin worm) (adult stages of *Dirofilaria repens*)

The product should be administered monthly for six consecutive months.

Reduction of microfilariae (skin worm) (*D. repens*)

The product should be administered monthly for four consecutive months. Efficacy against adult worms has not been shown. Adult worms may continue to produce microfilariae.

Treatment and prevention of *Angiostrongylus vasorum*

A single dose should be administered. A further veterinary examination 30 days after treatment is recommended as some animals may require a second treatment. In endemic areas regular 4 weekly application will prevent angiostrongylosis and patent infection with *Angiostrongylus vasorum*.

Treatment of *Crenosoma vulpis*

A single dose should be administered.

Prevention of spirocercosis (*Spirocerca lupi*):

The product should be administered monthly.


Roundworm, hookworm and whipworm treatment (*Toxocara canis*, *Ancylostoma caninum*, *Uncinaria stenocephala*, *Toxascaris leonina* and *Trichuris vulpis*).

In areas endemic for heartworm, monthly treatment may significantly reduce the risk of re-infection caused by the respective round-, hook- and whipworms. In areas non-endemic for heartworm, the product can be used as part of a seasonal prevention programme against fleas and gastrointestinal nematodes.

Studies have shown that monthly treatment of dogs will prevent infections caused by *Uncinaria stenocephala*.

9. ADVICE ON CORRECT ADMINISTRATION

For external use only. Do not remove the pipette from the sachet until ready to use. Remove the pipette from the outer sachet using scissors or fold along diagonal line to expose nick; tear back at nick.



Hold the pipette upright. Tap the narrow part of the pipette to ensure the contents remain within the main body of the pipette. Twist or snap back the tip.




For dogs up to 25 kg:

With the dog in a standing position, part the coat between the shoulder blades until the skin is visible. Wherever possible apply to undamaged skin. Place the tip of the pipette on the skin and squeeze the pipette firmly several times to empty its contents directly onto the skin.



For dogs of more than 25 kg:

For easy application the dog should be standing. The entire contents of the pipette should be applied evenly as 3 or 4 spots along the top of the back, from between the shoulders to the base of the tail. At each spot, part the coat until the skin is visible. Wherever possible apply to undamaged skin. Place the tip of the pipette on the skin and gently squeeze the pipette to expel a portion of its contents directly onto the skin. Do not apply an excessive amount of solution at any one spot, as that could cause some of the product to run down the animal's side.



10. WITHDRAWAL PERIOD

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Do not store above 25°C.

Store in the original package in order to protect from light and moisture.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton after {EXP}.

12. SPECIAL WARNINGS

For Animal Treatment Only.

Special warning for each target species:

Brief contact of the animal with water on one or two occasions between monthly treatments is unlikely to significantly reduce the efficacy of the product. However, frequent shampooing or immersion of the animal in water after treatment may reduce the efficacy of the product.

Parasite resistance to any particular class of anthelmintic may develop following frequent, repeated use of an anthelmintic of that class. Therefore, the use of this product should be based on the assessment of each individual case and on local epidemiological information about the current susceptibility of the target species in order to limit the possibility of a future selection for resistance. The use of the product should be based on the confirmed diagnosis of mixed infection (or risk of infection, where prevention applies) at the same time (see also the "INDICATIONS" and "DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION" sections of this package leaflet for additional information)..

Efficacy against adult *Dirofilaria repens* has not been tested under field conditions.

Special precautions for use in animals

The treatment of animals weighing less than 1 kg should be based on a risk-benefit assessment.

There is limited experience on the use of the product in sick and debilitated animals, thus the product should only be used based on a risk-benefit assessment for these animals.

Care should be taken that the contents of the pipette or the applied dose does not come into contact with the eyes or mouth of the recipient and/or other animals. Do not allow recently treated animals to groom each other. When the product is applied in 3 to 4 separate spots, specific care should be taken to prevent the animal licking the application sites.

This product contains moxidectin (a macrocyclic lactone), therefore special care should be taken with Collies, Old English Sheepdogs and related breeds or crossbreeds, to correctly administer the product as described under the section "ADVICE ON CORRECT ADMINISTRATION"; in particular, oral uptake by the recipient and/or other animals in close contact should be prevented.

The product should not enter water courses as it has harmful effects on aquatic organisms: moxidectin is highly toxic to aquatic organisms. Dogs should not be allowed to swim in surface waters for 4 days after treatment.

The safety of the product has only been evaluated in dogs classified as either Class 1 or 2 for heartworm disease in laboratory studies and in a few Class 3 dogs in a field study. Therefore the use in dogs with obvious or severe symptoms of the disease should be based on a careful benefit risk assessment by the treating veterinarian.

Although experimental overdose studies have shown that the product may be safely administered to dogs infected with adult heartworms, it has no therapeutic effect against adult *Dirofilaria immitis*. It is therefore recommended that all dogs 6 months of age or more, living in areas endemic for heartworm, should be tested for existing adult heartworm infection before being treated with the product. At the discretion of the veterinarian, infected dogs should be treated with an adulticide to remove adult heartworms. The safety of the product has not been evaluated when administered on the same day as an adulticide.

Imidacloprid is toxic for birds, especially canaries.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Avoid contact with skin, eyes or mouth.

Do not eat, drink or smoke during application. Wash hands thoroughly after use.

After application do not stroke or groom animals until the application site is dry. In case of accidental spillage onto skin, wash off immediately with soap and water. People with a known hypersensitivity to benzyl alcohol, imidacloprid or moxidectin should administer the product with caution. In very rare cases the product may cause skin sensitisation or transient skin reactions (for example numbness, irritation or burning/tingling sensation).

In very rare cases the product may cause respiratory irritation in sensitive individuals.

If the product accidentally gets into eyes, they should be thoroughly flushed with water.

If skin or eye symptoms persist, or the product is accidentally swallowed, seek medical advice immediately and show the package leaflet or label to the physician. Do not ingest. In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

In order to prevent children from getting access to pipettes, keep the pipette in the original packaging until ready for use and dispose of used pipettes immediately.

Treated animals should not be handled until the application site is dry, and children should not be allowed to play with treated animals until the application site is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals are not allowed to sleep with owners, especially children.

The solvent in the product may stain or damage certain materials including leather, fabrics, plastics and finished surfaces. Allow the application site to dry before permitting contact with such materials.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy and lactation. Laboratory studies with either imidacloprid or moxidectin in rats and rabbits have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. Use only accordingly to the benefit-risk assessment by the responsible veterinarian.

Interaction with other medicinal products and other forms of interaction:

During treatment with Moxiclear no other antiparasitic macrocyclic lactone should be administered. No interactions between Moxiclear and routinely used veterinary medicinal products or medical or surgical procedures have been observed. Safety of Moxiclear when administered on the same day as an adulticide to remove adult heartworms has not been evaluated.

Overdose (symptoms, emergency procedures, antidotes):

Up to 10 times the recommended dose was tolerated in adult dogs with no evidence of adverse effects or undesirable clinical signs. Five times the recommended minimum dose applied at weekly intervals for 17 weeks was investigated in dogs aged over 6 months and tolerated with no evidence of adverse effects or undesirable clinical signs.

The product was administered to puppies at up to 5 times the recommended dose, every 2 weeks for 6 treatments, and there were no serious safety concerns. Transient mydriasis, salivation, vomiting and transient rapid respiration were observed. After accidental oral ingestion or overdose, neurological signs (most of which are transient) such as ataxia, generalised tremors, ocular signs (dilated pupils, little pupillary reflex, nystagmus), abnormal respiration, salivation and vomiting may occur in very rare cases. Ivermectin-sensitive Collie dogs tolerated up to 5 times the recommended dose repeated at monthly intervals without any adverse effects, but the safety of application at weekly intervals has not been investigated in ivermectin-sensitive Collie dogs. When 40 % of the unit dose was given orally, severe neurological signs were observed. Oral administration of 10 % of the recommended dose produced no adverse effects. Dogs infected with adult heartworms tolerated up to 5 times the recommended dose, every 2 weeks for 3 treatments, without any adverse effects. In case of accidental oral uptake, symptomatic treatment should be administered. There is no known specific antidote. The use of activated charcoal may be beneficial.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Medicines should not be disposed of via wastewater or household waste. Moxiclear should not enter water courses as this may be dangerous for fish and other aquatic organisms. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

July 2019


15. OTHER INFORMATION

Imidacloprid is effective against larval flea stages and adult fleas. Flea larvae in the pet's surroundings are killed after contact with a pet treated with the product.

2.5 ml clear pipette with a film composed of 3 layers: a polypropylene/COC, solvent free lacquer laminate and a copolymer of polyethylene/EVOH/polyethylene. The pipettes are sealed within a child resistant 4-ply foil sachet composed of LDPE/nylon/aluminium foil/polyester film and presented in an outer box.

Pipettes are presented in cartons containing 1, 2, 3, 4, 6, 9, 12, 21 or 42 pipettes. Each pipette is individually sealed in a foil sachet.

Not all pack sizes may be marketed.



Approved: 29 July 2022