

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

1000ml or 125ml

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Optomease Vet 200mg/ml concentrate for solution for fish treatment

Benzocaine

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Benzocaine 200mg/ml

**3. PHARMACEUTICAL FORM**

Concentrate for solution for fish treatment

**4. PACKAGE SIZE**

<1000ml> <125ml>

**5. TARGET SPECIES**

Atlantic Salmon and Rainbow Trout

**6. INDICATION(S)**

For the anaesthesia and sedation of atlantic salmon and rainbow trout.

The product is not for use in open water and should always be used in an isolated treatment vessel.

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read package leaflet before use.

**8. WITHDRAWAL PERIOD**

Meat: 7 degree days

Do not use during stripping of fish eggs intended for human consumption.

**9. SPECIAL WARNING(S), IF NECESSARY**

Anaesthetics must be handled correctly. See package leaflet for user warnings.

**10. EXPIRY DATE**

EXP {month/year}

**11. SPECIAL STORAGE CONDITIONS**

Do not freeze.  
Store in the original packaging.  
Keep the packaging tightly closed.  
Protect from light.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only. To be supplied only on veterinary prescription.  
<POM>

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Virbac  
Premiere Avenue  
2065M-L I D  
06516 Carros Cedex  
France

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 05653/3008

**17. MANUFACTURER’S BATCH NUMBER**

Lot {number}

**B. PACKAGE LEAFLET**

## **PACKAGE LEAFLET**

### **Optomease Vet 200mg/ml concentrate for solution for fish treatment**

Benzocaine

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing Authorisation Holder:

Virbac

Premiere Avenue, 2065M-L I D, F-06516 Carros Cedex, France

Manufacturer responsible for batch release:

<Biofaktor Sp z.o.o.

Czysta 4 Str., 96-100 Skierniewice, Poland.>

<Naqua Ltd

Laboratory E12, Ground Floor, East Block, Building 500, Discovery Park, Ramsgate Road, Sandwich, Kent, CT13 9ND, UK.>

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Optomease Vet 200mg/ml concentrate for solution for fish treatment

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)**

Each ml contains:

**Active substance:**

Benzocaine 200mg

**Excipients:**

Patent blue V (E131) 0.008mg

The solution is a clear, blue liquid.

**4. INDICATION(S)**

For the anaesthesia and sedation of atlantic salmon and rainbow trout.

The product is not for use in open water and should always be used in an isolated treatment vessel.

**5. CONTRAINDICATIONS**

Avoid deep anaesthesia of young salmon during the final smolting period.

Sedating fish with the purpose of milking them for eggs intended for consumption is prohibited.

**6. ADVERSE REACTIONS**

None known.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon. Alternatively you can report via your national reporting system <*national system details*>.

## **7. TARGET SPECIES**

Atlantic salmon and rainbow trout.

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

For the anaesthesia and sedation of atlantic salmon and rainbow trout.

The product is designed to be dissolved in water. Depending on the desired level of anaesthesia, dilute 15 – 20ml per 100 litres of water of the veterinary medicinal product (equivalent to 30-40 mg of benzocaine per litre of water).

Dilute immediately after opening the product container.  
Use the diluted solution immediately.

The product is not for use in open water and should always be used in an isolated treatment vessel.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Do not feed the fish for at least 48 hours prior to anaesthesia and keep the water well oxygenated before exposure of the fish; immobilization usually occurs within a few minutes of immersion.

The maximum exposure time must not exceed 15 minutes. In general, the total exposure time of the fish to the solution should be controlled as there are different levels of tolerance among species and rearing conditions.

For recovery, transfer the anaesthetized fish into clear, well oxygenated water.

## **10. WITHDRAWAL PERIOD**

Meat: 7 degree days.  
Do not use during stripping of fish eggs intended for human consumption.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Do not freeze.  
Store in the original packaging.  
Keep the packaging tightly closed.  
Protect from light.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month.

## 12. SPECIAL WARNING(S)

### Special precautions for use in animals:

During anaesthesia, the fish must be closely monitored. It is recommended to test a few fish with the product before general application with the selected drug concentration and exposure time before large numbers are medicated, particularly when water temperature is at the upper or lower ends of the normal temperature ranges for the species being treated. The product should be dissolved in water of the same composition and characteristics as that to which the fish are accustomed.

To minimise damage and loss when medicated the level of sedation should allow fish to maintain their equilibrium and swimming position.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Benzocaine may cause hypersensitivity (allergy) following ingestion or contact with the skin or mucosa. In rare cases, benzocaine may cause methaemoglobinaemia (excessive methaemoglobin in the blood) in people who are hypersensitive. Cyanosis (blue skin or lips) may occur as a result of excessive methaemoglobin concentrations.

Avoid handling this product if you know you are sensitised to benzocaine, or if you have been advised not to work with such preparations.

As an anaesthetic, benzocaine may cause adverse reactions, including nervous, respiratory and cardiovascular alterations. This includes direct contact with skin, which can cause local anaesthesia and prolonged exposure may cause dermatitis.

This veterinary medicinal product also contains DMSO as an excipient, which may be irritant to the skin.

Avoid contact with the eyes, skin and clothing.

Do not eat, drink or smoke when handling the product or the anaesthetic bath.

Always wear impermeable gloves when handling the product and preparing stock solution or the anaesthetic bath.

In case of contact with the eyes, flush immediately with plenty of clean water. In case of skin contact, wash the exposed area immediately with plenty of water. Remove contaminated clothes immediately. If you develop symptoms following exposure, you should seek medical advice and show the package leaflet or the label to the doctor.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet / label to the doctor. Vomiting may be induced; activated charcoal may be administered.

Foetal deaths may occur following administration of benzocaine in human beings.



Pregnant women should be extra careful to avoid exposure if handling the product.

Wash hands thoroughly after use.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established for breeder fish. Use only according to the benefit/risk assessment by the responsible veterinarian.

Overdose (symptoms, emergency procedures, antidotes):

In case of overdose, immediately transfer the fish to clear, well oxygenated flowing water and make sure the mouth and gills are open.

High concentrations or prolonged exposure may cause medullary collapse, cardiac arrest and death. No antidote is available.

Incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

Other Precautions

In order to protect the environment, used solution must be transferred to a holding tank filled with water with subsequent controlled release for dilution in the effluent to be discharged from the facility. See disposal advice (section 13) for further details.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

This product is dangerous to fish and other aquatic organisms in the concentrated form. Do not contaminate ponds, streams, lochs or inlets with product or used packaging.

Used solution must be transferred to a holding tank filled with water with subsequent controlled release for dilution in the effluent to be discharged from the facility. The use of the product is limited to facilities equipped with suitable technology to guarantee safe discharges as indicated below.

Transfer of used solution to a holding tank filled with water and controlled release for dilution in effluent will ensure that the concentration of spent benzocaine in discharge water does not exceed 1 µg/L.

Dilution in effluent must ensure that the concentration of spent benzocaine in discharge water does not exceed the trigger of 1 µg/L safeguarding environmentally good water quality.

When releasing the solution from the holding tank, flow rates are calculated based on the following equation:

$$\text{Discharge (L/hr)} = \frac{\text{Farm flow rate (L/min)} \times 0.90 \text{ (safety factor)}}{\text{Holding tank concentration (mg/L)}} \times 60$$

| <b>Eg. Holding tank concentration (mg/L)</b> | <b>Farm flow rate (L/min)</b> | <b>Discharge flow from holding tank (L/h)</b> |
|--|-------------------------------|---|
| 5  | 10,000 / 20,000 / 30,000      | 108 / 216 / 324                               |
| 10   | 10,000 / 20,000 / 30,000      | 54 / 108 / 162                                |
| 20   | 10,000 / 20,000 / 30,000      | 27 / 54 / 81                                  |
| 40   | 10,000 / 20,000 / 30,000      | 14 / 27 / 41                                  |

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

November 2020

**15. OTHER INFORMATION**

Package size:

Bottle of 125ml with an integral 10ml doser

Bottle of 1000ml with an integral 25ml doser.

Not all pack sizes may be marketed.

To be supplied only on veterinary prescription.

*to be completed Nationally*