

## **LABELLING AND PACKAGE LEAFLET**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**CARDBOARD BOX – for 1 x 50 ml or 1 x 100 ml bottles**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Bravoxin suspension for injection

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each 1 ml contains:

<i>C. perfringens</i> type A ( $\alpha$ ) toxoid	$\geq 0.5$ IU
<i>C. perfringens</i> type B & C ( $\beta$ ) toxoid	$\geq 18.2$ IU
<i>C. perfringens</i> type D ( $\epsilon$ ) toxoid	$\geq 5.3$ IU
<i>C. chauvoei</i> whole culture, inac	$\geq 90\%$ protection
<i>C. novyi</i> toxoid	$\geq 3.8$ IU
<i>C. septicum</i> toxoid	$\geq 4.6$ IU
<i>C. tetani</i> toxoid	$\geq 4.9$ IU
<i>C. sordellii</i> toxoid	$\geq 4.4$ U
<i>C. haemolyticum</i> toxoid	$\geq 17.4$ U

**3. PACKAGE SIZE**

50 ml

100 ml

**4. TARGET SPECIES**

Cattle and sheep

**5. INDICATIONS**

**6. ROUTES OF ADMINISTRATION**

Subcutaneous use.

**7. WITHDRAWAL PERIODS**

Withdrawal period: zero days.

**8. EXPIRY DATE**

Exp. {mm/yyyy}

Once broached use within 8 hours.

**9. SPECIAL STORAGE PRECAUTIONS**

Store and transport refrigerated.  
Protect from light.  
Do not freeze.

**10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”**

Read the package leaflet before use.

**11. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**13. NAME OF THE MARKETING AUTHORISATION HOLDER**

MSD Animal Health UK Ltd.

**14. MARKETING AUTHORISATION NUMBERS**

Vm 01708/5094

**15. BATCH NUMBER**

Lot {number}

**16. SPECIAL WARNING(S), IF NECESSARY**

**17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: Read package leaflet.

**18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

**POM-VPS** Veterinary medicinal product subject to prescription.

## **PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

### **BOTTLE LABEL – 100 ml**

#### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Bravoxin suspension for injection

#### **2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Each 1 ml contains:

<i>C. perfringens</i> type A ( $\alpha$ ) toxoid	$\geq 0.5$ IU
<i>C. perfringens</i> type B & C ( $\beta$ ) toxoid	$\geq 18.2$ IU
<i>C. perfringens</i> type D ( $\epsilon$ ) toxoid	$\geq 5.3$ IU
<i>C. chauvoei</i> whole culture, inac	$\geq 90\%$ protection
<i>C. novyi</i> toxoid	$\geq 3.8$ IU
<i>C. septicum</i> toxoid	$\geq 4.6$ IU
<i>C. tetani</i> toxoid	$\geq 4.9$ IU
<i>C. sordellii</i> toxoid	$\geq 4.4$ U
<i>C. haemolyticum</i> toxoid	$\geq 17.4$ U

#### **3. TARGET SPECIES**

Cattle and sheep

#### **4. ROUTES OF ADMINISTRATION**

Subcutaneous use.

Read the package leaflet before use.

#### **5. WITHDRAWAL PERIODS**

Withdrawal period: zero days.

#### **6. EXPIRY DATE**

Exp. {mm/yyyy}

Once broached use within 8 hours.

#### **7. SPECIAL STORAGE PRECAUTIONS**

Store and transport refrigerated.

Do not freeze.

Protect from light.

Keep the vial in the outer carton.

**8. NAME OF THE MARKETING AUTHORISATION HOLDER**

MSD Animal Health UK Ltd.

**9. BATCH NUMBER**

Lot {number}

**10. SPECIAL WARNING(S), IF NECESSARY**

Shake well before use.

**11. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

**12. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only.

**POM-VPS** Veterinary medicinal product subject to prescription.

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**BOTTLE LABEL – 50 ml**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Bravoxin



**2. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES**

Clostridial cells and toxoids; see package leaflet.

50 ml

**3. BATCH NUMBER**

Lot {number}

**4. EXPIRY DATE**

Exp. {mm/yyyy}

Once broached use within 8 hours.

**5. ROUTE(S) OF ADMINISTRATION**

SC use.

**6. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

## **A. LABELLING**



## **PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Bravoxin suspension for injection for cattle and sheep

### **2. COMPOSITION**

Each 1 ml contains:

#### **Active substances:**

<i>C. perfringens</i> type A ( $\alpha$ ) toxoid	$\geq 0.5$ IU <sup>#</sup>
<i>C. perfringens</i> type B & C ( $\beta$ ) toxoid	$\geq 18.2$ IU*
<i>C. perfringens</i> type D ( $\epsilon$ ) toxoid	$\geq 5.3$ IU*
<i>C. chauvoei</i> whole culture, inactivated	$\geq 90\%$ protection**
<i>C. novyi</i> toxoid	$\geq 3.8$ IU*
<i>C. septicum</i> toxoid	$\geq 4.6$ IU*
<i>C. tetani</i> toxoid	$\geq 4.9$ IU*
<i>C. sordellii</i> toxoid	$\geq 4.4$ U <sup>1</sup>
<i>C. haemolyticum</i> toxoid	$\geq 17.4$ U <sup>#</sup>

\* ELISA According to Ph.Eur.

<sup>1</sup> In house ELISA

\*\* Guinea pig challenge test according to Ph.Eur.

<sup>#</sup> *In vitro* toxin neutralisation test based on haemolysis of sheep erythrocytes.

#### **Adjuvant:**

Aluminium<sup>1</sup> 3.026 – 4.094 mg

<sup>1</sup> from aluminium potassium sulphate (alum)

#### **Excipient:**

Thiomersal 0.05 – 0.18 mg

Light brown aqueous suspension that settles on storage.

### **3. TARGET SPECIES**

Cattle and sheep.

### **4. INDICATIONS FOR USE**

For the active immunisation of sheep and cattle against disease caused by *Clostridium perfringens* type A, *C. perfringens* type B, *C. perfringens* type C, *C. perfringens* type D, *C. chauvoei*, *C. novyi* type B, *C. septicum*, *C. sordellii* and *C. haemolyticum* and against tetanus caused by *C. tetani*. For the passive immunisation of lambs and calves against infections caused by the above mentioned clostridial species (except *C. haemolyticum* in sheep).



Onset of immunity:

Sheep and Cattle: 2 weeks after the basic vaccination course (as demonstrated by serology only).

Duration of active immunity:

As demonstrated by serology only:

Sheep: 1 year against *C. perfringens* type A, B, C and D, *C. novyi* type B, *C. sordellii*, *C. tetani*;  
< 6 months against *C. septicum*, *C. haemolyticum*, *C. chauvoei*;  
Cattle: 1 year against *C. tetani* and *C. perfringens* type D;  
< 1 year against *C. perfringens* type A, B and C;  
< 6 months against *C. novyi* type B, *C. septicum*, *C. sordellii*, *C. haemolyticum*, *C. chauvoei*.

An anamnestic humoral immune response (immunological memory) to all components was demonstrated 1 year following the basic course of vaccination.

Duration of passive immunity:

As demonstrated by serology only:

Lambs: At least 2 weeks for *C. septicum* and *C. chauvoei*;  
At least 8 weeks for *C. perfringens* type B and *C. perfringens* type C;  
At least 12 weeks for *C. perfringens* type A, *C. perfringens* type D, *C. novyi* type B, *C. tetani* and *C. sordellii*;  
No passive immunity was observed for *C. haemolyticum*.  
Calves: At least 2 weeks for *C. sordellii* and *C. haemolyticum*;  
At least 8 weeks for *C. septicum* and *C. chauvoei*;  
At least 12 weeks for *C. perfringens* type A, *C. perfringens* type B, *C. perfringens* type C, *C. perfringens* type D, *C. novyi* type B, and *C. tetani*.

## 5. CONTRAINDICATIONS

Do not use in sick or immunodeficient animals.

## 6. SPECIAL WARNINGS

Special warnings:

Vaccinate healthy animals only.

The effectiveness of the vaccine in providing passive immunity to young lambs and calves depends on these animals ingesting adequate amounts of colostrum on the first day of life.

Clinical trials have demonstrated that the presence of maternal derived antibodies (MDA), particularly against *C. tetani*, *C. novyi* type B, *C. perfringens* type A (calves only), *C. chauvoei* (lambs only) and *C. perfringens* type D may reduce the antibody response to vaccination in young lambs and calves. Therefore, to ensure an optimal response in young animals with high levels of MDA, the basic vaccination should be

delayed until the levels wane (which is after about 8-12 weeks of age, see section “Indications for use”).

Special precautions for safe use in the target species:

It is good practice to observe animals regularly for adverse reactions at the injection site following vaccination. It is recommended to seek medical advice from a veterinarian in case of a severe injection site reaction.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Pregnancy:

No side effects other than those already described in section “Adverse events” were seen when the vaccine was used in sheep and cattle between 8 and 2 weeks prior to parturition. In the absence of specific data, the use of the vaccine is not recommended during the first or second third of pregnancy. Avoid stress in pregnant ewes and cows.

Interaction with other medicinal products and other forms of interaction:

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Overdose:

In calves and lambs, local reactions may increase slightly if twice the recommended dose is administered (see section “Adverse events”).

Major incompatibilities:

Do not mix with any other veterinary medicinal product.

## 7. ADVERSE EVENTS

Cattle and sheep:

Very common (>1 animal / 10 animals treated):	Injection site swelling <sup>1</sup> .
Common (1 to 10 animals / 100 animals treated):	Injection site abscess, injection site skin discolouration <sup>2</sup> . Hyperthermia (elevated temperature) <sup>3</sup> .
Uncommon (1 to 10 animals / 1,000 animals treated):	Injection site pain <sup>4</sup> .
Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Anaphylactic type reaction (severe allergic reaction) <sup>5</sup> .

<sup>1</sup> Up to a mean value of 6 cm in sheep and 15 cm diameter in cattle; sometimes reactions of up to 25 cm diameter may be seen in cattle. Most local reactions resolve

within 3-6 weeks in sheep and in less than 10 weeks in cattle. In a minority of animals they may persist longer.

<sup>2</sup> Returns to normal as the local reaction resolves.

<sup>3</sup> Mild.

<sup>4</sup> For 1-2 days post first vaccination.

<sup>5</sup> In such cases appropriate treatment such as adrenaline should be administered without delay.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: Email: [adverse.events@vmd.gov.uk](mailto:adverse.events@vmd.gov.uk)  
Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

## **8. DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION**

Subcutaneous use.

*Dose:*

- Sheep: 1 ml – from 2 weeks of age
- Cattle: 2 ml – from 2 weeks of age

Administer by subcutaneous injection preferably in the loose skin on the side of the neck, observing aseptic precautions.

*Basic vaccination:* Two doses should be administered, 4-6 weeks apart (see section “Indications for use” and “Special Warnings”).

*Re-vaccination:* A single dose should be administered at 6 to 12 month intervals after the basic vaccination (see section “Indications for use”).

*Use in pregnancy:*

To provide passive protection of the offspring, via the colostrum, a single re-vaccination should be administered between 8 and 2 weeks before parturition, provided that animals have received a full basic vaccination course before pregnancy.

## **9. ADVICE ON CORRECT ADMINISTRATION**

Shake the bottle thoroughly before use.

Syringes and needles should be sterile before use and the injection should be made through an area of clean, dry skin taking precautions against contamination.

## **10. WITHDRAWAL PERIODS**

Zero days.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Store and transport refrigerated (2 °C – 8 °C). Do not freeze.  
Protect from light.

Do not use this veterinary medicinal product after the expiry date which is stated on the label and the carton. The expiry date refers to the last day of that month.

Shelf life after first opening the immediate packaging: 8 hours.

## **12. SPECIAL PRECAUTIONS FOR DISPOSAL**

Medicines should not be disposed of via wastewater.

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

## **13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.

## **14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES**

Vm 01708/5094

### Pack sizes:

Cardboard box with one bottle of 50 ml (50 doses of 1 ml or 25 doses of 2 ml).

Cardboard box with one bottle of 100 ml (100 doses of 1 ml or 50 doses of 2 ml).

Not all pack sizes may be marketed.

## **15. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST REVISED**

November 2023

Find more product information by searching for the 'Product Information Database' or 'PID' on [www.gov.uk](http://www.gov.uk)

## 16. CONTACT DETAILS

Marketing authorisation holder:

MSD Animal Health UK Ltd.  
Walton Manor, Walton  
Milton Keynes  
Buckinghamshire  
MK7 7AJ

Manufacturer responsible for batch release:

Intervet International B.V.  
Wim de Körverstraat 35  
5831 AN Boxmeer  
The Netherlands

Contact details to report suspected adverse reactions:

MSD Animal Health UK Ltd.  
Tel.: +44 (0)1908 685685

## 17. OTHER INFORMATION

For animal treatment only.

POM-VPS Veterinary medicinal product subject to prescription.