

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Carton for 20 ml, 50 ml, 100 ml and 250 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Inflacam 20 mg/ml solution for injection for cattle, pigs and horses
Meloxicam

2. STATEMENT OF ACTIVE SUBSTANCES

Meloxicam	20 mg/ml,
Ethanol (96%)	159.8 mg/ml.

3. PHARMACEUTICAL FORM

Solution for injection

4. PACKAGE SIZE

20 ml
50 ml
100 ml
250 ml

5. TARGET SPECIES

Cattle, pigs and horses

6. INDICATIONS

Read the package leaflet before use.

7. METHOD AND ROUTES OF ADMINISTRATION

Cattle
Single subcutaneous or intravenous injection.

Pigs

Single intramuscular injection. If required, a second administration can be given after 24 hours.

Horses

Single intravenous injection.

Read the package leaflet before use.

8. WITHDRAWAL PERIODS

Withdrawal period:

Cattle: meat and offal: 15 days; milk: 5 days.

Pigs: meat and offal: 5 days.

Horses: meat and offal: 5 days.

Not authorised for use in horses producing milk for human consumption.

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

10. EXPIRY DATE

EXP {month/year}

Shelf life after first opening the immediate packaging: 28 days.

Once broached, use by...

11. SPECIAL STORAGE CONDITIONS

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Read the package leaflet before use.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OF RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only. To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Chanelle Pharmaceuticals Manufacturing Ltd
Loughrea
Co. Galway
Ireland

16. MARKETING AUTHORISATION NUMBERS

Vm 08749/5013

17. MANUFACTURER’S BATCH NUMBER

Lot {number}

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

Label for 50 ml, 100 ml and 250 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Inflacam 20 mg/ml solution for injection for cattle, pigs and horses
Meloxicam

2. STATEMENT OF ACTIVE SUBSTANCES

Meloxicam	20 mg/ml,
Ethanol (96%)	159.8 mg/ml.

3. PHARMACEUTICAL FORM

Solution for injection

4. PACKAGE SIZE

50 ml
100 ml
250 ml

5. TARGET SPECIES

Cattle, pigs and horses

6. INDICATIONS

Read package leaflet before use.

7. METHOD AND ROUTES OF ADMINISTRATION

Cattle
SC or IV injection.

Pigs
IM injection.

Horses
IV injection.

8. WITHDRAWAL PERIODS

Withdrawal period:

Cattle: meat and offal: 15 days; milk: 5 days.

Pigs: meat and offal: 5 days.

Horses: meat and offal: 5 days.

Not authorised for use in horses producing milk for human consumption.

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

10. EXPIRY DATE

EXP {month/year}

Shelf life after first opening the immediate packaging: 28 days.

Once broached, use by...

11. SPECIAL STORAGE CONDITIONS

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Read the package leaflet before use.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OF RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only-to be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Chanelle Pharmaceuticals Manufacturing Ltd.
Loughrea
Co. Galway
Ireland

16. MARKETING AUTHORISATION NUMBER

Vm 08749/5013

17. MANUFACTURER'S BATCH NUMBER

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

Label for 20 ml bottles

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Inflacam 20 mg/ml solution for injection for cattle, pigs and horses
Meloxicam

2. QUANTITY OF THE ACTIVE SUBSTANCE

Meloxicam 20 mg/ml

3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES

20 ml

4. ROUTES OF ADMINISTRATION

Cattle
SC or IV injection.

Pigs
IM injection.

Horses
IV injection.

5. WITHDRAWAL PERIODS

Withdrawal period:
Cattle: meat and offal: 15 days; milk: 5 days.
Pigs: meat and offal: 5 days.
Horses: meat and offal: 5 days.

Not authorised for use in horses producing milk for human consumption.

6. BATCH NUMBER

Lot {number}

7. EXPIRY DATE

EXP {month/year}

Shelf life after first opening the immediate packaging: 28 days.

Once broached use by ...

8. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

PACKAGE LEAFLET:

Inflacam 20 mg/ml solution for injection for cattle, pigs and horses

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder

Chanelle Pharmaceuticals Manufacturing Ltd.,
Loughrea, Co. Galway, Ireland

Manufacturers responsible for batch release:

Chanelle Pharmaceuticals Manufacturing Ltd.,
Loughrea, Co. Galway, Ireland

and

Eurovet Animal Health B.V.
Handelsweg 25, 5531 AE Bladel, The Netherlands

and

Labiana Life Sciences, S.A., C/ Venus, 26, Pol. Ind. Can Parellada, Tarrasa, 08228
Barcelona

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Inflacam 20 mg/ml solution for injection for cattle, pigs and horses
Meloxicam

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS

One ml contains:

Meloxicam 20 mg,
Ethanol (96%) 159.8 mg.
Clear, yellow solution.

4. INDICATIONS

Cattle

For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in cattle.

For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in calves of over one week of age and young, non-lactating cattle.

For adjunctive therapy in the treatment of acute mastitis, in combination with antibiotic therapy.

For the relief of post-operative pain following dehorning in calves.

Pigs

For use in non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation.

For adjunctive therapy in the treatment of puerperal septicaemia and toxæmia (mastitis-metritis-agalactia syndrome) with appropriate antibiotic therapy.

Horses

For use in the alleviation of inflammation and relief of pain in both acute and chronic musculo-skeletal disorders.

For the relief of pain associated with equine colic.

5. CONTRAINDICATIONS

Do not use in horses less than 6 weeks of age.

Do not use in pregnant or lactating mares.

Do not use in animals suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

For the treatment of diarrhoea in cattle, do not use in animals of less than one week of age.

6. ADVERSE REACTIONS

A slight transient swelling at the injection site following subcutaneous administration was observed in less than 10 % of the cattle treated in clinical studies.

In horses, a transient swelling at the injection site can occur but resolves without intervention.

In very rare cases anaphylactoid reactions may occur and should be treated symptomatically.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

7. TARGET SPECIES

Cattle, pigs and horses

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Cattle

Single subcutaneous or intravenous injection at a dosage of 0.5 mg meloxicam/kg body weight (i.e. 2.5 ml/100 kg body weight) in combination with antibiotic therapy or with oral re-hydration therapy, as appropriate.

Pigs

Single intramuscular injection at a dosage of 0.4 mg meloxicam/kg body weight (i.e. 2.0 ml/100 kg body weight) in combination with antibiotic therapy, as appropriate. If required, a second administration of meloxicam can be given after 24 hours.

Horses

Single intravenous injection at a dosage of 0.6 mg meloxicam/kg body weight (i.e. 3.0 ml/100 kg body weight).

For use in the alleviation of inflammation and the relief of pain in both acute and chronic musculo-skeletal disorders, Inflacam 15 mg/ml oral suspension may be used for continuation of treatment at a dosage of 0.6 mg meloxicam/kg body weight, 24 hours after administration of the injection.

9. ADVICE ON CORRECT ADMINISTRATION

Avoid introduction of contamination during use.

10. WITHDRAWAL PERIODS

Cattle: meat and offal: 15 days; milk: 5 days.

Pigs: meat and offal: 5 days.

Horses: meat and offal: 5 days.

Not authorised for use in horses producing milk for human consumption.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Do not use after the expiry date (EXP) stated on the carton and vial.

Shelf life after first opening the container: 28 days.

12. SPECIAL WARNINGS

Treatment of calves with Inlacam 20 minutes before dehorning reduces post-operative pain. Inlacam alone will not provide adequate pain relief during the dehorning procedure. To obtain adequate pain relief during surgery co-medication with an appropriate analgesic is needed.

Special precautions for use in animals:

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

Avoid use in very severely dehydrated, hypovolaemic or hypotensive animals which require parenteral rehydration, as there may be a potential risk of renal toxicity.

In case of inadequate relief of pain when used in the treatment of equine colic, careful re-evaluation of the diagnosis should be made as this could indicate the need for surgical intervention.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Accidental self-injection may give rise to pain. People with known hypersensitivity to non-steroidal anti-inflammatory drugs (NSAIDs) should avoid contact with the veterinary medicinal product.

In case of accidental self-injection, seek medical advice immediately and show this package leaflet or the label to the physician.

Use during pregnancy and lactation:

Cattle and pigs: Can be used during pregnancy and lactation.

Horses: See section "Contraindications".

Interaction with other medicinal products and other forms of interaction:

Do not administer concurrently with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anti-coagulant agents.

Overdose (symptoms, emergency procedures, antidotes):

In the case of overdose, symptomatic treatment should be initiated.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Medicines should not be disposed of via wastewater or household waste.

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

October 2022

15. OTHER INFORMATION

Cardboard box containing one colourless glass injection vial of 20 ml, 50 ml, 100 ml or 250 ml.

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

United Kingdom (Northern Ireland)


VIRBAC

1 ère avenue 2065 m LID

FR-06516 Carros

France

Tel: + 33-(0)4 92 08 73 00

A handwritten signature in black ink, consisting of several loops and a final horizontal stroke.

Approved: 05 October 2022