

ANNEX II

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. STATEMENT OF THE MRLs**

A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer responsible for batch release

Elanco France S.A.S
26 Rue de la Chapelle
68330 Huningue
France

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Veterinary medicinal product subject to prescription.

C. STATEMENT OF THE MRLs

The Committee for Medicinal Products for Veterinary Use has recommended the inclusion of Monensin in Kexxtone in table 1 (Allowed substances) of the annex to Commission Regulation (EU) No 37/2010 as follows:

Pharmacologically active substance	Marker residue	Animal species	MRLs	Target tissues	Other provisions	Therapeutic classification
Monensin	Monensin A	Bovine	2 µg/kg 10 µg/kg 50 µg/kg 10 µg/kg 2 µg/kg	Muscle Fat Liver Kidney Milk	None	Anti-infectious agent/antibiotic

The excipients listed in section 6.1 of the SPC are either allowed substances for which table 1 of the annex to Commission Regulation (EU) No 37/2010 indicates that no MRLs are required, or are considered as not falling within the scope of Regulation (EC) No 470/2009 when used as in this veterinary medicinal product.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

FOIL-BAG CONTAINING 1,3 OR 5 CONTINUOUS-RELEASE INTRARUMINAL DEVICES

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Kexxtone 32.4g continuous-release intraruminal device for cattle
monensin

2. STATEMENT OF ACTIVE SUBSTANCES

35.2 g monensin sodium (equivalent to 32.4 g monensin)

3. PHARMACEUTICAL FORM

Continuous-release intraruminal device.

4. PACKAGE SIZE

1 intraruminal device
3 intraruminal devices
5 intraruminal devices

5. TARGET SPECIES

Cattle (dairy cows and heifers)

6. INDICATION(S)

7. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Intraruminal use

8. WITHDRAWAL PERIOD(S)

Meat and offal: zero days
Milk: zero days

9. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

Ingestion or oral exposure to monensin can be fatal in dogs, horses, other equines or guinea fowl. Do not allow dogs, horses, other equines or guinea fowl access to veterinary medicinal products containing monensin.

Due to the risk of bolus regurgitation, do not allow these species access to areas where treated cattle have been kept.

Exposure to the active substance may elicit an allergic response in susceptible persons. People with known hypersensitivity to monensin or any of the excipients should avoid contact with the veterinary medicinal product.

Do not eat, drink or smoke when handling the veterinary medicinal product.

Use gloves when handling an intraruminal device, including during retrieval of a regurgitated intraruminal device.

Remove gloves and wash hands and exposed skin after handling intraruminal devices.

10. EXPIRY DATE

EXP {month/year}

Once opened use by...

Shelf life after first opening the immediate packaging: 6 months

11. SPECIAL STORAGE CONDITIONS

Keep the foil tightly closed.

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of waste material in accordance with local requirements.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only. To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Elanco GmbH
Heinz-Lohmann-Str. 4
27472 Cuxhaven
Germany

16. MARKETING AUTHORISATION NUMBER(S)
--

Vm 52127/5032

17. MANUFACTURER'S BATCH NUMBER
--

Lot {number}

B. PACKAGE LEAFLET

PACKAGE LEAFLET:
Kexxtone 32.4g continuous-release intraruminal device for cattle

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER
AND OF THE MANUFACTURING AUTHORISATION HOLDER
RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Elanco GmbH
Heinz-Lohmann Strasse 4
Groden
Cuxhaven
D-27472
Germany

Manufacturer responsible for batch release:

Elanco France S.A.S
26 Rue de la Chapelle
68330 Huningue
France

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Kexxtone 32.4g continuous-release intraruminal device for cattle
monensin

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER
INGREDIENT(S)**

Monensin 32.4 g (equivalent to 35.2 g monensin sodium)

A cylindrical orange polypropylene intraruminal device uniquely identified with a number, fitted with wings, consisting of a core which presents as a stack of 12 subunits.

4. INDICATION(S)

For the reduction in the incidence of ketosis in the peri-parturient dairy cow/heifer which is expected to develop ketosis.

5. CONTRAINDICATIONS

Do not use in animals weighing less than 300 kg bodyweight.
Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

6. ADVERSE REACTIONS

In rare cases, digestive signs (e.g. diarrhoea, ruminant stomach disorder) have been observed.

In very rare cases, oesophagus obstruction has been observed.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

7. TARGET SPECIES

Cattle (dairy cows and heifers)

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Intraruminal use.

A single intraruminal device is to be administered to a dairy cow/heifer 3-4 weeks prior to expected calving, using an appropriate administration tool.

Kexxtone delivers an approximate average dose of 335 mg of monensin per day for approximately 95 days.

9. ADVICE ON CORRECT ADMINISTRATION

Follow directions carefully.

Adequate animal restraint is required to properly administer this intraruminal device. Such restraint must limit forward/backward motion and allow the animal's head to be held in the forward extended position and without pressure on the neck to prevent choking.

Each intraruminal device has an individual number along the intraruminal device body. This should be recorded with the corresponding animal identification number so that, should an intraruminal device be regurgitated, the animal can be identified. Fold wings down along the intraruminal device body and insert the intraruminal device into an appropriate administration tool, orifice end first.

Standing to one side of the animal, restrain it with its head and neck stretched forward and held firmly against your side. Grasp the animal with one hand in the corner of the animal's mouth. Introduce the administration tool into the mouth avoiding the front teeth. In order to avoid trauma and damage to the pharynx and oesophagus, do not use excessive force.

Insert the administration tool past the base of the tongue making sure to avoid the molar teeth. As the animal swallows the administration tool will move easily over the

base of the tongue. DO NOT USE EXCESSIVE FORCE. If resistance is encountered, withdraw the tool slightly and repeat the procedure. Be sure the head of the administration tool is past the base of the tongue. When the animal swallows, eject the intraruminal device from the administration tool.

Hold treated cattle in a confined area for 1 hour after administration to observe for failure to swallow or regurgitation. If this occurs re-administer the intraruminal device if undamaged. If damaged, administer a new intraruminal device. Recheck cattle for up to 4 days after dosing to observe for signs of an intraruminal device lodging in the oesophagus. In the event of early regurgitation, identify the animal by matching the animal ID number with the number on the intraruminal device.

Signs of lodging may include bloat which may be followed by coughing, drooling, inappetence and unthriftiness.

10. WITHDRAWAL PERIOD(S)

Meat and offal: zero days
Milk: zero days

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children

Keep the foil tightly closed.
Shelf life after first opening the immediate packaging. 6 months

Do not use this veterinary medicinal product after the expiry date which is stated on the package after "EXP". The expiry date refers to the last day of that month.

12. SPECIAL WARNING(S)

Special precautions for use in animals:

Identification of animals for treatment should be at veterinary discretion. Risk factors may include a history of energy-deficiency-related diseases, high body condition score and parity.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Exposure to the active substance may elicit an allergic response in susceptible individuals. People with known hypersensitivity to monensin or any of the excipients should avoid contact with the veterinary medicinal product.

Do not eat, drink or smoke when handling the veterinary medicinal product.
Use gloves when handling an intraruminal device, including during retrieval of a regurgitated intraruminal device.

Remove gloves and wash hands and exposed skin after handling intraruminal devices.

Other precautions

Ingestion or oral exposure to monensin can be fatal in dogs, horses, other equines or guinea fowl. Do not allow canine, horses, other equines or guinea fowl access to veterinary medicinal products containing monensin.

Due to the risk of bolus regurgitation, do not allow these species access to areas where treated cattle have been kept.

Keep dogs away from treated animals. Accidental ingestion of active ingredient by dogs has resulted in fatal consequences. In case of suspected ingestion by dogs, seek veterinary advice immediately.

Pregnancy and lactation:

Can be used during pregnancy and lactation.

Overdose (symptoms, emergency procedures, antidotes):

Accidental administration of more than one intraruminal device could result in some adverse reactions which are typical of monensin overdose, including decreased appetite, scouring and lethargy. These are generally transient. The highest tolerated dose is typically between 1 mg and 2 mg monensin/kg bodyweight/day.

Incompatibilities:

Not applicable

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product, waste materials derived from such veterinary medicinal products or regurgitated intraruminal devices should be disposed of in accordance with local requirements.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

<DD/MM/YY>

15. OTHER INFORMATION

Aluminium foil bag containing 1, 3 or 5 intraruminal device(s).

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the marketing authorisation holder.

Approved 12 July 2021

