

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

Carton for 100 ml and 250 ml

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Metacam 15 mg/ml oral suspension  
for pigs Meloxicam

**2. STATEMENT OF ACTIVE SUBSTANCES**

Meloxicam 15 mg/ml

**3. PHARMACEUTICAL FORM**

Oral suspension

**4. PACKAGE SIZES**

100 ml  
250 ml

**5. TARGET SPECIES**

Pigs

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Shake well before use.

Preferably mixed with small quantity of feed. Alternatively, prior to feeding or directly into the mouth. After use, close the bottle by replacing the cap, wash the measuring syringe with warm water and let it dry.

Read the package leaflet before use.

**8. WITHDRAWAL PERIOD(S)**

Withdrawal  
periods: Meat and  
offal: 5 days.

**9. SPECIAL WARNING(S), IF NECESSARY**

**10. EXPIRY DATE**

EXP. {month/year}  
Once opened use within 6 months.

**11. SPECIAL STORAGE CONDITIONS**

**12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet before use.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim Vetmedica  
GmbH 55216 Ingelheim/Rhein

GERMANY

**16. MARKETING AUTHORISATION NUMBERS**

EU/2/97/004/041 100 ml  
EU/2/97/004/042 250 ml

**17. MANUFACTURER'S BATCH NUMBER**

Lot {number}

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

**Bottle, 100 ml and 250 ml**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Metacam 15 mg/ml oral suspension  
for pigs Meloxicam

**2. STATEMENT OF ACTIVE SUBSTANCES**

Meloxicam 15 mg/ml

**3. PHARMACEUTICAL FORM**

**4. PACKAGE SIZES**

100 ml  
250 ml

**5. TARGET SPECIES**

Pigs

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Shake well before use.  
After use, close the bottle by replacing the cap, wash the measuring syringe with warm water and let it dry.

Read the package leaflet before use.

**8. WITHDRAWAL PERIOD(S)**

Withdrawal  
periods: Meat and  
offal: 5 days.

**9. SPECIAL WARNING(S), IF NECESSARY**

**10. EXPIRY DATE**

EXP {month/year}  
Once opened use within 6 months.

**11. SPECIAL STORAGE CONDITIONS**

**12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

**13. THE WORDS "FOR ANIMAL TREATMENT ONLY" AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"**

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim Vetmedica  
GmbH GERMANY

**16. MARKETING AUTHORISATION NUMBERS**

EU/2/97/004/041 100 ml  
EU/2/97/004/042 250 ml

**17. MANUFACTURER'S BATCH NUMBER**

Lot {number}



## **B. PACKAGE LEAFLET**

**PACKAGE LEAFLET:  
Metacam 15 mg/ml oral suspension for  
pigs**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION  
HOLDER AND OF THE MANUFACTURING AUTHORISATION  
HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder and manufacturer  
responsible for batch release Boehringer Ingelheim  
Vetmedica GmbH  
55216  
Ingelheim/  
Rhein  
GERMANY

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Metacam 15 mg/ml oral  
suspension for pigs  
Meloxicam

**3. STATEMENT OF THE ACTIVE SUBSTANCE AND OTHER INGREDIENTS**

One ml contains:  
Meloxicam 15 mg

Yellowish viscous oral suspension with a green tinge.

**4. INDICATION(S)**

For use in non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation.

For adjunctive therapy in the treatment of puerperal septicaemia and toxæmia (Mastitis-Metritis- Agalactia syndrome MMA) with appropriate antibiotic therapy.

**5. CONTRAINDICATIONS**

Do not use in pigs suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

## **6. ADVERSE REACTIONS**

None.

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Pigs

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Oral suspension to be administered at a dosage of 0.4 mg/kg body weight (i.e. 2.7 ml/100 kg) in combination with antibiotic therapy, as appropriate. If required, a second administration of meloxicam can be given after 24 hours. In cases of MMA with severely disturbed general demeanour (e.g. anorexia) the use of Metacam 20 mg/ml solution for injection is recommended.

Shake well before use.

After administration of the veterinary medicinal product, close the bottle by replacing the cap, wash the measuring syringe with warm water and let it dry.

## **9. ADVICE ON CORRECT ADMINISTRATION**

To be administered preferably mixed with a small quantity of feed. Alternatively to be given prior to feeding, or directly into the mouth. The suspension should be given using the measuring syringe provided in the package. The syringe fits onto the bottle and has a kg-body weight scale.

## **10. WITHDRAWAL PERIOD(S)**

Meat and offal: 5 days.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions. Shelf life after first opening of the container: 6 months.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and the bottle after EXP.

## **12. SPECIAL WARNING(S)**

Special precautions for use in animals:

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

Avoid use in very severely dehydrated, hypovolaemic or hypotensive pigs which require parenteral rehydration, as there may be a potential risk of renal toxicity.

Special precautions to be taken by the person administering the veterinary medicinal product to animals: People with known hypersensitivity to non-steroidal anti-inflammatory drugs (NSAIDs) should avoid contact with the veterinary medicinal product.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

This product can cause eye irritation. In case of contact with the eyes, immediately rinse thoroughly with water.

Pregnancy and lactation:

Can be used during pregnancy and lactation.

Interaction with other medicinal products and other forms of interaction:

Do not administer concurrently with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anticoagulant agents.

Overdose (symptoms, emergency procedures, antidotes):

In case of overdose symptomatic treatment should be initiated.

## **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

## **14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>.

## **15. OTHER INFORMATION**

100 ml or 250 ml bottle.  
Not all pack sizes may be marketed.