

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**CARDBOARD CARTON FOR BLISTER**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Apoquel 5.4 mg chewable tablets for dogs

oclacitinib

**2. STATEMENT OF ACTIVE SUBSTANCES**

5.4 mg oclacitinib per tablet (as oclacitinib maleate).

**3. PHARMACEUTICAL FORM**

Chewable tablet

**4. PACKAGE SIZE**

20 tablets  
50 tablets  
100 tablets

**5. TARGET SPECIES**

Dogs

**6. INDICATION(S)**

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use.

**8. WITHDRAWAL PERIOD(S)**

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

**10. EXPIRY DATE**

EXP {month/year}

**11. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture.  
Remaining tablet parts should be stored in the blister and be given at the next administration.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read the package leaflet.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Zoetis UK Limited  
1st Floor, Birchwood Building  
Springfield Drive  
Leatherhead  
Surrey  
KT22 7LP

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 42058/5001

**17. MANUFACTURER'S BATCH NUMBER**

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Apoquel 5.4 mg chewable tablets for dogs.

oclacitinib



**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Zoetis

**3. EXPIRY DATE**

EXP {month/year}

**4. BATCH NUMBER**

Lot {number}

**5. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

## **B. PACKAGE LEAFLET**

**PACKAGE LEAFLET:**

**Apoquel 3.6 mg chewable tablets for dogs**  
**Apoquel 5.4 mg chewable tablets for dogs**  
**Apoquel 16 mg chewable tablets for dogs**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder and manufacturer responsible for batch release:

Zoetis Belgium SA  
Rue Laid Burniat 1  
1348 Louvain-la-Neuve  
BELGIUM

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Apoquel 3.6 mg chewable tablets for dogs  
Apoquel 5.4 mg chewable tablets for dogs  
Apoquel 16 mg chewable tablets for dogs

oclacitinib

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)**

Each chewable tablet contains 3.6 mg, 5.4 mg or 16 mg (as oclacitinib maleate).

Light to dark brown pentagon shaped mottled tablets with score lines on both sides. The tablets are debossed with the corresponding strength ("S S" for 3.6 mg, "M M" for 5.4 mg and "L L" for 16 mg).

The tablets can be divided into equal halves.

**4. INDICATION(S)**

Treatment of pruritus associated with allergic dermatitis in dogs.  
Treatment of clinical manifestations of atopic dermatitis in dogs.

**5. CONTRAINDICATIONS**

Do not use in cases of hypersensitivity to oclacitinib or to any of the excipients.  
Do not use in dogs less than 12 months of age or less than 3 kg bodyweight.  
Do not use in dogs with evidence of immune suppression such as hyperadrenocorticism or with evidence of progressive malignant neoplasia as the active substance has not been evaluated in these cases.



## 6. ADVERSE REACTIONS

The common adverse reactions seen up to day 16 of the field trials are listed in the following table:

	Adverse reactions observed in atopic dermatitis study up to day 16		Adverse reactions observed in pruritus study up to day 7	
	Apoquel (n = 152)	Placebo (n = 147)	Apoquel (n = 216)	Placebo (n = 220)
Diarrhoea	4.6%	3.4%	2.3%	0.9%
Vomiting	3.9%	4.1%	2.3%	1.8%
Anorexia	2.6%	0%	1.4%	0%
New cutaneous or subcutaneous lumps	2.6%	2.7%	1.0%	0%
Lethargy	2.0%	1.4%	1.8%	1.4%
Polydipsia	0.7%	1.4%	1.4%	0%

After day 16, the following adverse reactions have been observed:

- pyoderma and non-specified dermal lumps have been observed very commonly;
- otitis, vomiting, diarrhoea, histiocytoma, cystitis, yeast skin infections, pododermatitis, lipoma, polydipsia, lymphadenopathy, nausea, increased appetite and aggression have been observed commonly.

Treatment-related clinical pathology changes were restricted to an increase in mean serum cholesterol and a decrease in mean leukocyte count, however, all mean values remained within the laboratory reference range. The decrease in mean leukocyte count observed in oclacitinib-treated dogs was not progressive, and affected all white blood cell counts (neutrophil, eosinophil and monocyte counts) except lymphocyte counts. Neither of these clinical pathology changes appeared clinically significant.

The development of papillomas was noted in a number of dogs in a laboratory study.

Anaemia and lymphoma have been reported very rarely in spontaneous reports.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

## 7. TARGET SPECIES

Dogs.



## 8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For oral use.

### Dosage and treatment schedule:

The recommended initial dose of Apoquel tablets to be given to the dog is to achieve 0.4 to 0.6 mg oclacitinib/kg bodyweight, administered orally, twice daily for up to 14 days.

For maintenance therapy (after the initial 14 days of treatment), the same dose (0.4 to 0.6 mg oclacitinib/kg bodyweight) should then be administered only once a day. The requirement for long-term maintenance therapy should be based on an individual benefit-risk assessment by the responsible veterinarian.

Apoquel tablets are chewable, palatable and readily consumed by the majority of dogs.

These tablets can be administered with or without food.

Please see dosing table below for the number of tablets required to achieve the recommended dose. The tablets are breakable along the score line.

Bodyweight (kg) of dog	Strength and number of tablets to be administered:		
	Apoquel 3.6 mg tablets	Apoquel 5.4 mg tablets	Apoquel 16 mg tablets
3.0–4.4	½		
4.5–5.9		½	
6.0–8.9	1		
9.0–13.4		1	
13.5–19.9			½
20.0–26.9		2	
27.0–39.9			1
40.0–54.9			1½
55.0–80.0			2

## **9. ADVICE ON CORRECT ADMINISTRATION**

Dogs should be carefully observed following administration to ensure that each tablet is swallowed.

## **10. WITHDRAWAL PERIOD(S)**

Not applicable.

## **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Store in the original package in order to protect from moisture.

Remaining tablet parts should be stored in the blister and be given at the next administration.

Do not use this veterinary medicinal product after the expiry date which is stated on the blister after EXP.

## **12. SPECIAL WARNING(S)**

### Special precautions for use in animals:

Oclacitinib modulates the immune system and may increase susceptibility to infection and exacerbate neoplastic conditions. Dogs receiving Apoquel tablets should therefore be monitored for the development of infections and neoplasia.

When treating pruritus associated with allergic dermatitis with oclacitinib, investigate and treat any underlying causes (e.g. flea allergic dermatitis, contact dermatitis, food hypersensitivity). Furthermore, in cases of allergic dermatitis and atopic dermatitis, it is recommended to investigate and treat complicating factors, such as bacterial, fungal or parasitic infections/infestations (e.g. flea and mange).

Given the potential for effects on certain clinicopathological parameters (see section 6), periodic monitoring with complete blood counts and serum biochemistry is recommended when dogs are on long-term treatment.

The tablets are flavoured. In order to avoid accidental ingestion, store tablets in a safe place out of reach of animals.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Wash hands after administration.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Ingestion of this product may be harmful for children. To avoid accidental ingestion, administer the tablet(s) to the dog immediately after removal from the blister packaging.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy and lactation, or in breeding male dogs, therefore its use is not recommended during pregnancy, lactation or in dogs intended for breeding.

Interaction with other medicinal products and other forms of interaction:

No drug interactions were observed in field studies where oclacitinib was administered concomitantly with veterinary medicinal products such as endo- and ectoparasiticides, antimicrobials and anti-inflammatories.

The impact of oclacitinib administration on vaccination with modified live vaccines, canine parvovirus (CPV), canine distemper virus (CDV) and canine parainfluenza (CPI) and inactivated rabies vaccine (RV), on 16 week old vaccine naive puppies has been studied. An adequate immune response (serology) to CDV and CPV vaccination was achieved when puppies were administered oclacitinib at 1.8 mg/kg bodyweight (bw) twice daily for 84 days. However, the findings of this study indicated a reduction in serological response to vaccination with CPI and RV in puppies being treated with oclacitinib compared to untreated controls. The clinical relevance of these observed effects for animals vaccinated while being administered oclacitinib (in accordance with the recommended dosing regimen) is unclear.

Overdose (symptoms, emergency procedures, antidotes):

Oclacitinib tablets were administered to healthy, one year old Beagle dogs twice daily for 6 weeks, followed by once per day for 20 weeks, at 0.6 mg/kg bw, 1.8 mg/kg bw and 3.0 mg/kg bw for a total of 26 weeks. Clinical observations that were considered likely to be related to oclacitinib treatment included: alopecia (local), papilloma, dermatitis, erythema, abrasions and scabbing/crusts, interdigital "cysts", and oedema of the feet.

Dermatitis lesions were mostly secondary to the development of interdigital furunculosis on one or more feet during the study with the number and frequency of observations increasing with increasing dose. Lymphadenopathy of peripheral nodes was noted in all groups, increasing in frequency with increasing dose, and was frequently associated with interdigital furunculosis.

Papilloma was considered treatment related, but not dose related.

There is no specific antidote and in case of signs of overdose the dog should be treated symptomatically.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

May 2022

## **15. OTHER INFORMATION**

Apoquel chewable tablets are supplied in blister packs with 20, 50 or 100 tablets per pack.

Not all pack sizes may be marketed.

Oclacitinib is a Janus kinase (JAK) inhibitor. It can inhibit the function of a variety of cytokines dependent on JAK enzyme activity. For oclacitinib, the target cytokines are those that are proinflammatory or have a role in allergic responses/pruritis. However, oclacitinib may also exert effects on other cytokines (for example, those involved in host defence or haematopoiesis) with the potential for unwanted effects.

Approved 30 June 2022

A handwritten signature in black ink, appearing to read "Hunter.", is positioned below the approval date. The signature is stylized and written in a cursive-like font.