

PARTICULARS TO APPEAR ON THE OUTER PACKAGE {Plastic box of 10 bottles of lyophilisate and 10 bottles of solvent; Plastic box of 50 bottles of lyophilisate and 50 bottles of solvent}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Purevax RCP FeLV lyophilisate and solvent for suspension for injection

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Per dose of 1 ml or 0.5 ml:

FHV (F2 strain)	$\geq 10^{4.9}$ CCID ₅₀
FCV (431 and G1 strains)	≥ 2.0 ELISA U.
FPV (PLI IV)	$\geq 10^{3.5}$ CCID ₅₀ .
FeLV recombinant canarypox virus (vCP97)	$\geq 10^{7.2}$ CCID ₅₀ .

3. PACKAGE SIZE

Lyophilisate (10 x 1 dose) + solvent (10 x 1 ml)
Lyophilisate (50 x 1 dose) + solvent (50 x 1 ml)
Lyophilisate (10 x 1 dose) + solvent (10 x 0.5 ml)
Lyophilisate (50 x 1 dose) + solvent (50 x 0.5 ml)

4. TARGET SPECIES

Cats.

5. INDICATION(S)

6. ROUTES OF ADMINISTRATION

Subcutaneous use.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {dd/mm/yyyy}
Once reconstituted use immediately.

9. SPECIAL STORAGE PRECAUTIONS

Store and transport refrigerated.
Protect from light.
Do not freeze.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Boehringer Ingelheim Vetmedica GmbH.

14. MARKETING AUTHORISATION NUMBERS

Vm 04491/5054

15. BATCH NUMBER

Lot {number}

16. SPECIAL WARNING(S), IF NECESSARY

17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: Read package leaflet

18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

POM-V

Veterinary medicinal product subject to prescription

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING
UNITS {Lyophilisate bottle}**

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Purevax RCP FeLV



2. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

1 dose

1 ml or 0.5 ml

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {dd/mm/yyyy}

5. ROUTE(S) OF ADMINISTRATION

6. THE WORDS “FOR ANIMAL TREATMENT ONLY”

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING
UNITS {Solvent bottle}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Purevax RCP FeLV solvent



2. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

1 ml or 0.5 ml

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {dd/mm/yyyy}

5. ROUTE(S) OF ADMINISTRATION

6. THE WORDS “FOR ANIMAL TREATMENT ONLY”

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Purevax RCP lyophilisate and solvent for suspension for injection.

2. COMPOSITION

Per dose of 1 ml or 0.5 ml:

Active substances:

Lyophilisate:

Attenuated feline rhinotracheitis herpesvirus (FHV F2 strain)	$\geq 10^{4.9}$ CCID ₅₀ ¹
Inactivated feline calicivirus (FCV 431 and G1 strains) antigens	≥ 2.0 ELISA U.
Attenuated feline panleucopenia virus (PLI IV)	$\geq 10^{3.5}$ CCID ₅₀ ¹

Solvent:

FeLV recombinant canarypox virus (vCP97)	$\geq 10^{7.2}$ CCID ₅₀ ¹
--	---

¹ cell culture infective dose 50%

Lyophilisate: friable pellet, homogeneous from beige to white.

Solvent: clear colourless liquid with presence of cell debris in suspension.

3. TARGET SPECIES

Cats.

4. INDICATIONS FOR USE

Active immunisation of cats aged 8 weeks and older:

- against feline viral rhinotracheitis to reduce clinical signs,
- against calicivirus infection to reduce clinical signs,
- against feline panleucopenia to prevent mortality and clinical signs.
- against leukaemia to prevent persistent viraemia and clinical signs of the related disease.

Onset of immunity:

- Rhinotracheitis, calicivirus and panleucopenia component: 1 week after primary vaccination course.
- Feline Leukaemia component: 2 weeks after primary vaccination course.

Duration of immunity:

- Rhinotracheitis, calicivirus and panleucopenia component: 1 year after the primary vaccination course and 3 years after the last re-vaccination.
- Feline leukaemia component: 1 year after the last re-vaccination.

5. CONTRAINDICATIONS

None.

6. SPECIAL WARNINGS

Vaccinate healthy animals only.

Special precautions for safe use in the target species:

It is recommended that a test for FeLV antigenaemia be carried out prior to vaccination.

Vaccination of FeLV positive cats is of no benefit.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Pregnancy and lactation:

Do not use during the whole pregnancy and lactation.

Interaction with other medicinal products and other forms of interaction:

Safety and efficacy data are available which demonstrate that this vaccine can be administered the same day but not mixed with Boehringer Ingelheim adjuvanted vaccine against rabies.

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product except the products mentioned above. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Overdose (symptoms, emergency procedures, antidotes):

No adverse event other than those already mentioned in section on "Adverse events" have been observed, except hyperthermia that may exceptionally last 5 days.

Major incompatibilities:

Do not mix with any other veterinary medicinal product, except the solvent supplied for use with the veterinary medicinal product and except those mentioned above.

7. ADVERSE EVENTS

Cats

Common (1 to 10 animals / 100 animals treated):

Apathy, anorexia, and hyperthermia¹.

Injection site reactions (pain, itching, oedema)².

Uncommon (1 to 10 animals / 1 000 animals treated):

Hypersensitivity reaction³.

Very rare (<1 animal / 10 000 animals treated, including isolated reports):

Emesis⁴.

¹ lasting usually for 1 or 2 days.

² slight pain at palpation, itching or limited oedema disappearing within 1 or 2 weeks at most.

³ may require appropriate symptomatic treatment.

⁴ mostly within 24 to 48 hours.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system:

Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

e-mail: adverse.events@vmd.gov.uk

8. DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION

Subcutaneous route.

After reconstitution of the lyophilisate with 0.5 ml or 1 ml of the solvent (depending on the presentation chosen), inject one dose of vaccine according to the following vaccination schedule:

Primary vaccination course:

- first injection: from 8 weeks of age,
- second injection: 3 to 4 weeks later.

Where high levels of maternal antibodies against rhinotracheitis, calicivirosis or panleucopenia components are expected to be present (e.g. in kittens of 9 to 12 weeks of age born from queens, which were vaccinated before pregnancy and/or with known or suspected previous exposure to the pathogen(s)), the primary vaccination course should be delayed until 12 weeks of age.

Revaccination:

- the first revaccination must be carried out one year after the primary vaccination course,

Subsequent revaccinations:

- Feline leukaemia component: every year
- Rhinotracheitis, calicivirosis and panleucopenia components: at intervals of up to three years.

9. ADVICE ON CORRECT ADMINISTRATION

Reconstitute gently the vaccine in order to obtain a uniform suspension with limited foam formation.

Visual appearance after reconstitution: clear slightly yellow suspension with presence of cell debris in suspension.

10. WITHDRAWAL PERIODS

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.
Store and transport refrigerated (2 °C – 8 °C).
Protect from light.
Do not freeze.

Do not use this veterinary medicinal product after the expiry date, which is stated on the carton and the bottle after Exp.

Shelf life after reconstitution according to directions: use immediately.

12. SPECIAL PRECAUTIONS FOR DISPOSAL

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 04491/5054

Plastic box containing:

10 x 1 dose of lyophilisate and 10 x 1 ml of solvent or
50 x 1 dose of lyophilisate and 50 x 1 ml of solvent or
10 x 1 dose of lyophilisate and 10 x 0.5 ml of solvent or
50 x 1 dose of lyophilisate and 50 x 0.5 ml of solvent.

Not all pack sizes may be marketed.

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. CONTACT DETAILS

Marketing authorisation holder:
Boehringer Ingelheim Vetmedica GmbH
55216 Ingelheim/Rhein
Germany

Manufacturer responsible for the batch release:
Boehringer Ingelheim Animal Health France SCS
Laboratoire Porte des Alpes
Rue de l'aviation

69800 Saint-Priest
France

Local representatives and contact details to report suspected adverse reactions:

United Kingdom (Great Britain)

Boehringer Ingelheim Animal Health UK Ltd., United Kingdom

Tel: + 44 1344 746957

17. OTHER INFORMATION

The feline leukaemia vaccine strain is a recombinant canarypox virus expressing the env and gag genes of FeLV-A. Under field conditions, only sub-group A is infective and immunisation against sub-group A provides full protection against A, B and C. After inoculation, the virus expresses the protective proteins, but does not replicate in the cat. As a consequence, the vaccine induces an immune status against feline leukaemia virus.

The product was shown to reduce excretion of feline calicivirus at onset of immunity and for one year after vaccination.

POM-V

For animal treatment only

Gavin Hall

Approved: 20 December 2024