

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **B. PACKAGE LEAFLET**

N.B. for the product there is no package leaflet. All information is directly on the container. Therefore the structure taken from the QRD8 template is the one foreseen for the leaflet

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**  
(Pressurised aluminium spray container containing 200 ml product)

Engemycin Spray, 25 mg/mL, cutaneous spray, suspension for cattle, sheep and pigs

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER  
AND OF THE MANUFACTURING AUTHORISATION HOLDER  
RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

MSD Animal Health UK Ltd.  
Walton Manor  
Walton  
Milton Keynes  
Buckinghamshire  
MK7 7AJ

Manufacturer for the batch release:

Intervet Productions S.r.l.  
Via Nettunense, Km 20,300  
04011 Aprilia (LT)  
Italy

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Engemycin Spray 25 mg/mL  
Cutaneous spray suspension for cattle, sheep and pigs

Oxytetracycline hydrochloride

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER  
INGREDIENT(S)**

Each mL contains:

Active substance:

Oxytetracycline hydrochloride	25.00 mg
(equivalent to oxytetracycline)	23.15 mg

Excipients:

Patent blue V (E131) as colouring agent	1.25 mg
---	---------

**4. PHARMACEUTICAL FORM**

Cutaneous spray, suspension  
Green to green-blue suspension

**5. INDICATIONS**

For the treatment of the following infections caused by, or associated with, organisms sensitive to oxytetracycline in cattle, sheep and pigs:

- Treatment of foot infections caused in particular by: *Dichelobacter nodosus*, *Fusobacterium necrophorum* and other *Fusobacterium* spp., and *Bacteroides* spp.
- Supporting treatment of superficial wound infections following surgery or physical injuries, e.g., tail biting in pigs, scratches and abrasions.

## 6. CONTRAINDICATIONS

Do not use for treatment of teats in order to prevent the product from getting into milk. Do not use in animals in cases of hypersensitivity to oxytetracycline or to any of the excipients.

## 7. ADVERSE REACTIONS

None known.

If you notice any serious effects or other effects not mentioned on this container, please inform your veterinary surgeon.

## 8. TARGET SPECIES

Cattle, sheep and pigs

## 9. DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION

For topical use only.

Shake well before use. The spray container is suitable to be used in upright and inverted positions. Before application properly clean the surface to be treated, spray the product for 1-2 seconds, at a distance of 15-20 cm, until the area has a homogeneous colour. Repeat the treatment every 12 hours for 1 to 3 days, depending on the healing process.

## 10. ADVICE ON CORRECT ADMINISTRATION

To achieve the best results in case of pedal lesions the following instructions are recommended:

- fully clean the foot area, completely removing foreign material, exudates and necrotic tissue
- keep the animal on dry ground for 12 hours after each application.

## 11. WITHDRAWAL PERIODS

### **Cattle, sheep:**

Meat and offal: Zero days

Milk: Zero hours

### **Pigs:**

Meat and offal: Zero days

Stained part of the pig skin must be removed prior to the rest of the animal being used for human consumption.

## **12. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Pressurised container: protect from sunlight and do not expose to temperatures exceeding 50°C. Keep away from sources of ignition - No smoking. Do not use after the expiry date stated on the container.

## **13. SPECIAL WARNING(S)**

For topical use only.

The animals should be treated in a well ventilated area.

Do not spray in or near the eyes of the animals.

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

No negative effects due to the use of the product during pregnancy and lactation are known.

Because of the risk of sensitisation and contact dermatitis, the user should avoid skin contact. Wear appropriate impermeable gloves whilst handling the product.

Because of risk of eye irritation, contact with the eyes should be avoided.

Protect the eyes and face.

Do not spray on a naked flame or any incandescent material.

Do not pierce or burn the container, even after use.

Avoid inhaling vapours.

Apply the product in the open air or in a well ventilated area.

Wash hands after use.

Do not eat or smoke whilst administering the product.

In case of accidental ingestion or in case of contact with eyes, seek medical advice immediately and show the label to the physician.

## **14. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIAL, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

## **15. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

August 2020

## **16. OTHER INFORMATION**

Pack size is one container containing 200 ml.

**17. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

For animal treatment only - to be supplied only on veterinary prescription.

**18. MARKETING AUTHORISATION NUMBER**

Vm 01708/4545

**19. MANUFACTURER’S BATCH NUMBER**

<Batch> <Lot> <BN> {number}

**20. EXPIRY DATE**

EXP: end MM/YY