LABELLING AND PACKAGE LEAFLET

A. LABELLING

<PARTICULARS TO APPEAR ON THE OUTER PACKAGE>

{CARTON FOR PACK SIZES OF 2,4, 6 AND 8 TABLETS}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Cazitel Plus Tablets

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

50 mg/tablet Praziquantel, 50 mg/tablet Pyrantel (equivalent to 144 mg pyrantel embonate) and 150 mg/tablet Febantel.

3. PACKAGE SIZE

2,4, 6, 8, 10, 12, 14, 16, 18, 20, 24, 28, 30, 32, 36, 40, 42, 44, 48, 50, 52, 56, 60, 70, 80, 84, 90, 98, 100, 104, 106, 120, 140, 150, 180, 200, 204, 206, 250, 280, 300, 500 and 1000 tablets.

4. TARGET SPECIES

Dogs.

5. INDICATION(S)

Treatment of mixed infections by nematodes and cestodes.

6. ROUTES OF ADMINISTRATION

Oral use.

1 tablet per 10 kg bodyweight.

The tablets can be given directly to the dog or disguised in food.

7. WITHDRAWAL PERIOD

N/A

8. EXPIRY DATE

EXP {month/year}

Discard any unused divided tablets immediately.

9. SPECIAL STORAGE PRECAUTIONS

Keep the blister in the outer carton.

10. THE WORDS "READ THE PACKAGE LEAFLET BEFORE USE"

Read the package leaflet before use

11. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

12. THE WORDS "KEEP OUT OF THE SIGHT AND REACH OF CHILDREN"

Keep out of the sight and reach of children.

13. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Chanelle Pharmaceuticals Manufacturing Ltd Loughrea Co. Galway Ireland

14. MARKETING AUTHORISATION NUMBER(S)

Vm 08749/5038

15. BATCH NUMBER

BN{number}

16. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

17. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE [Distribution category]

NFA-VPS

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

{BLISTER FOIL TEXT}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT



2. QUALITATIVE AND QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCE(S)

Each tablet contains 50 mg Praziquantel, 50 mg Pyrantel (equivalent to 144 mg pyrantel embonate) and 150 mg Febantel.

3. BATCH NUMBER

BN {number}

4. EXPIRY DATE

EXP {month/year}

5. NAME OF THE MARKETING AUTHORISATION HOLDER

Chanelle Pharmaceuticals Manufacturing Ltd.

6. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For Animal Treatment Only.

B. PACKAGE LEAFLET

PACKAGE LEAFLET Cazitel Plus Tablets for dogs

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Cazitel Plus Tablets For Dogs.

2. COMPOSITION

Each tablet contains Praziquantel 50 mg, 50 mg Pyrantel (equivalent to 144 mg Pyrantel Embonate and Febantel 150 mg.

The veterinary medicinal product is a pale yellow tablet with a cross breakline on one side.

The tablets can be divided into 2 or 4 equal parts.

3. TARGET SPECIES

Dogs

4. INDICATIONS FOR USE

Treatment of mixed infections by nematodes and cestodes of the following species **Nematodes:**

Ascarids: Toxocara canis, Toxascaris leonina (adult and late immature forms).

Hookworms: Uncinaria stenocephala, Ancylostoma caninum (adults).

Whipworms: Trichuris vulpis (adults).

Cestodes:

Tapeworms: *Echinococcus* species, *(E. granulosus, E. multilocularis), Taenia* species,

(*T. hydatigena, T. pisiformis, T. taeniformis) Dipylidium caninum* (adult and immature forms).

5. CONTRAINDICATIONS

Do not use simultaneously with piperazine compounds.

Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

6. SPECIAL WARNING(S)

Special warnings:

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*.

Tapeworm infestation is certain to re-occur unless control of intermediate hosts such as fleas, mice etc. is undertaken.

Tapeworm infestation is unlikely in pups less than 6 weeks of age.

Parasite resistance to any particular class of anthelmintic may develop following frequent, repeated use of an anthelmintic of that class.

Unnecessary use of antiparasitics or use deviating from the instructions given in the SPC may increase the resistance selection pressure and lead to reduced efficacy.

The decision to use the product should be based on confirmation of the parasitic species and burden, or of the risk of infection based on its epidemiological features, for each individual animal.

In the absence of risk of co-infection with nematodes or cestodes, a narrow spectrum product should be used.

The possibility that other animals in the same household can be a source of reinfection with nematodes and cestodes should be considered, and these should be treated as necessary with an appropriate product.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In case of accidental ingestion, seek medical advice and show the package leaflet to the physician.

In the interests of good hygiene, persons administering the tablets directly to a dog or adding them to the dog's food should wash their hands afterwards. For animal treatment only.

Pregnancy:

Teratogenic effects attributed to high doses of febantel have been reported in sheep and rats. No studies have been performed in dogs during early pregnancy. Use of the product during pregnancy should be in accordance with a benefit risk assessment by the responsible veterinarian. The use is not recommended during the first 4 weeks of pregnancy in dogs. Do not exceed the stated dose when treating pregnant bitches.

Interactions with other medicinal products and other forms of interaction:

Do not use simultaneously with piperazine compounds as the anthelmintic effects of pyrantel and piperazine may be antagonised.

Concurrent use with other cholinergic compounds(e.g.foxim) can lead to toxicity. If you are uncertain, and your dog uses other veterinary medicinal products, check with a veterinary surgeon or pharmacist.

<u>Overdose:</u>

The combination of praziquantel, pyrantel embonate and febantel is well tolerated in dogs. In safety studies, a single dose of 5 times the recommended dose or greater gave rise to occasional vomiting.

Other Precautions:

Echinococcosis represents a hazard for humans. As echinococcosis is a notifiable disease to the World Organisation for Animal Health (OIE), specific guidelines on the treatment and follow-up, and on the safeguard of persons, need to be obtained from the relevant competent authority.

7. ADVERSE EVENTS

Dogs:

Very rare (<1	animal	/ 10,000	Digestive tract disorders (diarrhoea, emesis)
animals tre	ated,	including	
isolated reports):			

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Oral use.

. . .

The recommended dose rates are: 15 mg/kg bodyweight febantel, 5 mg/kg pyrantel (equivalent to 14.4 mg/kg pyrantel embonate) and 5 mg/kg praziquantel. 1 Cazitel Plus tablet per 10 kg (22 lbs) bodyweight. The tablets can be given directly to the dog ordisguised in food. No starvation is needed before, or after, treatment. The tablet can be divided in two or four equal parts.

Dosage table:			
Bodyweight (kg)	Tablets		
1⁄2 - 2.5	1/4		
2.6-5.0	1/2		
5.1-10.0	1		
10.1-15.0	11/2		
15.1-20.0	2		
20.1-25.0	21/2		
25.1-30.0	3		
30.1-35.0	31/2		
35.1-40.0	4		
>40.1	1 tablet per 10 kg		

Underdosing could result in ineffective use and may favour resistance development. The advice of a veterinarian should be sought regarding the need for and the frequency of repeat administration.

9. ADVICE ON CORRECT ADMINISTRATION

To ensure a correct dosage, body weight should be determined as accurately as possible.

10. WITHDRAWAL PERIOD

Not applicable.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions. Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month. Discard any unused divided tablets immediately.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

NFA-VPS

14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 08749/5038

The veterinary medicinal product is presented in either:

Individual strips composed of aluminium foil 30 μ m/30 gsm extruded polythene, containing 2, 4, 6, 8, 10, 12, 14, 16, 18 or 20 tablets.

or

Individual blisters composed of 45 μ m, soft temper aluminium foil and 25 μ m hard temper aluminium foil, containing 2 or 8 tablets.

The strips or blisters are packed into cartons containing either 2,4, 6, 8, 10, 12, 14, 16, 18, 20, 24, 28, 30, 32, 36, 40, 42, 44, 48, 50, 52, 56, 60, 70, 80, 84, 90, 98, 100, 104, 106, 120, 140, 150, 180, 200, 204, 206, 250, 280, 300, 500 or 1000 tablets.

Not all pack sizes may be marketed.

15. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST REVISED

February 2023

16. CONTACT DETAILS

Marketing authorisation holder and manufacturer responsible for batch release Chanelle Pharmaceuticals Manufacturing Ltd., Dublin Road, Loughrea, Co. Galway, Ireland

Local representatives and contact details to report suspected adverse reactions: Zoetis UK Limited 1st Floor, Birchwood Building Springfield Drive Leatherhead Surrey KT22 7LP

Revised: June 2023 AN: 01230/2022

Approved 20 June 2023