ANNEX III

LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Box containing 1 litre bottle
Box containing 2.5 L, 4.5 L or 8 L pouches

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Neoprinil pour-on 5 mg/ml pour-on solution

2. STATEMENT OF ACTIVE SUBSTANCES

Each ml contains:

Eprinomectin 5.00 mg

3. PACKAGE SIZE

1 litre
2.5 L
4.5 L
8 L

4. TARGET SPECIES

Cattle

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

For external use.
Pour-on use.

7. WITHDRAWAL PERIODS

Withdrawal period:

Meat and offal : 15 days.
Milk : zero hours.

8. EXPIRY DATE

Exp. :{mm/yyyy}

(Bottle only:) Once opened, use within 1 year.
(Pouches only:) Once opened, use within 2 years.

9. SPECIAL STORAGE PRECAUTIONS

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Virbac

14. MARKETING AUTHORISATION NUMBER

Vm 05653/3047

15. BATCH NUMBER

Lot {number}

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

2.5 and 5 L cans

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Neoprinil pour-on 5 mg/ml pour-on solution

2. STATEMENT OF ACTIVE SUBSTANCES

Each ml contains:

Eprinomectin 5.00 mg

3. PACKAGE SIZE

2.5 L
5L

4. TARGET SPECIES

Cattle

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

For external use.
Pour-on use.

7. WITHDRAWAL PERIODS

Withdrawal period:

Meat and offal : 15 days.
Milk : zero hours.

8. EXPIRY DATE

Exp. :{mm/yyyy}

Once opened, use within 1 year.
Once opened, use by . . .

9. SPECIAL STORAGE PRECAUTIONS

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Virbac

14. MARKETING AUTHORISATION NUMBER

Vm 05653/3047

15. BATCH NUMBER

Lot {number}

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

1 litre bottle
2.5, 4.5 and 8 L pouches

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Neoprinil pour-on 5 mg/ml pour-on solution

2. STATEMENT OF ACTIVE SUBSTANCES

Each ml contains:

Eprinomectin 5.00 mg

3. TARGET SPECIES

Cattle

4. ROUTES OF ADMINISTRATION

For external use.
Pour-on use.

Read the package leaflet before use.

5. WITHDRAWAL PERIODS

Withdrawal period:
Meat and offal : 15 days.
Milk : zero hours.

6. EXPIRY DATE

Exp. :{mm/yyyy}

(Bottle only:) Once opened, use within 1 year.
(Pouches only:) Once opened, use within 2 years.

Once opened, use by . . .

7. SPECIAL STORAGE PRECAUTIONS

8. NAME OF THE MARKETING AUTHORISATION HOLDER

Virbac

9. BATCH NUMBER

Lot {number}

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Neoprinil pour-on 5mg/ml pour-on solution for cattle

2. Composition

Each ml contains:

Active substance:
Eprinomectin 5.00 mg

Excipients:
Butylhydroxytoluene (E321) 0.10 mg
All-rac-alpha-tocopherol (E307) 0.06 mg

Slightly yellowish, clear oily solution.

3. Target species

Cattle.

4. Indications for use

In beef and dairy cattle :
Treatment of infestations by the following parasites sensitive to eprinomectin :

Gastrointestinal roundworms (adults and L4 larvae): *Ostertagia ostertagi* (including inhibited L4 larvae), *Ostertagia lyrata* (only adults), *Haemonchus placei, Trichostrongylus axei, Trichostrongylus colubriformis, Cooperia sp.* (including inhibited L4), *Cooperia oncophora, Cooperia punctata, Cooperia pectinata, Cooperia surnabada, Bunostomum phlebotomum, Nematodirus helvetianus, Oesophagostomum radiatum, Oesophagostomum* sp. (only adults), *Trichuris discolor* (only adults);
Lungworms : *Dictyocaulus viviparus* (adults and L4);
Warbles (parasitic stages) : *Hypoderma bovis, Hypoderma lineatum*;
Mange mites : *Chorioptes bovis, Sarcoptes scabiei* var. *bovis*;
Sucking lice : *Linognathus vituli, Haematopinus eurysternus, Solenopotes capillatus*;
Biting lice : *Damalinia bovis*;
Flies : *Haematobia irritans*

The veterinary medicinal product protects the animals against reinfestations with:
- *Nematodirus helvetianus* for 14 days;
- *Trichostrongylus axei and Haemonchus placei* for 21 days;
- *Dictyocaulus viviparus, Cooperia oncophora, Cooperia punctata, Cooperia surnabada, Oesophagostomum radiatum* and *Ostertagia ostertagi* for 28 days.

5. Contraindications

Do not use in animal species other than those listed in sections “Target species” and “Indications for use”.
Do not administer orally or by injection.
Do not use in cases of hypersensitivity to the active substance or to any of the excipients.
Avermectins may not be well tolerated in non-target species (including dogs, cats and horses). Cases of mortality are reported in dogs, especially Collies, bobtail and related breeds and crosses, and also in turtles/tortoises.

6. Special warnings

Special warnings:

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:
- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.
- Underdosing, which may be due to underestimation of bodyweight, misadministration of the veterinary medicinal product, or lack of calibration of the dosing device (if any).

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used.

To date no resistance to eprinomectin (a macrocyclic lactone) has been reported in cattle within the EU. However, resistance to other macrocyclic lactones has been reported in parasite species in cattle within the EU. Therefore, use of this veterinary medicinal product should be based on local (regional, farm) epidemiological information about susceptibility of nematodes and recommendations on how to limit further selection for resistance to anthelmintics.

If there is a risk for re- infection, the advice of a veterinarian should be sought regarding the need for and frequency of repeat administration.

For the best results the veterinary medicinal product should be part of a planned programme to control both internal and external parasites of cattle based on the epidemiology of these parasites.

Special precautions for safe use in the target species:

For external use only.
To avoid adverse reactions due to the death of warble larvae in the oesophagus or backbone, it is recommended to administer the veterinary medicinal product after the end of the swarming of the warbles and before the larvae reach their sites in the body; consult a veterinary surgeon to know the appropriate treatment period.
For effective use, the veterinary medicinal product should not be applied to areas of the backline covered with mud or manure. The veterinary medicinal product should be applied only on healthy skin.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product may be irritating to the skin and eyes and may cause hypersensitivity (allergic reactions).
Avoid direct contact with the skin or eyes during treatment and when handling recently treated animals.
People with known hypersensitivity to eprinomectin should avoid contact with the veterinary medicinal product.
Personal protective equipment consisting of rubber gloves, boots and a waterproof coat should be worn when applying the veterinary medicinal product.
If accidental skin contact occurs, wash the affected area immediately with soap and water. If accidental eye exposure occurs, flush eyes immediately with water.
Should clothing become contaminated, remove as soon as possible and launder before re-use. This veterinary medicinal product may affect the central nervous system if accidentally ingested. Avoid accidental ingestion of the veterinary medicinal product, including by hand to mouth contact. In case of accidental ingestion, wash the mouth out with water, seek medical advice immediately and show the package leaflet or the label to the physician.
Do not smoke, eat or drink while handling the veterinary medicinal product.
Wash hands after use

Special precautions for the protection of the environment:

Eprinomectin is very toxic to aquatic organisms, is persistent in soils and may accumulate in sediments. Faeces containing eprinomectin excreted onto pasture by treated animals may temporarily reduce the abundance of dung feeding organisms. Following treatment of cattle with the veterinary medicinal product, levels of eprinomectin that are potentially toxic to dung fly species may be excreted over a period of more than 4 weeks and may decrease dung fly abundance during that period. In case of repeated treatments with eprinomectin (as with veterinary medicinal products of the same anthelmintic class) it is advisable not to treat animals every time on the same pasture to allow dung fauna populations to recover.
Eprinomectin is inherently toxic to aquatic organisms. The veterinary medicinal product should be used only according to the label instructions. Based on the excretion profile of eprinomectin when administered as the pour-on formulation, treated animals should not have access to watercourses during the first 7 days after treatment.

Pregnancy and lactation:

Laboratory studies in rats and rabbits have not produced any evidence of teratogenic or embryotoxic effects due to the use of eprinomectin at therapeutic doses. The safety of the veterinary medicinal product in cattle has been established during pregnancy and lactation and in reproductive bulls. Can be used during pregnancy and lactation as well as in reproductive bulls.

Interaction with other medicinal products and other forms of interaction:

Since eprinomectin binds strongly to plasmatic proteins, this should be taken into account if it is used in association with other molecules having the same characteristics.

Overdose:

No signs of toxicity have been observed after administration of up to 5 times the recommended dose. No specific antidote has been identified.

Major incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

7. Adverse events

Cattle:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):
Licking at application site[1], Application site twitching[1], Application site reaction (e.g. application site skin scaling, application site skin squamosis)[2]

[1] Transient.
[2] Minor.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: {national system details}.

8. Dosage for each species, routes and method of administration

For external use.
Pour-on use.
To be administered topically in one single treatment at the dose rate of 500 µg eprinomectin per kg bodyweight equivalent to 1 ml per 10 kg bodyweight.
Apply the pour-on solution along the mid-line of the back in a narrow strip between the withers and tail head.
To ensure a correct dosage, bodyweight should be determined as accurately as possible and accuracy of the dosing device should be checked. If animals are to be treated collectively rather than individually, they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- and over- dosing.

All the animals belonging to the same group should be treated at the same time.

9. Advice on correct administration

Squeeze-Measure Pour-on system (1 litre bottle)
1 and 2. Remove the protective aluminium seal from the bottle.
3 and 4. Screw the dosing cup to the bottle.

Set the dose by turning the top section of the cup to align the correct bodyweight with the pointer.
When body weight is between marking, use the higher setting.
5. Hold the bottle upright and squeeze it to deliver a slight excess of the required dose as indicated by the calibration lines.
6 and 7. By releasing the pressure, the dose automatically adjusts to the correct level.
Take the dosing cup off the bottle after use and screw the cap on the bottle.



Can (2.5 litre and 5 litre can)
Connect an appropriate dosing gun and draw-off tubing to the back-pack as follows.
1 and 2. Remove the protective aluminium seal from the bottle.
3. Replace the shipping cap with the cap having the draw-off tubing. Tighten the draw-off cap.
4. Connect one side of the tube to the draw-off cap and the other side to the dosing gun.
5. Gently prime the dosing gun, checking before use that all connections are tight.
Follow the gun manufacturer's instructions for adjusting the dose and proper use and maintenance of the dosing gun and draw-off tubing.
When body weight is between marking, use the higher setting.

2.5 L & 5 L



FlexiBag (2.5 litre, 4.5 litre and 8 litre flexible pouch)
Connect an appropriate dosing gun to the FlexiBag as follows.
1 to 4. Connect one side of the tube to the coupling E-lock system draw-off and the other side to the dosing gun.
5 and 6. Plug the E-lock coupling system to the FlexiBag.
7. Gently prime the dosing gun, checking before use that all connections are tight.
Follow the gun manufacturer's instructions for adjusting the dose and proper use and maintenance of the dosing gun.
When body weight is between marking, use the higher setting.



10. Withdrawal periods

Meat and offal: 15 days.
Milk: zero hours.

11. Special storage precautions

Keep out of the sight and reach of children.

This veterinary medicinal product does not require any special storage conditions.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp. The expiry date refers to the last day of that month.

Shelf life after first opening the immediate packaging (bottles and cans): 1 year.
Shelf life after first opening the immediate packaging (pouches): 2 years.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater or household waste.

This veterinary medicinal product should not enter water courses as eprinomectin is extremely dangerous for fish and other aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty container.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack sizes

- 1 L bottles with a dosing device,
- 2.5 L and 5 L cans,
- 2.5 L, 4.5 L and 8 L pouches.

Not all pack sizes may be marketed.

Vm 05653/3047

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. Contact details

Marketing authorisation holder and manufacturer responsible for batch release and contact details to report suspected adverse reactions:

VIRBAC
1ère avenue 2065 m LID
06516 Carros
France

Local representatives and contact details to report suspected adverse reactions:

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

17. Other information

Like other macrocyclic lactones, eprinomectin has the potential to adversely affect non-target organisms. Following treatment, excretion of potentially toxic levels of eprinomectin may take place over a period of several weeks. Faeces containing eprinomectin excreted onto pasture by treated animals may reduce the abundance of dung feeding organisms which may impact on the dung degradation.
Eprinomectin is very toxic to aquatic organisms and may accumulate in sediments.
Eprinomectin is persistent in soils.

Gavin Hall

Approved: 21 July 2025