

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

**CARTON**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Benazecare Flavour 20 mg Tablets for Dogs

**2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Benazepril hydrochloride 20mg

**3. PHARMACEUTICAL FORM**

Tablet  
Beef flavoured, beige, divisible oval tablets.

**4. PACKAGE SIZE**

14 tablets  
28 tablets  
56 tablets  
140 tablets

**5. TARGET SPECIES**

Dogs

**6. INDICATION**

For the treatment of congestive heart failure in dogs.

**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use. Read the package leaflet before use.

**8. WITHDRAWAL PERIOD**

Not applicable

**9. SPECIAL WARNING(S), IF NECESSARY**

Read the package leaflet before use.

**10. EXPIRY DATE**

EXP {month/year}

**11. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C.

Store in a dry place.

Divided tablets should be stored in the blister pack. Any divided tablet portion remaining after 48 hours should be discarded.

Keep the blister pack in the outer carton.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable**

For animal treatment only - to be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE REACH AND SIGHT OF CHILDREN”**

Keep out of the reach and sight of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Ecuphar NV  
Legeweg 157-i  
8020 Oostkamp  
Belgium

**16. MARKETING AUTHORISATION NUMBER**

Vm 32742/4037

POM-V

VPA 10778/003/002

POM

**17. MANUFACTURER’S BATCH NUMBER**

BN:

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**Blister pack**

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Benazecare Flavour 20 mg Tablets for Dogs

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Ecuphar NV

**3. EXPIRY DATE**

EXP {month/year}

**4. BATCH NUMBER**

BN:

**5. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

## **B. PACKAGE LEAFLET**

## **PACKAGE LEAFLET**

Benazecare Flavour 20 mg Tablets for Dogs

### **1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder:

Ecuphar NV  
Legeweg 157-i  
8020 Oostkamp  
Belgium

Manufacturer for the batch release:

Lelypharma B.V.  
Zuiveringweg 42,  
8243 PZ  
Lelystad  
The Netherlands

### **2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Benazecare Flavour 20 mg Tablets for Dogs

### **3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)**

Each tablet contains 20 mg benazepril hydrochloride.  
Beef flavoured, beige, divisible oval tablets.

### **4. INDICATION**

Benazecare Flavour 20mg tablets belong to a group of medicines called Angiotensin Converting Enzyme (ACE) inhibitors. It is prescribed by the veterinary surgeon for the treatment of congestive heart failure in dogs.

### **5. CONTRAINDICATIONS**

Do not use in case of hypersensitivity to the active substance benazepril hydrochloride or to any ingredient of the tablets.

Do not use in cases of hypotension (low blood pressure), hypovolemia (low blood volume), hyponatraemia (low blood sodium levels) or acute renal failure.

Do not use in cases of cardiac output failure due to aortic or pulmonary stenosis.

Do not use in pregnant or lactating dogs because the safety of benazepril hydrochloride has not been established during pregnancy or lactation in this species.

## 6. ADVERSE REACTIONS

Some dogs with congestive heart failure may exhibit vomiting or fatigue during treatment.

In dogs with chronic kidney disease there may be a moderate increase in levels of creatinine, an indicator of kidney function, in the blood. This is likely due to the effect of the medication in reducing the blood pressure within the kidney and is therefore not necessarily a reason for treatment to be stopped, unless the animal is showing other adverse reactions.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

## 7. TARGET SPECIES

Dogs

## 8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Benazecare Flavour 20mg tablets should be given orally once daily, with or without food. The duration of treatment is unlimited.

Benazecare Flavour 20mg tablets should be administered orally at a minimum dose of 0.25 mg (range 0.25-0.5) benazepril hydrochloride/kg body weight once daily, according to the following table:

Weight of dog (kg)	BENAZECARE FLAVOUR 20mg	
	Standard Dose	Double Dose
>20 – 40	0.5 tablet	1 tablet
>40 – 80	1 tablet	2 tablets

In dogs with congestive heart failure, the dose may be doubled, still administered once daily, to a minimum dose of 0.5 mg (range 0.5-1.0) benazepril hydrochloride/kg body weight if judged necessary and advised by the veterinary surgeon. Always follow the dosing instructions given by the veterinary surgeon.

## 9. ADVICE ON CORRECT ADMINISTRATION

Not applicable.

## 10. WITHDRAWAL PERIOD

Not applicable.

## 11. SPECIAL STORAGE PRECAUTIONS

Keep out of the reach and sight of children.  
Do not store above 25°C

Store in a dry place

Divided tablets should be stored in the blister pack. Any divided tablet portion remaining after 48 hours should be discarded. Keep the blister pack in the outer carton.

Do not use after the expiry date stated on the carton.

## **12. SPECIAL WARNINGS**

### **Special warnings for dogs**

The efficacy and safety of benazepril hydrochloride has not been established in dogs below 2.5 kg body weight.

### **Special precautions for use in animals**

In cases of chronic kidney disease, your veterinarian will check the hydration status of your pet before starting therapy, and may recommend that regular blood tests are carried out during therapy in order to monitor plasma creatinine concentrations and blood erythrocyte counts.

### **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

Wash hands after use.

In case of accidental oral ingestion, seek medical advice immediately and show the label or the package leaflet to the physician.

Pregnant women should take special care to avoid accidental oral exposure because ACE inhibitors have been found to affect the unborn child during pregnancy in humans.

### **Use during pregnancy, lactation**

Do not use during pregnancy or lactation. The safety of benazepril hydrochloride has not been established in breeding, pregnant or lactating dogs.

### **Interactions**

Inform the veterinary surgeon if the animal is taking, or has recently taken, any other medicines.

In dogs with congestive heart failure, benazepril hydrochloride has been given in combination with digoxin, diuretics, pimobendan and anti-arrhythmic products without evidence of associated adverse reactions.

In humans, the combination of ACE inhibitors and NSAIDs (Non-Steroidal Anti-Inflammatory Drugs) can lead to reduced anti-hypertensive efficacy or impaired kidney function. The combination of benazepril hydrochloride and other anti-hypertensive agents (e.g. calcium channel blockers,  $\beta$ -blockers or diuretics), anaesthetics or sedatives may lead to additive hypotensive effects. Therefore, concurrent use of NSAIDs or other medications with a hypotensive effect should be considered with care.

Your veterinary surgeon may recommend to closely monitor kidney function and for signs of hypotension (lethargy, weakness etc) and treat these if necessary.

Interactions with potassium-preserving diuretics like spironolactone, triamterene or amiloride cannot be ruled out. Your veterinary surgeon may recommend to monitor

plasma potassium concentrations when using benazepril hydrochloride in combination with a potassium-sparing diuretic because of the risk of hyperkalaemia (high blood potassium).

### **Overdose**

Transient reversible hypotension (low blood pressure) may occur in cases of accidental overdose. Therapy should consist of intravenous infusion of warm isotonic saline.

### **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIAL, IF ANY**

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

### **14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

October 2022

### **15. OTHER INFORMATION**


#### **Pharmacodynamic properties**

Benazepril hydrochloride is a prodrug hydrolysed in vivo to its active metabolite, benazeprilat. Benazeprilat is a highly potent and selective inhibitor of the angiotensin converting enzyme (ACE), thus preventing the conversion of inactive angiotensin I to active angiotensin II and thereby also reducing synthesis of aldosterone. Therefore, it blocks effects mediated by angiotensin II and aldosterone, including vasoconstriction of both arteries and veins, retention of sodium and water by the kidney and remodelling effects (including pathological cardiac hypertrophy and degenerative renal changes).

Benazepril hydrochloride causes long-lasting inhibition of plasma ACE activity in dogs, with more than 95% inhibition at peak effect and significant activity (>80% in dogs) persisting 24 hours after dosing.

Benazepril hydrochloride reduces the blood pressure and volume load on the heart in dogs with congestive heart failure.

In contrast with other ACE inhibitors, benazeprilat is excreted equally by both biliary and urinary routes in dogs, and therefore no adjustment of the dose of benazepril hydrochloride is necessary in the treatment of cases with renal insufficiency.



Approved 12 October 2022



