

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

CARDBOARD BOX

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Enrox Max 100 mg/ml solution for injection

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Each ml contains 100 mg of enrofloxacin.

Excipients:

Benzyl alcohol (E1519) 20 mg

Butyl alcohol 30 mg

3. PACKAGE SIZE

100 ml

4. TARGET SPECIES

Cattle and pigs



5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Cattle: **s.c., i.v.**

Pigs: **i.m.**

7. WITHDRAWAL PERIODS

Withdrawal period:

Cattle:

Meat and offal: s.c.: 14 days

i.v.: 7 days

Milk: s.c.: 120 hours

i.v.: 72 hours

Pigs:

Meat and offal: i.m.: 12 days

8. EXPIRY DATE

Exp.
Once opened use within 28 days.

9. SPECIAL STORAGE PRECAUTIONS

Store in the original package. Do not freeze.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE»

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

KRKA

14. MARKETING AUTHORISATION NUMBERS

Vm 01656/5066

15. BATCH NUMBER

Lot

16. SPECIAL WARNING(S), IF NECESSARY

17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND SAFE USE IF APPLICABLE

POM-V

Veterinary medicinal product subject to prescription'

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

GLASS VIAL 100 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Enrox Max 100 mg/ml solution for injection

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Each ml contains 100 mg of enrofloxacin.

Excipients:

Benzyl alcohol (E1519) 20 mg

Butyl alcohol 30 mg

3. TARGET SPECIES

Cattle and pigs



4. ROUTES OF ADMINISTRATION

Cattle: **s.c., i.v.**

Pigs: **i.m.**

Read the package leaflet before use.

5. WITHDRAWAL PERIODS

Withdrawal period:

Cattle:

Meat and offal: s.c.: 14 days

i.v.: 7 days

Milk: s.c.: 120 hours

i.v.: 72 hours

Pigs:

Meat and offal: i.m.: 12 days

6. EXPIRY DATE

Exp

Once opened use within 28 days.

7. SPECIAL STORAGE PRECAUTIONS

Store in the original package. Do not freeze.

8. NAME OF THE MARKETING AUTHORISATION HOLDER

KRKA

9. BATCH NUMBER

Lot

10 SPECIAL WARNING(S), IF NECESSARY

11. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

12. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE

For animal treatment only.

POM-V ('Veterinary medicinal product subject to prescription')

B. PACKAGE LEAFLET

PACKAGE LEAFLET

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Enrox Max 100 mg/ml solution for injection for cattle and pigs

2. COMPOSITION

Each ml contains:

Active substance:

Enrofloxacin 100 mg

Excipients:

Benzyl alcohol (E1519) 20 mg

Butyl alcohol 30 mg

Clear, yellow solution.

3. TARGET SPECIES

Cattle and pigs.



4. INDICATIONS FOR USE

Cattle:

For the treatment of respiratory tract infections caused by enrofloxacin-susceptible *Histophilus somni*, *Mannheimia haemolytica*, *Pasteurella multocida*, and *Mycoplasma* spp. For the treatment of mastitis caused by enrofloxacin-susceptible *E. coli*.

Pigs:

For the treatment of bacterial bronchopneumonia caused by enrofloxacin-susceptible *Actinobacillus pleuropneumoniae*, *Pasteurella multocida* and complicated by *Haemophilus parasuis* as secondary pathogen in pigs.

5. CONTRAINDICATIONS

Do not use for prophylaxis.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in animals with central nervous system-associated seizure disorders. Do not use in the presence of existing disorders of cartilage development or musculoskeletal damage around functionally significant or weight-bearing joints.

Do not use in known cases of resistance against other (fluoro)quinolones due to the potential for cross-resistance.

6. SPECIAL WARNING(S)

Special precautions for safe use in the target species:

Official and local antimicrobial policies should be taken into account when the veterinary medicinal product is used.

Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly, to other classes of antimicrobials.

Whenever possible, fluoroquinolones should only be used based on susceptibility testing.

Use of the veterinary medicinal product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to the fluoroquinolones and may decrease the effectiveness of treatment with other quinolones due to the potential for cross resistance.

If there is no clinical improvement observed within 2-3 days of therapy, re-evaluation of treatment and susceptibility testing may be necessary.

Enrofloxacin is eliminated renally. As with all fluoroquinolones, delayed excretion can therefore be expected in the presence of existing renal damage.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Direct contact with the skin should be avoided due to sensitisation, contact dermatitis and possible hypersensitivity reactions.

People with known hypersensitivity to (fluoro)quinolones should avoid contact with the veterinary medicinal product.

Wash hands after use.

In the event of accidental splash into the eye, rinse with large amounts of clean water. If irritation occurs, seek medical advice.

Do not eat, drink or smoke while handling the veterinary medicinal product.

Take care to avoid accidental self-injection.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or label to the physician.

Pregnancy and lactation:

Can be used during pregnancy and lactation.

Interactions with other medicinal products and other forms of interaction:

Antagonist effects due to concurrent administration of macrolides and tetracyclines may occur. Enrofloxacin may interfere with the metabolism of theophylline, decreasing theophylline clearance resulting in increased plasma levels of theophylline.

Overdose:

In cattle, a dose of 25 mg/kg bodyweight administered by the subcutaneous route for 15 consecutive days is tolerated without any clinical symptoms. Clinical signs seen in gross overdosage include lethargy, lameness, ataxia, slight salivation and muscle tremors.

In pigs, doses of around 25 mg active ingredient per kg body weight and above may cause lethargy, loss of appetite and ataxia.

Do not exceed the recommended dose. In accidental overdose there is no antidote and treatment should be symptomatic.

Major incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

7. ADVERSE EVENTS

Cattle:

Rare (1 to 10 animals / 10,000 animals treated):

Injection site inflammation (swelling, redness)¹

Circulatory shock²

Very rare (<1 animal / 10,000 animals treated, including isolated reports):

Digestive tract disorder³

¹Resolves within a few days, no treatment required

²Following intravenous administration

³Reported in calves.

Pigs:

Rare (1 to 10 animals / 10,000 animals treated):

Injection site inflammation (swelling, redness)¹

¹Resolves within a few days, no treatment required

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system {<https://www.gov.uk/report-veterinary-medicine-problem>}.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

To ensure a correct dosage, body weight should be determined as accurately as possible.

Cattle:

For respiratory infections: administer by subcutaneous injection (s.c.):

A single dose of 7.5 mg enrofloxacin /kg body weight/day (7.5 ml of the veterinary medicinal product/100 kg body weight/day)

In cases of severe or chronic respiratory tract infections, a second injection may be required after 48 hours.

For *E. coli* mastitis in cattle: administer by slow intravenous injection (i.v.).

5 mg enrofloxacin/kg body weight/day (5.0 ml of the veterinary medicinal product/100 kg body weight/day) daily for 2-3 days.

Pigs:

For respiratory infections: administer by intramuscular injection (i.m.) in neck musculature behind the ear:

A single dose of 7.5 mg enrofloxacin /kg bodyweight/day (0.75 ml of the veterinary medicinal product/10 kg body weight/day)

In cases of severe or chronic respiratory tract infections, a second injection may be required after 48 hours.

9. ADVICE ON CORRECT ADMINISTRATION

Repeated injections should be administered at different sites.

Cattle: not more than 15 ml of the veterinary medicinal product (7.5 ml in calves) should be administered at one subcutaneous injection site.

Pigs: not more than 7.5 ml of the veterinary medicinal product should be administered at one intramuscular injection site.

10. WITHDRAWAL PERIOD(S)

Cattle:

| | |
|-----------------|---|
| Meat and offal: | after subcutaneous administration (s.c.): 14 days |
| | after intravenous administration (i.v.): 7 days |
| Milk: | after subcutaneous administration (s.c.): 120 hours |
| | after intravenous administration (i.v.): 72 hours |

Pigs:

| | |
|-----------------|--|
| Meat and offal: | after intramuscular administration (i.m.): 12 days |
|-----------------|--|

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Store in the original package. Do not freeze.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp. The expiry date refers to the last day of that month.

Shelf life after first opening the immediate packaging: 28 days.

12. SPECIAL PRECAUTIONS FOR DISPOSAL

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 01656/5066

Cardboard box with one 100 ml vial.

15. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST REVISED

December 2023

Find more product information by searching for the 'Product Information Database' or 'PID' on www.gov.uk.

16. CONTACT DETAILS

Marketing authorisation holder.:

KRKA, d.d., Novo mesto,
Šmarješka cesta 6,
8501 Novo mesto,
Slovenia

Manufacturer responsible for batch release:

KRKA d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia
TAD Pharma GmbH, Heinz-Lohmann-Straße 5, 27472 Cuxhaven, Germany


Local representatives and contact details to report suspected adverse reactions:

KRKA UK Ltd
United Kingdom
Tel: 02071 646 156
info.uk@krka.biz

17. OTHER INFORMATION

POM-V ('Veterinary medicinal product subject to prescription')

For animal treatment only.



Approved: 21 March 2024