

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

Polyester/ Aluminium/ Polyethylene Bags (10 sponges, 25 sponges and 50 sponges presentations) [*text appearing on label as no carton box will be used*].

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Chronogest CR 20 mg controlled release vaginal sponge for sheep

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Flugestone acetate, 20 mg/sponge.

17.9 mg flugestone equivalent to 20 mg flugestone acetate.

Excipients:

Hydroxypropylcellulose 20 mg/sponge

Macrogol 4000 20 mg/sponge

3. PACKAGE SIZE

Polyethylene bag containing 10, 25 or 50 sponges.

4. TARGET SPECIES

Sheep (ewe and ewe lamb).

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Vaginal use.

7. WITHDRAWAL PERIODS

Withdrawal period:

Meat and offal: 2 days after withdrawal of sponges.

Milk: zero hours, including the treatment time.

8. EXPIRY DATE

Exp. {mm/yyyy}

9. SPECIAL STORAGE PRECAUTIONS

Store below 25 °C.
Store in the original package.
Store in a dry place.
Once packaging is opened, any unused product should be discarded.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

MSD Animal Health UK Ltd.

14. MARKETING AUTHORISATION NUMBERS

Vm 01708/5087

15. BATCH NUMBER

Lot {number}

16. SPECIAL WARNING(S), IF NECESSARY

Read the package leaflet before use.

The product should not be administered by pregnant women or women suspected to be pregnant.

17. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Disposal: Read package leaflet.

18. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

POM-V Veterinary medicinal product subject to prescription.

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Chronogest CR 20 mg controlled release vaginal sponge for sheep

2. COMPOSITION

Each sponge contains:

Active substance(s)

17.9 mg flugestone equivalent to 20 mg flugestone acetate.

List of excipients

Hydroxypropylcellulose, 20 mg

Macrogol 4000, 20 mg

White cylindrical polyester polyurethane medicated sponge equipped with string.

3. TARGET SPECIES

Sheep (ewe and ewe lamb).

4. INDICATIONS FOR USE

In ewes and ewe lambs, in combination with PMSG (Pregnant Mare Serum Gonadotropin)

- Induction and synchronization of oestrus and ovulation (non cycling ewes during seasonal anoestrus and ewe lambs).
- Synchronization of oestrus and ovulation (cycling ewes and ewe-lambs).

5. CONTRAINDICATIONS

None.

6. SPECIAL WARNINGS

Special precautions for safe use in the target species:

- The repeated use of sponges within one year has not been studied.
- The use of a vaginal applicator designed for ewes or ewe lambs is recommended to correctly insert sponges and to avoid vaginal injuries.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Personal protective equipment consisting of single use gloves should be worn when handling the veterinary medicinal product.

The veterinary medicinal product should not be administered by pregnant women or women suspected to be pregnant.

Direct contact with the skin should be avoided. If accidental contact with the skin occurs, wash the affected zone with soap and water. Wash hands after treatment and before meals.

Human exposure to this product can affect fertility.

Pregnancy and lactation:

The use is not recommended during pregnancy.

Can be used during lactation.

Fertility:

- The repeated treatment with the product combined with PMSG may trigger the appearance of PMSG antibodies in some ewes. This in turn may affect the time of ovulation and result in reduced fertility when combined with fixed time artificial insemination at 55h following sponge removal.

Interaction with other medicinal products and other forms of interaction:

The sponges should not be used in conjunction with alcohols, cresols, phenols, sheep dips or similar disinfectants.

Overdose:

A five time overdose of flugestone acetate (100 mg/sponge) did not result in observable side effects.

Major incompatibilities:

None known.

7. ADVERSE EVENTS

Sheep (ewe and ewe lamb):

Very rare
(<1 animal / 10,000 animals treated, including isolated reports):
Vaginal discharge ¹

¹ muco-purulent discharge may be observed at sponge removal. It is not associated with clinical signs and does not alter fertility.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system: Email: adverse.events@vmd.gov.uk

Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

8. DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION

The dose is one sponge per animal independent of the body weight, breed, type and season.

For vaginal use using an applicator.

9. ADVICE ON CORRECT ADMINISTRATION

The dose is one sponge per animal irrespective of body weight, breed, type (dairy or meat) and season.

The sponge is inserted intra-vaginally using an applicator.

Duration of sponge residence is 14 days. At the end of the administration period, the sponge is gently removed by pulling on its string.

To obtain an optimal synchronization of ovulation, an intra-muscular injection of PMSG (range 300-700 IU) is recommended (i.m.) at the time of sponge removal.

In case fixed time artificial insemination is applied, it is recommended to perform it 55 h after sponge removal.

10. WITHDRAWAL PERIODS

Meat and offal: 2 days after withdrawal of sponges.

Milk: zero hours, including the treatment time.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Store below 25 °C.

Store in the original package.

Store in a dry place.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after Exp. The expiry date refers to the last day of that month.

Once packaging is opened, any unused product should be discarded.

12. SPECIAL PRECAUTIONS FOR DISPOSAL

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

These measures should help to protect the environment.

13. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 01708/5087

Not all pack sizes may be marketed.

15. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST REVISED

October 2023

Find more product information by searching for the 'Product Information Database' or 'PID' on www.gov.uk.

16. CONTACT DETAILS

Marketing authorisation holder:

MSD Animal Health UK Ltd.

Walton Manor, Walton

Milton Keynes

Buckinghamshire

MK7 7AJ

Manufacturer responsible for batch release:

Intervet Productions S.A.

Rue De Lyons

Igoville

27460

France

Contact details to report suspected adverse reactions:

MSD Animal Health UK Ltd.

Tel.: +44 (0)1908 685685

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

17. OTHER INFORMATION

For animal treatment only.

POM-V Veterinary medicinal product subject to prescription.

Approved 10 January 2024

