

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

Polyethylene back pack container

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

CLiKZiN 12.5 mg/ml Pour-On Suspension for Sheep.

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Dicyclanil 12.5mg/ml (1.25% w/v).

3. PHARMACEUTICAL FORM

Pour-on Suspension.

4. PACKAGE SIZE

0.8 litres

2.2 litres

5 litres

5. TARGET SPECIES

Sheep

6. INDICATION(S)

Prevention of blowfly strike on sheep due to *Lucilia sericata*.

7. METHOD AND ROUTE(S) OF ADMINISTRATION

CLiKZiN 12.5 mg/ml Pour-On Suspension for Sheep is applied according to the following recommendations:

Bodyweight (kg)	Dose Volume (ml)
10-20	20
21-30	24
31-50	30
>50	36

(Guide dose volumes correspond to 0.7–2 ml (7.5–25 mg dicyclanil) per kg bodyweight.)

To ensure administration of a correct dose, body weight should be determined as accurately as possible; accuracy of the dosing device should be checked.

If animals are to be treated collectively rather than individually, they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- or overdosing.

Recommended for treatment in sheep 3 weeks post shearing. The product must be applied with a manual or automatic dosing gun, fitted with a spray nozzle, which guarantees the correct spreading of the product on the fleece. Best results will be achieved by holding the gun approximately 45 cm from the sheep during application. Apply as a fan spray along the spine of the animal in bands at least 10 cm wide from the middle of the shoulders and in an arc around the crutch and tail. Half the dose should be applied along the spine with the remainder over the tail and crutch area.

CLiKZiN should be administered before or at the start of predicted fly activity but is also suitable for use during the fly season. The product will protect against fly strike for 8 weeks. It is good agricultural practice to check animals regularly for fly strike.

Please refer to section “Special Warnings” for further advice on administration.

8. WITHDRAWAL PERIOD

Meat and offal: 7 days

Not authorised for use in animals producing milk for human consumption.

9. SPECIAL WARNING(S), IF NECESSARY

For external use only.

If you notice any serious effects or other effects not mentioned on this label, please inform your veterinary surgeon.

Special warnings for each target species

CLiKZiN is best applied before an anticipated blowfly challenge, or when a blowfly challenge is identified on or in the vicinity of the farm. Established strikes may require a separate treatment with a knockdown insecticide.

It is recommended that animals with dirty back-ends are dagged or crutched prior to application. If dagging or crutching is undertaken in the weeks following application, these animals should be re-treated, otherwise protection could be lost.

Do not apply during heavy rainfall, or when such conditions are expected. The resulting wash out may reduce the protection period.

Shake the container well before use.

User Warnings

- Redness and irritation may develop after skin or eye contact with the product.
- Contact with skin and eyes should be avoided.
- Personal protective equipment consisting of synthetic rubber gloves and PVC trousers should be worn when handling the product.
- In case of skin contact remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water.
- In case of eye contact, wash immediately with clean water.
- Always wash hands and exposed skin with soap and water after work.
- Do not eat, drink or smoke whilst using the product.

- It is good agricultural practice to minimise handling of sheep after treatment. If you need to handle sheep within 2 months after treatment, wear synthetic rubber gloves and long trousers or coveralls. If sheep are wet wear waterproof trousers.

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

The safety of the product has not been established during pregnancy or lactation. Use only accordingly to the benefit/risk assessment by the responsible veterinarian. Laboratory studies have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects.

Other precautions

Treated sheep **must** be kept away from watercourses for at least one hour after treatment. There is a **serious** risk to aquatic life if this advice is not followed.

The use of the product has harmful effects on dung flies and beetles.

10. EXPIRY DATE

<EXP {month/year}>

Once opened use within 12 months.

11. SPECIAL STORAGE CONDITIONS

Store in the original container.

Keep the container tightly closed, away from food, drink and animal feedstuffs.

Protect from frost.

Protect from direct sunlight.

Do not use after the expiry date stated on the label.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Harmful to aquatic invertebrates. Do not contaminate ponds or other waterways with product or empty containers.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Marketing authorisation holder

Elanco Europe Ltd
Form 2, Bartley Way
Bartley Wood Business Park
Hook
RG27 9XA
United Kingdom

Manufacturer responsible for batch release

Elanco France S.A.S
26 Rue de la Chapelle
68330 Huingue
France

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

16. MARKETING AUTHORISATION NUMBER

Vm 00879/4071

17. MANUFACTURER'S BATCH NUMBER

<Batch> <Lot> <BN> {number}

Approved: 26/04/21

