

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Frusedale 40 mg Oral Tablets

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each tablet contains:

#### Active ingredient

Furosemide (Frusemide) 40 mg

For a full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Tablets.

White, circular, biconvex, flat-faced tablets with bevelled edges and a breakline.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Cats and dogs.

#### **4.2 Indications for use, specifying the target species**

For the treatment of oedema associated with cardiac insufficiency, renal dysfunction and trauma in cats and dogs.

In animals with pulmonary oedema of cardiac origin, combined therapy with other medicinal products may be indicated.

#### **4.3 Contraindications**

Do not use in animals with acute glomerular nephritis, renal failure with anuria, electrolyte deficiency disease or in animals that have received an overdose of digitalis.

Do not use concurrently with aminoglycoside antibiotics.

Do not use in animals weighing less than 4 kg.

#### **4.4 Special warnings for each target species**

None.

#### **4.5 Special precautions for use**

i. Special precautions for use in animals

Do not exceed the recommended dosage.

Therapeutic efficacy may be impaired by increased intake of drinking water. Where the animal's condition permits, water intake should be restricted during treatment with Frusedale 40 mg oral tablets.

Monitoring of plasma potassium levels is advisable during periods of prolonged treatment of combined therapy with cardiac glycosides. Potassium supplements may be necessary.

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wear gloves, or wash your hands immediately after handling the tablets.

iii. Other precautions

None.

#### **4.6 Adverse reactions (frequency and seriousness)**

During the post-authorisation period and following use of the product in dogs displaying signs of cardiac disease, lethargy, malaise and a fast or irregular heartbeat have been recorded rarely.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

#### **4.7 Use during pregnancy, lactation or lay**

Frusedale 40 mg oral tablets are not contraindicated in pregnant or lactating animals.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

Concurrent administration with corticosteroids may increase the risk of hypokalaemia.

Concurrent administration with aminoglycoside antibiotics may result in ototoxicity.

Concurrent administration with cephalosporin antibiotics may result in nephrotoxicity.

Concurrent administration with digoxin enhances the cardiac glycoside.

Concurrent administration with sulphonamide antibacterials may result in sulphonamide allergy.

#### **4.9 Amounts to be administered and administration route**

For oral administration only. Cats and dogs:

Dosage up to 5 mg/kg bodyweight, or one Frusedale 40 mg oral tablet, per 8 kg bodyweight, one to two times daily with an interval of 6 - 8 hours between administrations.

For maintenance, the dosage should be reduced to 1 - 2 mg/kg per day.

For animals weighing between 4 and 8 kg, one half of one tablet should be administered. The tablets may be divided by breaking along the score line.

Not to be used in animals under 4 kg bodyweight.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

Transitory deafness may occur in animals administered the product at doses higher than those recommended.

Cardiovascular side effects may occur in weak and old animals following overdosage.

Excessive doses can lead to hypovolaemia and decompensate renal function.

Management of signs of overdosage is symptomatic.

#### **4.11 Withdrawal periods**

Not applicable.

## **5. PHARMACOLOGICAL PARTICULARS**

Pharmacotherapeutic group: Furosemide

ATC Vet Code: QC03CA01

### **5.1 Pharmacodynamic properties**

Frusedale 40 mg oral tablets contain Furosemide. Furosemide is a potent loop diuretic with a rapid action.

### **5.2 Pharmacokinetic properties**

Furosemide inhibits electrolyte resorption in the proximal and distal renal tubules and in the ascending Loop of Henle. Excretion of sodium, potassium and chloride ions is enhanced, and also water excretion is enhanced. Furosemide has no effect on carbonic anhydrase. Diuretic activity begins one hour after oral administration and continues for four hours. The potency ensures diuretic action even when renal function is poor. Loop diuretics may cause severe potassium loss.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Maize starch  
Pregelatinised maize starch  
Magnesium stearate  
Lactose monohydrate

### **6.2 Incompatibilities**

There are no known incompatibilities.

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 4 years.

### **6.4 Special precautions for storage**

Do not store above 25°C.  
Protect from light.

### **6.5 Nature and contents of immediate packaging**

Polypropylene containers containing 1000 white, circular, biconvex, flat-faced tablets with bevelled edges, a breakline and which are embossed F40 on one face, CP or DP on the reverse.

**6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products, if appropriate**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

Dechra Limited  
Snaygill Industrial Estate  
Keighley Road  
Skipton  
North Yorkshire  
BD23 2RW  
United Kingdom

**8. MARKETING AUTHORISATION NUMBER**

Vm 10434/4033

**9. DATE OF FIRST AUTHORISATION**

14 August 1998

**10. DATE OF REVISION OF THE TEXT**

March 2022

Approved 03 March 2022

A handwritten signature in black ink, appearing to read "A. Hunter.", is positioned below the approval date.