

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Equipalazone 1 g Oral Powder for Horses and Ponies

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Active substance: per sachet

Phenylbutazone 1 g

Excipient(s)

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Oral powder.
White/cream powder.

4. CLINICAL PARTICULARS

4.1 Target species

Horses and ponies (non-food producing).

4.2 Indications for use, specifying the target species

For the treatment of musculoskeletal disorders in horses and ponies where the anti-inflammatory and analgesic properties of phenylbutazone can offer relief. Examples of conditions normally considered suitable for treatment with phenylbutazone include lameness associated with osteoarthritic conditions, acute and chronic laminitis, bursitis and carpalis, and in the reduction of post-surgical soft tissue reaction.

4.3 Contraindications

Do not administer with other non-steroidal anti-inflammatory drugs (NSAIDs) concurrently or within 24 hours of each other.
Do not use in animals suffering from cardiac, hepatic or renal disease; where there is the possibility of gastrointestinal ulceration or bleeding or where there is evidence of a blood dyscrasia.
Do not use in cases of known hypersensitivity to the active substance or to any of the excipients.

4.4 Special warnings for each target species

The clinical effect of phenylbutazone can be evident for at least three days following cessation of administration. This should be borne in mind when examining horses for soundness.

4.5 Special precautions for use

Special precautions for use in animals

The therapeutic index of phenylbutazone is low. Do not exceed the stated dose or the duration of treatment.
Use in any animal under six weeks of age, or in aged animals, may involve additional risks. If such use cannot be avoided, animals may require a reduced dosage and special clinical management.
Avoid use in any dehydrated, hypovolaemic or hypotensive animal as there is a risk of increased toxicity.
It is preferable that NSAIDs which inhibit prostaglandin synthesis are not administered to animals undergoing general anaesthesia until fully recovered. Response to long-term therapy should be monitored at regular intervals by a veterinary practitioner.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

The product may cause hypersensitivity (allergic) reactions in those sensitized to phenylbutazone, either via skin contact or accidental inhalation. People with known hypersensitivity to phenylbutazone, or any of the excipients, should avoid contact with this product.
If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes, or difficulty breathing, are more serious symptoms and require urgent medical attention.
This product can be irritating to the skin and eyes. Avoid contact with the eyes. In case of accidental eye contact, rinse eyes with plenty of water. If irritation persists seek medical advice. Wash any exposed skin and hands after use. Care should be taken to avoid ingesting the powder. In the event of accidental ingestion, seek medical advice and show the product packaging to the physician.
The safety of phenylbutazone in pregnancy has not been established. The veterinary medicinal product should not be administered by pregnant women or women attempting to conceive.

4.6 Adverse reactions (frequency and seriousness)

In common with other NSAIDs that inhibit prostaglandin synthesis, there may be gastric and/or renal intolerance. This is usually associated with overdose and such events are rare. Recovery is usual on cessation of treatment and following the initiation of supportive symptomatic therapy (see 4.10 for further information).

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports)

4.7 Use during pregnancy, lactation or lay

The safety of phenylbutazone in pregnancy and lactation has not been established. Use during pregnancy should be avoided whenever possible, particularly during the first trimester.

Use phenylbutazone in pregnant and lactating mares only according to a benefit/risk assessment by the responsible veterinarian.

4.8 Interaction with other medicinal products and other forms of interaction

Some NSAIDs may be highly bound to plasma proteins and compete with other highly bound drugs to produce an increase in non-bound pharmacologically active concentrations which can lead to toxic effects. Concurrent administration of potential nephrotoxic drugs (e.g. aminoglycoside antibiotics) should be avoided.

Adverse reactions caused by phenylbutazone are exacerbated by concurrent administration of glucocorticoids, other non-steroidal antiphlogistics, or anticoagulants.

Gastrointestinal tract ulceration may be exacerbated by corticosteroids in animals given NSAIDs.

4.9 Amounts to be administered and administration route

For oral administration only. When mixed with a concentrate feed, the product was shown to be palatable to horses.

The dosage should be adjusted according to the individual animal's response, but the following may be taken as a guide:

Horses 450 kg (1000 lb) body weight: the contents of two sachets to be administered twice on day 1 of treatment (equivalent to 8.8 mg/kg/day) followed by the contents of one sachet twice daily for four days (4.4 mg/kg/day), then one sachet daily, or on alternate days, sufficient to keep the horse comfortable (2.2 mg/kg/day).

Ponies 225 kg (500 lb) body weight, one sachet (4.4 mg/kg) on alternate days.

Discontinue treatment if no response is evident after four to five days treatment.

For ease of administration mix the powder with a small quantity of feed.

Dampening of the veterinary medicinal product in feed 5 minutes prior to feeding has been shown to have no detrimental influence on the palatability of the product. However, the influence of prolonged dampening on palatability or stability of the product is not known.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Overdosing may result in gastric and large intestinal ulceration and general enteropathy. Renal papillary damage may also occur with impaired renal function. Subcutaneous oedema, especially under the jaw may become evident due to plasma protein loss.

There is no specific antidote. If signs of possible overdosage occur, treat the animal symptomatically.

The therapeutic index of phenylbutazone is low. In man, charcoal haemoperfusion in conjunction with dopamine has been used successfully to treat overdosage with phenylbutazone, but there is no experience of the use of this technique in the horse.

4.11 Withdrawal periods

Not for use in horses intended for human consumption.

Treated horses may never be slaughtered for human consumption.

The horse must have been declared as not intended for human consumption under national horse passport legislation.

5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Anti-inflammatory and antirheumatic products, non-steroids.

ATC Vet Code: QM01AA01.

5.1 Pharmacodynamic properties

Phenylbutazone is a pyrazolone non-steroidal anti-inflammatory drug (NSAID) which acts by the nonselective inhibition of prostaglandin synthetases (cyclooxygenases COX-1 and COX-2). Prostaglandins possess a wide variety of physiological properties, including those involved in the production of pain, inflammation and pyrexia. The main metabolite, oxyphenbutazone, possesses similar pharmacological properties.

5.2 Pharmacokinetic particulars

Phenylbutazone is generally well absorbed following oral administration. The rate, but not the extent, of absorption may be affected due to binding of phenylbutazone to food and the contents of the gastrointestinal tract.

Therefore, it is recommended that Equipalazone Powder is administered mixed with a small amount of bran or oats. Phenylbutazone is highly bound to plasma proteins.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Acacia
Gelatin
Silicon dioxide
Sucralose
Apple flavour

6.2 Major incompatibilities

None known.

6.3 Shelf life

Shelf life of veterinary medicinal product as packaged for sale: 4 years.

6.4 Special precautions for storage

Do not store above 25°C.
Store in a dry place.

6.5 Nature and composition of immediate packaging

Sachets of a paper/polyethylene outer layer and aluminium/polyethylene inner layer, in packs of 100 sachets (25 strips of four sachets) and of 32 sachets (8 strips of four sachets). Each sachet contains 1.5 g Equipalazone Powder. Not all pack sizes may be marketed.

6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Dechra Limited
Snaygill Industrial Estate
Keighley Road
Skipton
North Yorkshire
BD23 2RW
United Kingdom

8. MARKETING AUTHORISATION NUMBER


Vm 10434/4090

9. DATE OF FIRST AUTHORISATION

15 June 2017

10. DATE OF REVISION OF THE TEXT

October 2022

A handwritten signature in black ink, consisting of several loops and a final horizontal stroke.

Approved: 05 October 2022