

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Drontal Plus XL Flavour Tablets for Dogs

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each tablet contains:	mg per tablet
Febantel	525.0
Pyrantel embonate	504.0
Praziquantel	175.0

#### **Excipients**

Artificial beef flavour Irradiated	408.0
------------------------------------	-------

For the full list of excipients, see section 6.1

### **3. PHARMACEUTICAL FORM**

Tablet.

Pale brown to brown oval shaped divisible tablet scored on both sides.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Dogs.

#### **4.2 Indications for use, specifying the target species**

For the control of the following roundworms and tapeworms in adult dogs:

Ascarids: *Toxocara canis*, *Toxascaris leonina* (adult and late immature forms).

Hookworms: *Uncinaria stenocephala*, *Ancylostoma caninum* (adults)

Whipworms: *Trichuris vulpis* (adults)

Tapeworms: *Echinococcus* spp. *Taenia* spp. and *Dipylidium caninum* (adult and immature forms).

### **4.3 Contraindications**

Do not use simultaneously with piperazine compounds.

### **4.4 Special warnings for each target species**

Fleas serve as intermediate hosts for one common type of tapeworm - *Dipylidium caninum*. Tapeworm infestation is certain to re-occur unless control of intermediate hosts such as fleas, mice etc. is undertaken.

Since it contains praziquantel, the product is effective against *Echinococcus multilocularis*, which does not occur in the UK or Ireland but is becoming more common in some European countries. As a precautionary measure to prevent establishment of *E multilocularis* in the UK and Ireland, it is recommended that all dogs entering the country be treated with praziquantel.

### **4.5 Special precautions for use**

#### **i) Special precautions for use in animals**

Any part-used tablets should be discarded.  
Do not exceed the stated dose when treating pregnant bitches.

#### **ii) Special precautions to be taken by the person administering the medicinal product to animals**

In case of accidental ingestion, seek medical advice and show package leaflet to the physician.

In the interests of good hygiene, persons administering the tablet directly to a dog or by adding it to the dog's food, should wash their hands afterwards.

### **4.6 Adverse reactions (frequency and seriousness)**

In very rare cases slight and transient digestive tract disorders such as vomiting and/or diarrhoea may occur. In individual cases these signs can be accompanied by nonspecific signs such as lethargy, anorexia or hyperactivity.

### **4.7 Use during pregnancy and lactation**

Consult a veterinary surgeon before treating pregnant animals for roundworms.

Drontal Plus XL Flavour Tablets may be used during lactation (see Section 4.9 below).

### **4.8 Interaction with other medicinal products and other forms of interaction**

Do not use simultaneously with piperazine.  
Concurrent use with other cholinergic compounds is not recommended.

#### 4.9 Amount(s) to be administered and administration route

##### *Dosage*

The recommended dose rates are: 15 mg/kg bodyweight febantel, 14.4 mg/kg pyrantel and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 35 kg bodyweight as follows:

Dosages are as follows:

Body weight (kg)	Tablet quantity
17.5	½
>17.5 - 35	1
>35 - 52.5	1 ½
>52.5 - 70	2

Drontal Plus Flavour Tablets should be used to achieve accurate dosing in dogs weighing less than 17.5 kg. The dose is equivalent to 1 tablet per 10 kg.

##### *Administration and Duration of Treatment*

Oral administration: the tablet(s) can be given directly to the dog or disguised in food. No starvation is needed before or after treatment.

For routine treatment a single dose is recommended.

For the control of *Toxocara*, nursing bitches should be dosed 2 weeks after giving birth and every 2 weeks until weaning.

For routine control adult dogs should be treated every 3 months. In the event of a heavy roundworm infestation, a repeat dose should be given after 14 days.

#### 4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Drontal Plus XL Flavour Tablets are well tolerated in dogs. In safety studies doses of 5 x or greater gave rise to occasional vomiting.

#### 4.11 Withdrawal period(s)

Not applicable.

### 5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group:  
Anthelmintics, tetrahydropyrimidine

ATC VetCode: QP52AF30

The product contains anthelmintics active against roundworms and tapeworms. The product contains three active substances:

- 1) Febantel, a pro-benzimidazole.
- 2) Pyrantel embonate (pamoate), a tetrahydropyrimidine derivative, and
- 3) Praziquantel, a partially hydrogenated pyrazino-isoquinoline derivative.

## 5.1 Pharmacodynamic Properties

In this fixed combination product pyrantel and febantel act synergistically against relevant nematodes (ascarids, hookworms and whipworms) in dogs. In particular, the activity spectrum covers *Toxocara canis*, *Toxascaris leonina*, *Uncinaria stenocephala*, *Ancylostoma caninum* and *Trichuris vulpis*.

The spectrum of activity of praziquantel covers all important cestode species in dogs, in particular all *Taenia* spp, *Dipylidium caninum*, *Echinococcus granulosus* and *Echinococcus multilocularis*. Praziquantel acts against adult and immature forms of these parasites.

Praziquantel is very rapidly absorbed and distributed throughout the parasite. Both *in vivo* and *in vitro* studies have shown that praziquantel causes severe damage to the parasite integument, resulting in contraction and paralysis. There is an almost instantaneous tetanic contraction of the parasite musculature and a rapid vacuolisation of the syncytial tegument. This rapid contraction has been explained by changes in divalent cation fluxes, especially calcium.

Pyrantel acts as a cholinergic agonist. Its mode of action is to stimulate nicotinic cholinergic receptors of the parasite, induce spastic paralysis and thereby allow removal from the gastro-intestinal (GI) system by peristalsis.

Within the mammalian system febantel undergoes ring closure forming fenbendazole and oxfendazole. It is these chemical entities which exert the anthelmintic effect by inhibition of tubulin polymerisation. Formation of microtubules is thereby prevented, resulting in disruption to structures vital to the normal functioning of the helminth. Glucose uptake, in particular, is affected, leading to a depletion in cell ATP. The parasite dies upon exhaustion of its energy reserves, which occurs 2-3 days later.

## 5.2 Pharmacokinetic Properties

No data available.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

Artificial beef flavour Irradiated  
Maize starch  
Lactose monohydrate  
Microcrystalline cellulose  
Povidone K25  
Magnesium stearate

Sodium laurilsulfate  
Silica colloidal anhydrous

## **6.2 Incompatibilities**

None known.

## **6.3 Shelf-life**

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.

## **6.4 Special precautions for storage**

Do not store above 25°C.  
Any part used tablets should be discarded

## **6.5 Nature and composition of immediate packaging**

Container material: PCTFE/PVC-aluminium foil strip

Container colour: White

Container sizes: Cartons containing 2, 8, 48 and 96 tablets  
Not all pack sizes may be marketed.

## **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products, if appropriate**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Vetoquinol UK Limited  
Steadings Barn  
Pury Hill Business Park  
Nr. Alderton  
Towcester  
Northamptonshire  
NN12 7LS

## **8. MARKETING AUTHORISATION NUMBER**

Vm 08007/4169

**9. DATE OF FIRST AUTHORISATION**

31 March 2008

**10. DATE OF LAST REVISION OF THE TEXT**

November 2020

Approved 05 November 2020

A handwritten signature in black ink, appearing to read "Hunter.", is positioned below the approval date. The signature is stylized and written in a cursive-like font.