

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Bob Martin Clear Spot on 268 mg spot-on solution for large dogs

UK: Bob Martin Fipronil 268 mg spot-on solution for large dogs

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

One 2.68 ml pipette contains:

#### **Active substance:**

Fipronil 268 mg

#### **Excipients:**

Butylhydroxyanisole E320 0.536 mg

Butylhydroxytoluene E321 0.268 mg

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Spot-on solution.

Clear, pale amber solution.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Dogs.

#### **4.2 Indications for use, specifying the target species**

Treatment of flea (*Ctenocephalides spp.*) and tick (*Rhipicephalus sanguineus* and *Ixodes ricinus*) infestations.

Fleas will be killed within 24 h. Insecticidal efficacy against new infestations with adult fleas persists for 8 weeks.

The product has a persistent acaricidal efficacy for 4 weeks against ticks (*Rhipicephalus sanguineus*, *Ixodes ricinus*, *Dermacentor reticulatus*).

Ticks will usually be killed within 48 h after contact with Fipronil. However, if ticks of some species (*Dermacentor reticulatus*) are already present when the product is applied, all of the ticks may not be killed within the first 48 hours.

The product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD), where this has been previously diagnosed by a veterinary surgeon.

### **4.3 Contraindications**

Do not use on puppies less than 2 months old and /or weighing less than 2kg. Do not use on sick (e.g. systemic diseases, fever...) or convalescent animals.

Do not use in rabbits, as adverse reactions and even death could occur.

This product is specifically developed for dogs. Do not use in cats, as this could lead to overdosing.

Do not use on animals with known hypersensitivity to the active substance or to any of the excipients.

### **4.4 Special warnings for each target species**

Avoid frequent swimming or shampooing the animal because the maintenance of effectiveness of the product in these cases has not been tested.

The product does not prevent ticks from attaching to the animals. If the animal has been treated prior to exposure to the ticks, the ticks will be killed in the first 24-48 hours after attachment. This will usually be prior to engorgement, minimising but not excluding the risk of transmission of diseases. Once dead, ticks will often drop off the animal, but any remaining ticks may be removed with a gentle twist.

Fleas from pets often infest the animal's basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated, in case of massive infestation and at the beginning of the control measures, with a suitable insecticide and vacuumed regularly.

For optimal control of flea infestation in multi-pet household, all dogs and cats in the household should be treated with a suitable insecticide.

When used as part of a strategy for the treatment of Flea Allergy Dermatitis, monthly applications to the allergic patient and to other dogs and cats in the household are recommended.

### **4.5 Special precautions for use**

#### **i) Special precautions for use in animals**

Animals should be weighed accurately prior to treatment.

Avoid contact with the animal's eyes. In the case of accidental eye contact, immediately and thoroughly flush the eyes with water.

It is important to make sure that the product is applied to an area where the animal cannot lick it off, and to make sure that animals do not lick each other following treatment.

Do not apply the product on wounds or damaged skin.

## **ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals**

This product can cause mucous membrane and eye irritation. Therefore, contact between the product and the mouth or eyes should be avoided.

In the case of accidental eye contact, immediately and thoroughly flush the eyes with water. If eye irritation persists seek medical advice and show the package leaflet or the label to the physician.

Do not smoke, drink or eat during application.

Avoid contents coming into contact with the fingers. If this occurs, wash hands with soap and water. Wash hands after use.

Animals or people with a known hypersensitivity to fipronil or any of the other ingredients should avoid contact with the product.

Treated animals should not be handled until the application site is dry, and children should not be allowed to play with treated animals until the application site is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals should not be allowed to sleep with owners, especially children.

Keep pipettes in the original packaging and dispose of used pipettes immediately.

## **iii) Environmental warning**

Fipronil may adversely affect aquatic organisms. Dogs should not be allowed to swim in water courses for 2 days after application.

## **iv) Other precautions**

The product may have adverse effects on painted, varnished or other household surfaces or furnishings.

## **4.6 Adverse reactions (frequency and seriousness)**

If licking occurs, a brief period of hypersalivation may be observed due mainly to the nature of the carrier.

Among the extremely rare suspected adverse reactions, transient cutaneous reactions on the application site (skin discoloration, local alopecia, pruritus, erythema) and general pruritus or alopecia have been reported after use.

Exceptionally, hypersalivation, reversible neurologic symptoms (hyperesthesia, depression, nervous symptoms), vomiting or respiratory symptoms have been observed after use.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals displaying adverse reactions during the course of one treatment)
- common (more than 1 but less than 10 animals in 100 animals)
- uncommon (more than 1 but less than 10 animals in 1,000 animals)
- rare (more than 1 but less than 10 animals in 10,000 animals)
- very rare (less than 1 animal in 10,000 animals, including isolated reports).

## **4.7 Use during pregnancy and lactation**

Laboratory studies using fipronil have not shown any evidence of teratogenic or embryotoxic effect. Studies have not been carried out with this product in pregnant

and lactating bitches. Use in pregnancy and lactation only in accordance with professional veterinary advice and a benefit/risk assessment.

#### 4.8 Interaction with other medicinal products and other forms of interaction

None known.

#### 4.9 Amounts to be administered and administration route

##### Route of administration and dosage:

External use only. Administer by topical application to the skin according to the bodyweight as follows: 1 pipette of 2.68 ml per dog weighing over 20 kg and up to 40 kg bodyweight.

For dogs over 60 kg, use two pipettes of 2.68 ml.

##### Method of administration:



Remove the pipette from the blister packaging. Hold the pipette in an upright position and snap off the top section of the cap, thereby opening the pipette.

Part your dog's coat until the skin is visible. Place the tip of the pipette directly against the bared skin and squeeze gently. Empty approximately half the contents at each of two points along the dog's back, preferably at the base of the head and between the shoulder blades. Squeeze the pipette several times to ensure dosing is complete. Avoid applying the solution onto the fur and do not rub into the skin.

Application of the solution as directed minimises the possibility that the animal will lick the solution off. Do not allow pets to lick each other after treatment.

Care should be taken to avoid excessive wetting of the hair with the product since this will cause a sticky appearance of hairs at the treatment spot. However, should this occur, it will usually disappear within 24-48 hours post application. Crystals may be seen on the hair and slight scaling may occur within 24-48 hours after application.

##### Treatment schedule:

For optimal control of flea and/or tick infestation the treatment schedule can be based on the local epidemiological situation.

The minimum interval between two treatments is 4 weeks.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

No adverse effects were observed in target animal safety studies in 2 month-old puppies, growing dogs and dogs weighing about 2 kg treated with 5 times the therapeutic dose once a month for 3 consecutive months. The risk of adverse effects (see section 4.6) may increase in cases of overdose.

#### **4.11 Withdrawal period(s)**

Not applicable.

### **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic group: Ectoparasiticides for topical use.  
ATCvet code: QP53AX15

#### **5.1 Pharmacodynamic properties**

Fipronil is an insecticide and acaricide belonging to the phenylpyrazole family. It acts by inhibiting the GABA complex, binding to the chloride channel and thereby blocking pre- and post-synaptic transfer of chloride ions across cell membranes. This results in uncontrolled activity of the central nervous system and death of insects or acarids. Fipronil exhibits an insecticidal and acaricidal activity against fleas (*Ctenocephalides spp.*), ticks (*Rhipicephalus spp.*, *Ixodes spp.* including *Ixodes ricinus*) in the dog.

#### **5.2 Pharmacokinetic particulars**

Fipronil is mainly metabolised to its sulfone derivative (RM1602), which also possesses insecticidal and acaricidal properties. The concentrations of fipronil on the hair decrease with time.

### **6. PHARMACEUTICAL PARTICULARS**

#### **6.1 List of excipients**

Butylhydroxyanisole E320  
Butylhydroxytoluene E321  
Benzyl alcohol E1519  
Diethylene glycol monoethyl ether

#### **6.2 Incompatibilities**

None known.

#### **6.3 Shelf life**

Shelf-life of the veterinary medicinal product as packaged for sale: 2 years.

#### **6.4 Special precautions for storage**

Store below 25°C, in a dry place in the original packaging.

## **6.5 Nature and composition of immediate packaging**

White opaque or purple or pink translucent polypropylene single-dose pipettes containing an extractable volume of 2.68 ml packaged in a clear PVC blister closed by heat sealing with aluminium foil and placed in a carton box or blister card. Blister cards or boxes of 1, 2, 3, 4, 5 or 6 pipettes.

Not all pack sizes may be marketed.

## **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements. Fipronil may adversely affect aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty container.

## **7. MARKETING AUTHORISATION HOLDER**

Pets Choice Limited  
Brentwood House  
Lower Philips Road  
Whitebirk Industrial Estate  
Blackburn  
Lancashire  
BB1 5UD  
United Kingdom

## **8. MARKETING AUTHORISATION NUMBER**


Vm 52797/4007

## **9. DATE OF FIRST AUTHORISATION**

15 December 2011

## **10. DATE OF REVISION OF THE TEXT**

May 2020



Approved: 29 May 2020