

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Canigen Bb suspension for injection in pre-filled syringe for dogs

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each dose (1 ml) contains:

Active substance:

Bordetella bronchiseptica fimbriae¹: 88 - 399 U²

¹ Purified from strain Bb7 92932

² Antigenic mass ELISA units

Adjuvant:

dl- α -tocopheryl acetate: 74.7 mg

Excipients:

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Suspension for injection in pre-filled syringe.
Aqueous, white to nearly white suspension, mild creaming.

4. CLINICAL PARTICULARS

4.1 Target species

Dogs.

4.2 Indications for use, specifying the target species

For active immunisation of dogs against *Bordetella bronchiseptica* to reduce clinical signs of upper respiratory tract disease and bacterial shedding post infection.

Onset of immunity: 2 weeks.

Duration of immunity: 7 months after primary vaccination.
1 year after revaccination.

4.3 Contraindications

None.

4.4 Special warnings for each target species

Vaccinate healthy animals only.

4.5 Special precautions for use

Special precautions for use in animals

Not applicable.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Not applicable.

Special precautions for the protection of the environment

Not applicable.

Other precautions

Not applicable.

4.6 Adverse reactions (frequency and seriousness)

Dogs:

| | |
|---|--|
| Very common (> 1 animal / 10 animals treated): | Injection site swelling (\leq 2 cm, occasionally firm, may be present up to 25 days post-vaccination). |
| Common (1 to 10 animals / 100 animals treated): | Injection site swelling (\leq 3.5 cm, may be present up to 25 days post-vaccination ¹ and can be painful). |
| Very rare (< 1 animal / 10,000 animals treated, including isolated reports): | Hypersensitivity reaction. ² |

¹ The swelling may uncommonly last for up to 35 days post-vaccination.

² If hypersensitivity reaction occurs, appropriate treatment should be administered without delay. Such reactions may evolve to a more severe condition which may be life-threatening.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

4.7 Use during pregnancy, lactation or lay

Pregnancy:

Can be used during pregnancy. The safety of this vaccine has not been investigated during the first 20 days of gestation.

4.8 Interaction with other medicinal products and other forms of interaction

Safety and efficacy data are available which demonstrate that this vaccine can be administered at the same time but not mixed with the live vaccines in the Canigen range against canine distemper, canine contagious hepatitis caused by canine adenovirus type 1, canine parvovirus disease and respiratory disease caused by canine adenovirus type 2, where authorised.

Safety data are available which demonstrate that this vaccine can be administered at the same time but not mixed with the Canigen range of vaccines mentioned above together with the live Canigen parainfluenza vaccine and the inactivated vaccines in the Canigen range against leptospirosis caused by *L. interrogans* serogroup Canicola serovar Canicola, *L. interrogans* serogroup Icterohaemorrhagiae serovar Copenhageni, *L. interrogans* serogroup Australis serovar Bratislava, and *L. kirschneri* serogroup Grippotyphosa serovar Bananal/Lianguang.

In addition, for the live canine parainfluenza vaccine there are antibody response data, and for the inactivated canine leptospirosis vaccines there are antibody response data and other immunity data which support the use of the vaccine at the same time but not mixed with the mentioned Canigen range of vaccines.

When this vaccine is administered in association with the relevant Canigen vaccines, the demonstrated safety and efficacy claims of the vaccine are the same as when this vaccine is administered alone.

The product information of the relevant Canigen vaccines used in association with this vaccine should be consulted before administration.

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product except the products mentioned above. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

4.9 Amount(s) to be administered and administration route

Subcutaneous use, 1 ml dose per vaccination.

Dogs can be vaccinated from the age of 6 weeks onwards.

Allow the vaccine to reach room temperature (15 °C – 25 °C) before use.

Primary vaccination:

Two vaccinations with an interval of 4 weeks.

Revaccination:

A single vaccination, administered 7 months after primary vaccination with this vaccine, is sufficient to maintain protection against *Bordetella bronchiseptica* for a further year. Thereafter, a single vaccination should be administered, annually. In case revaccination at 7 months is missed, a single vaccination within 12 months after primary vaccination is sufficient to extend protection against *Bordetella bronchiseptica* for a further year.

This vaccine can also be used for revaccination in a schedule where Canigen KC has been used for primary vaccination. A single vaccination, administered one year after

primary vaccination with Canigen KC, is sufficient to prolong immunity against *Bordetella bronchiseptica* for another year.

Revaccination after primary vaccination with Canigen KC:

One vaccination, annually.

For associated use:

When this vaccine is administered in associated use (i.e. not mixed) with another vaccine in the Canigen range as indicated under section 4.8, the vaccines should be given subcutaneously at the same time, at a different site. Dogs should not be younger than the minimum age recommended for the other Canigen vaccine, as stated in the respective product information.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Not applicable.

4.11 Withdrawal period(s)

Not applicable.

5. IMMUNOLOGICAL PROPERTIES

Pharmacotherapeutic group: Immunologicals for *Canidae*, inactivated bacterial vaccines (including mycoplasma, toxoid and chlamydia)

ATCvet code: QI07AB03.

The subunit vaccine stimulates active immunity against *Bordetella bronchiseptica* infection in dogs.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

dl- α -tocopheryl acetate
Sodium chloride
Disodium hydrogen phosphate dihydrate
Sodium dihydrogen phosphate dihydrate
Polysorbate 80
Water for injections

6.2 Major incompatibilities

Do not mix with any other veterinary medicinal product.

6.3 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

6.4 Special precautions for storage

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Store in the original package in order to protect from light.

6.5 Nature and composition of immediate packaging

Type I glass pre-filled syringe, containing a plunger with a halogenobutyl end and closed with a halogenobutyl stopper.

Pack sizes:

Cardboard box with:

5 single dose pre-filled syringes (5 x 1 ml) and needles.

10 single dose pre-filled syringes (10 x 1 ml) and needles.

Not all pack sizes may be marketed.

6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

MSD Animal Health UK Limited
Walton Manor
Walton
Milton Keynes
MK7 7AJ

8. MARKETING AUTHORISATION NUMBER

Vm 01708/5064

9. DATE OF FIRST AUTHORISATION

23 November 2021

10. DATE OF REVISION OF THE TEXT

December 2023


PROHIBITION OF SALE, SUPPLY AND/OR USE

Not applicable.

11. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCT

Veterinary medicinal product subject to prescription.

Find more product information by searching for the 'Product Information Database' or 'PID' on www.gov.uk.

A handwritten signature in black ink, consisting of several loops and a final flourish.

Approved: 28 April 2024