

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Metacam 40 mg/ml solution for injection for cattle and horses

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One ml contains:

Active substance:

Meloxicam 40 mg

Excipient:

Ethanol 150 mg

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for injection.

Clear yellow solution.

4. CLINICAL PARTICULARS

4.1 Target species

Cattle and horses

4.2 Indications for use, specifying the target species

Cattle:

For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in cattle.

For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in calves of over one week of age and young, non-lactating cattle.

For adjunctive therapy in the treatment of acute mastitis, in combination with antibiotic therapy.

For the relief of post-operative pain following dehorning in calves.

Horses:

For use in the alleviation of inflammation and relief of pain in both acute and chronic musculo-skeletal disorders.

For the relief of pain associated with equine colic.

4.3 Contraindications

Do not use in pregnant or lactating mares (see section 4.7).

Do not use in horses less than 6 weeks of age.

Do not use in animals suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

For the treatment of diarrhoea in cattle, do not use in animals of less than one week of age.

4.4 Special warnings for each target species

Treatment of calves with Metacam 20 minutes before dehorning reduces post-operative pain. Metacam

alone will not provide adequate pain relief during the dehorning procedure. To obtain adequate pain relief during surgery co-medication with an appropriate analgesic is needed.

4.5 Special precautions for use

Special precautions for use in animals

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

Avoid use in very severely dehydrated, hypovolaemic or hypotensive animals which require parenteral rehydration, as there may be a potential risk of renal toxicity.

In case of inadequate relief of pain when used in the treatment of equine colic, careful re-evaluation of the diagnosis should be made as this could indicate the need for surgical intervention.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Accidental self-injection may give rise to pain. People with known hypersensitivity to non-steroidal anti-inflammatory drugs (NSAIDs) should avoid contact with the veterinary medicinal product.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

In view of the risk of accidental self-injection and the known adverse class-effects of NSAIDs and other prostaglandin inhibitors on pregnancy and/or embryofetal development, the veterinary medicinal product should not be administered by pregnant women or women attempting to conceive.

This product can cause eye irritation. In case of contact with the eyes, immediately rinse thoroughly with water.

4.6 Adverse reactions (frequency and seriousness)

In cattle, only a slight transient swelling at the injection site following subcutaneous administration was observed in less than 10 % of the cattle treated in clinical studies.

In horses, a transient swelling at the injection site was observed in isolated cases in clinical studies, but resolved without intervention.

Anaphylactoid reactions, which may be serious (including fatal), have been observed very rarely from post-marketing safety experience and should be treated symptomatically.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reactions)
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

4.7 Use during pregnancy, lactation or lay

Cattle: Can be used during pregnancy and lactation.

Horses: Do not use in pregnant or lactating mares (see section 4.3).

4.8 Interaction with other medicinal products and other forms of interaction

Do not administer concurrently with glucocorticosteroids, other NSAIDs or with anticoagulant agents.

4.9 Amounts to be administered and administration route

Cattle:

Single subcutaneous or intravenous injection at a dose of 0.5 mg meloxicam/kg body weight (i.e.

1.25 ml/100 kg body weight) in combination with antibiotic therapy or with oral re-hydration therapy, as appropriate.

Horses:

Single intravenous injection at a dose of 0.6 mg meloxicam/kg body weight (i.e. 1.5 ml/100 kg body weight).

For use in the alleviation of inflammation and the relief of pain in both acute and chronic musculo-skeletal disorders, Metacam 15 mg/ml oral suspension may be used for continuation of treatment at a dose of 0.6 mg meloxicam/kg body weight, 24 hours after administration of the injection.

Avoid introduction of contamination during use.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

In case of overdose symptomatic treatment should be initiated.

4.11 Withdrawal period(s)

Cattle: Meat and offal: 15 days; milk: 5 days.

Horses: Meat and offal: 5 days.

Not authorised for use in horses producing milk for human consumption.

5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Anti-inflammatory and antirheumatic products, non-steroids (oxicams).
ATCvet code: QM01AC06.

5.1 Pharmacodynamic properties

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) of the oxicam class which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, anti-exudative, analgesic and antipyretic effects. It reduces leukocyte infiltration into the inflamed tissue. To a minor extent it also inhibits collagen-induced thrombocyte aggregation. Meloxicam also has anti-endotoxic properties because it has been shown to inhibit production of thromboxane B₂ induced by *E. coli* endotoxin administration in calves and lactating cows.

5.2 Pharmacokinetic particulars

Absorption

After a single subcutaneous dose of 0.5 mg meloxicam/kg, C_{max} values of 2.1 µg/ml and 2.7 µg/ml were reached after 7.7 hours and 4 hours in young cattle and lactating cows, respectively.

Distribution

More than 98 % of meloxicam is bound to plasma proteins. The highest meloxicam concentrations are to be found in liver and kidney. Comparatively low concentrations are detectable in skeletal muscle and fat.

Metabolism

Meloxicam is predominantly found in plasma. In cattle, meloxicam is also a major excretion product in milk and bile whereas urine contains only traces of the parent compound. Meloxicam is metabolised to an alcohol, an acid derivative and to several polar metabolites. All major metabolites have been shown to be pharmacologically inactive. The metabolism in horses has not been investigated.

Elimination

Meloxicam is eliminated with a half-life of 26 hours and 17.5 hours after subcutaneous injection in young

cattle and lactating cows, respectively.

In horses, after intravenous injection meloxicam is eliminated with a terminal half-life of 8.5 hours. Approximately 50 % of the administered dose is eliminated via urine and the remainder via faeces.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Ethanol
Poloxamer 188
Macrogol 300
Glycine
Disodium edetate
Sodium hydroxide
Hydrochloric acid
Meglumine
Water for injections

6.2 Major incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

6.3 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.
Shelf life after first opening the immediate packaging: 28 days.

6.4 Special precautions for storage

This veterinary medicinal product does not require any special storage conditions.

6.5 Nature and composition of immediate packaging

Pack sizes of 1 or 12 colourless glass injection vial(s) each containing 50 ml or 100 ml.
Each vial is closed with a rubber stopper and sealed with an aluminium cap.
Not all pack sizes may be marketed.

6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Boehringer Ingelheim Vetmedica GmbH
55216 Ingelheim/Rhein
GERMANY

8. MARKETING AUTHORISATION NUMBER(S)

Vm 04491/5023

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 07.01.1998
Date of last renewal: 06.12.2007

10. DATE OF REVISION OF THE TEXT

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>.

PROHIBITION OF SALE, SUPPLY AND/OR USE

Not applicable.