

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Vomend 5 mg/ml solution for injection for dogs and cats

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

1 ml contains:

#### **Active substance:**

Metoclopramide (as hydrochloride monohydrate)	4.457 mg
equivalent to metoclopramide hydrochloride	5 mg

#### **Excipient:**

Benzyl alcohol (E1519)	18 mg
------------------------	-------

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Solution for injection.  
Clear colourless aqueous solution.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Dogs and cats.

#### **4.2 Indications for use, specifying the target species**

Symptomatic treatment of vomiting and reduced gastrointestinal motility associated with gastritis, pyloric spasm, chronic nephritis and digestive intolerance to some drugs.

#### **4.3 Contraindications**

Do not use in cases of gastrointestinal perforation or obstruction.

#### **4.4 Special warnings for each target species**

None.

## **4.5 Special precautions for use**

### Special precautions for use in animals

The dosage must be adapted in animals with renal or hepatic insufficiency (due to an increase in the risk of side effects). Avoid administration to animals with epilepsy. The dosage should be carefully observed, especially in cats and small breed dogs. Following prolonged vomiting, consideration should be given to fluid and electrolyte replacement therapy.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash hands after administration to the animal.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

In case of accidental exposure by spillage onto the skin or eyes, wash immediately with abundant water. If an adverse effect occurs, seek medical advice immediately and show the package leaflet or the label to the physician.

## **4.6 Adverse reactions (frequency and seriousness)**

In some very rare cases (less than 1 animal in 10,000 animals, including isolated reports), extrapyramidal effects (agitation, ataxia, abnormal positions and/or movements, prostration, tremors and aggression, vocalisation) have been observed after treatment of dogs and cats.

The observed effects are transient and disappear when treatment is stopped. In very rare cases, allergic reactions may occur.

## **4.7 Use during pregnancy, lactation or lay**

Laboratory studies in laboratory animals have not produced any evidence of teratogenic or foetotoxic effects. However, studies in laboratory animals are limited and the safety of the active substance has not been evaluated in the target species. The use of the product during pregnancy and lactation must be made according to the benefit/risk assessment carried out by the veterinarian.

## **4.8 Interaction with other medicinal products and other forms of interaction**

In cases of gastritis, avoid the co-administration of anticholinergic drugs (atropine) as they may counteract the effects of metoclopramide on gastrointestinal motility.

In cases of simultaneous diarrhoea, there is no contraindication to the use of anticholinergic drugs.

Concurrent use of metoclopramide with neuroleptics derived from phenothiazine (acepromazine) and butyrophenones increases the risk of extrapyramidal effects (see section 4.6).

Metoclopramide can potentiate the action of central nervous system depressants. If used concurrently, it is advised to use the lowest dosage of metoclopramide to avoid excessive sedation.

## **4.9 Amounts to be administered and administration route**

Intramuscular or subcutaneous use.

0.5 mg metoclopramide hydrochloride per kg body weight, if necessary repeated every 6-8 hours.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

Most of the clinical signs reported after an overdose are well known extrapyramidal side effects (see section 4.6).

In the absence of a specific antidote, it is recommended to offer a calm environment to the animal until the extrapyramidal side effects disappear.

Metoclopramide being rapidly metabolised and eliminated, side effects generally disappear quickly.

#### **4.11 Withdrawal period**

Not applicable.

### **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic group: propulsives.  
ATCvet code: QA03FA01

#### **5.1 Pharmacodynamic properties**

Metoclopramide is an original orthopramide molecule.

The anti-emetic action of metoclopramide is mainly due to its antagonist activity at D2 receptors in the central nervous system, preventing nausea and vomiting triggered by most stimuli.

The prokinetic effect on the gastro-duodenal transit (increase in intensity and rhythm of stomach contractions and opening of the pylorus) is mediated by muscarinic activity, D2 receptor antagonist activity and 5-HT<sub>4</sub> receptor agonist activity at the gastrointestinal level.

#### **5.2 Pharmacokinetic particulars**

Metoclopramide is rapidly and completely absorbed after parenteral administration. After subcutaneous administration to dogs and cats, maximum concentrations are obtained after 15-30 minutes.

Metoclopramide is rapidly distributed into most tissues and fluids, crosses the blood-brain barrier and enters the central nervous system.

Metoclopramide is metabolised by the liver.

The elimination of metoclopramide is rapid, 65% of the dose being eliminated within 24 hours in the dog, primarily by the urinary route.

### **6. PHARMACEUTICAL PARTICULARS**

#### **6.1 List of excipients**

Benzyl alcohol (E1519)  
Sodium chloride  
Sodium hydroxide (for pH adjustment)  
Hydrochloric acid (for pH adjustment)  
Water for injections

## **6.2 Incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

## **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.  
Shelf life after first opening the immediate packaging: 28 days.

## **6.4 Special precautions for storage**

Store in the original package.

## **6.5 Nature and composition of immediate packaging**

Vials of clear colourless glass type I, filled with 5 ml, 10 ml, 20 ml, 25 ml, 30 ml and 50 ml.  
Bromobutyl rubber stoppers type I (the stoppers are secured with aluminium caps).  
1 vial in a cardboard box.

Not all pack sizes may be marketed.

## **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Eurovet Animal Health B.V.  
Handelsweg 25  
5531 AE Bladel  
The Netherlands

## **8. MARKETING AUTHORISATION NUMBER**

Vm 16849/4020

## **9. DATE OF FIRST AUTHORISATION**

20 December 2010

## **10. DATE OF REVISION OF THE TEXT**

October 2015

Approved: 21 October 2015

