

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Meloxaid 2 mg/ml Solution for Injection for Cats

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

One ml contains:

**Active substance:** Meloxicam 2 mg

**Excipients:**

| <b>Qualitative composition of excipients and other constituents</b> | <b>Quantitative composition if that information is essential for proper administration of the veterinary medicinal product</b> |
|---|--|
| Ethanol, anhydrous  | 150 mg   |
| Meglumine   |  |
| Macrogol 300  |  |
| Poloxamer 188   |  |
| Glycine   |  |
| Disodium edetate  |  |
| Sodium hydroxide  |  |
| Hydrochloric acid   |  |
| Water for Injections  |  |

Clear yellow solution.

### **3. CLINICAL INFORMATION**

#### **3.1 Target species**

Cats.

#### **3.2 Indications for use for each target species**

Alleviation of mild to moderate post-operative pain and inflammation following surgical procedures in cats, e.g. orthopaedic and soft tissue surgery.

#### **3.3 Contraindications**

Do not use in pregnant or lactating animals.

Do not use in cats suffering from gastrointestinal disorders such as irritation and haemorrhage, impaired hepatic, cardiac or renal function and haemorrhagic disorders.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in cats less than 6 weeks of age nor in cats of less than 2 kg.

### 3.4 Special warnings

None.

### 3.5 Special precautions for use

#### Special precautions for safe use in the target species:

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

Avoid use in any dehydrated, hypovolaemic or hypotensive cat, as there is a potential risk of renal toxicity.

During anaesthesia, monitoring and fluid therapy should be considered as standard practice.

In case additional pain relief is required, multimodal pain therapy should be considered.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Accidental self-injection may give rise to pain. People with known hypersensitivity to non-steroidal anti-inflammatory drugs (NSAIDs) should avoid contact with the veterinary medicinal product.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

This product can cause eye irritation. In case of contact with the eyes, immediately rinse thoroughly with water.

In view of the risk of accidental self-injection and the known adverse class-effects of NSAIDs and other prostaglandin inhibitors on pregnancy and/or embryofetal development, the veterinary medicinal product should not be administered by pregnant women or women attempting to conceive.

#### Special precautions for the protection of the environment:

Not applicable.

#### Other precautions:

Not applicable.

### 3.6 Adverse events

Cats

|   |  |
|---|--|
| Very rare (<1 animal / 10,000 animals treated, including isolated reports): | Appetite loss <sup>1</sup> , Lethargy <sup>1</sup><br>Renal failure <sup>1</sup><br>Vomiting <sup>1</sup> , Diarrhoea <sup>1</sup> , Blood in faeces <sup>1,2</sup> , Gastrointestinal ulceration <sup>1</sup><br>Elevated liver enzymes <sup>1</sup><br>Anaphylactoid reaction <sup>3</sup> |
|---|--|

<sup>1</sup>These adverse reactions are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal.

<sup>2</sup>Occult.

<sup>3</sup>Should be treated symptomatically.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

### **3.7 Use during pregnancy, lactation or lay**

The safety of the veterinary medicinal product has not been established during pregnancy and lactation (see section 3.3).

### **3.8 Interaction with other medicinal products and other forms of interaction**

Other NSAIDs, diuretics, anticoagulants, aminoglycoside antibiotics and substances with high protein binding may compete for binding and thus lead to toxic effects. The veterinary medicinal product must not be administered in conjunction with other NSAIDs or glucocorticosteroids. Concurrent administration of potential nephrotoxic veterinary medicinal products should be avoided. In animals at anaesthetic risk (e.g. aged animals) intravenous or subcutaneous fluid therapy during anaesthesia should be taken into consideration. When anaesthesia and NSAID are concomitantly administered, a risk for renal function cannot be excluded.

Pre-treatment with anti-inflammatory substances may result in additional or increased adverse effects and accordingly a treatment-free period with such veterinary medicinal products should be observed for at least 24 hours before commencement of treatment. The treatment-free period, however, should take into account the pharmacological properties of the products used previously.

### **3.9 Administration routes and dosage**

#### Reduction of post-operative pain and inflammation when administration of meloxicam is to be continued as an oral follow-up therapy:

Single subcutaneous injection at a dosage of 0.2 mg meloxicam/kg body weight (i.e. 0.1 ml/kg body weight) before surgery, for example at the time of induction of anaesthesia.

To continue treatment for up to five days, this initial dose may be followed 24 hours later by administration of an oral suspension veterinary medicinal product containing meloxicam authorised for cats at a dosage of 0.05 mg meloxicam/kg body weight. The oral follow-up dose may be administered for up to a total of four doses at 24-hour intervals.

#### Reduction of post-operative pain and inflammation where no oral follow-up treatment is possible e.g. feral cats:

Single subcutaneous injection at a dosage of 0.3 mg meloxicam/kg body weight (i.e. 0.15 ml/kg body weight) before surgery, for example at the time of induction of anaesthesia. In this case do not use oral follow up treatment.

Particular care should be taken with regard to the accuracy of dosing. Avoid introduction of contamination during use.

### **3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)**

In the case of overdose symptomatic treatment should be initiated.

### **3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance**

Not applicable.

### **3.12 Withdrawal periods**

Not applicable.

## **4. PHARMACOLOGICAL INFORMATION**

### **4.1 ATCvet code: QM01AC06**

### **4.2 Pharmacodynamics**

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) of the oxicam class which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, analgesic, anti-exudative and antipyretic effects. It reduces leukocyte infiltration into the inflamed tissue. To a minor extent it also inhibits collagen-induced thrombocyte aggregation. *In vitro* and *in vivo* studies demonstrated that meloxicam inhibits cyclooxygenase-2 (COX-2) to a greater extent than cyclooxygenase-1 (COX-1).

### **4.3 Pharmacokinetics**

#### Absorption

Following subcutaneous administration, meloxicam is completely bioavailable and maximal mean plasma concentrations of 1.1 µg/ml were reached approximately 1.5 hours post administration.

#### Distribution

There is a linear relationship between the dose administered and plasma concentration observed in the therapeutic dose range. More than 97 % of meloxicam is bound to plasma proteins. The volume of distribution is 0.09 l/kg.

#### Metabolism

Meloxicam is predominantly found in plasma and is also a major biliary excretion product whereas urine contains only traces of the parent compound. Five major metabolites were detected all having been shown to be pharmacologically inactive. Meloxicam is metabolised to an alcohol, an acid derivative and to several polar metabolites. As for other species investigated, the main pathway of meloxicam biotransformation in cat is oxidation.

### Elimination

Meloxicam is eliminated with a half-life of 24 hours. The detection of metabolites from the parent compound in urine and faeces, but not in plasma is indicative for their rapid excretion. 21 % of the recovered dose is eliminated in urine (2 % as unchanged meloxicam, 19 % as metabolites) and 79 % in the faeces (49 % as unchanged meloxicam, 30 % as metabolites).

## **5. PHARMACEUTICAL PARTICULARS**

### **5.1 Major incompatibilities**

None known.

### **5.2 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 1 year  
Shelf life after first opening the immediate packaging: 28 days

### **5.3 Special precautions for storage**

Keep the vial in the outer carton, in order to protect from light.  
This veterinary medicinal product does not require any special storage conditions.

### **5.4 Nature and composition of immediate packaging**

Cardboard box containing one colourless glass vial of 10 ml or 20 ml, closed with a bromobutyl rubber stopper and sealed with an aluminium cap.

Not all pack sizes may be marketed.

### **5.5 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Medicines should not be disposed of via wastewater.  
Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

## **6. NAME OF THE MARKETING AUTHORISATION HOLDER**

C&H Generics Ltd

## **7. MARKETING AUTHORISATION NUMBERS**

Vm 40162/3018 (NI)  
Vm 40162/5018 (GB)

**8. DATE OF FIRST AUTHORISATION**

06 November 2025

**9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS**

November 2025

**10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.  
Find more product information by searching for the 'Product Information Database'  
on [www.gov.uk](http://www.gov.uk).

*Gavin Hall*  
Approved: 17 December 2025