

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

PGF Veyx forte 0.250 mg/ml solution for injection for cattle and pigs

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

1 ml solution for injection contains:

**Active substance:**

Cloprostenol 0.250 mg (corresponding to 0.263 mg cloprostenol sodium)

**Excipients:**

Chlorocresol 1.0 mg

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Solution for injection.  
Clear, colourless aqueous solution.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Cattle (heifers, cows) and pigs (sows)

#### **4.2 Indications for use, specifying the target species**

Cattle (heifers, cows):

- To schedule the time of oestrous and ovulation and for cycle synchronization in animals with an ovulatory cycle when applied during the diestrus (induction of oestrus in non-detected oestrus, synchronisation of oestrus)
- Treatment of anoestrus and uterine disorders caused by a progesterone-induced oestrous cycle blockade (induction of oestrous in anoestrus, endometritis, pyometra, corpus luteal cysts, follicular luteal cysts, shortening of the sexual rest period)
- Induction of abortion up to day 150 of pregnancy
- Expulsion of mummified fetuses
- Induction of parturition

Pigs (sows):

- Induction or synchronisation of farrowing from day 114 of pregnancy onwards (day 1 of pregnancy is the last day of insemination).

### 4.3 Contraindications

- Do not use for intravenous administration
- Do not use in pregnant animals where the induction of abortion or parturition is not intended
- Do not use in case of spastic diseases of the respiratory tract and gastrointestinal tract
- Do not use in cases of hypersensitivity to the active substance or to any of excipients

### 4.4 Special warnings for each target species

None

### 4.5 Special precautions for use

#### Special precautions for use in

#### animals

To reduce the risk of anaerobic infections care should be taken to avoid injection through contaminated areas of skin. Clean and disinfect injection sites thoroughly before application.

#### **Pigs:**

Use only if the cover dates are known. Too early an administration could adversely affect the viability of the piglets. This is the case when the injection is given more than 2 days before the average gestation period of the stock. Day 1 of pregnancy is the last day of insemination. The gestation period is generally 111-119 days.

#### **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

- This product must be handled carefully to avoid accidental self-injection or contact with the skin or mucous membranes of the user.
- Prostaglandins of the F2 $\alpha$  type may be absorbed through the skin and may cause bronchospasm or miscarriage.
- Pregnant women, women in childbearing age, asthmatics and people with other respiratory tract diseases should wear waterproof gloves during administration of the product.
- Accidental spillage on the skin should be washed off immediately with soap and water.
- In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician. Should respiratory distress result from accidental inhalation or injection, a rapid acting bronchodilator, e.g. isoprenaline or salbutamol by inhalation is indicated.

#### **4.6 Adverse reactions (frequency and seriousness)**

Anaerobic infections may occur if anaerobic bacteria are introduced into the tissue by the injection, in particular following intramuscular injection.

Cattle:

When used for induction of parturition, the incidence of retained placenta may be increased depending on the time of treatment. In very rare cases, anaphylactic-type reactions can be observed which might be life threatening and require rapid medical care.

Pigs:

The abnormal behaviour that might occur in pigs immediately after treatment, when the drug has been used to induce parturition, is similar to that of sows before normal birth and normally subsides within one hour.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports)

#### **4.7 Use during pregnancy, lactation or lay**

Do not use in pregnant animals when abortion or induction of parturition is not intended. Safety of the product has not been established during lactation.

Use only accordingly to the benefit/risk assessment by the responsible veterinarian.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

Concurrent use of oxytocin and Cloprostenol increases the effects on the uterus. Do not use in animals being treated with non-steroidal anti-inflammatories, as the synthesis of endogenous prostaglandins is inhibited.

#### **4.9 Amounts to be administered and administration route**

For intramuscular injection in cattle (heifers, cows).

For deep intramuscular injection in pigs (sows) (with a needle at least 4 cm long).

Cattle (heifers, cows): 0.5 mg Cloprostenol/animal corresponding to 2.0 ml of the product/animal

In order to synchronise oestrus in a cattle herd, it is recommended that the product is administered on two occasions with an 11-day interval between treatments.

Pigs (sows): 0.175 mg Cloprostenol/animal corresponding to 0.7 ml of the product/animal Use automatic syringe equipment for the 50 ml vials.

Single administration.

The rubber stopper of the vial may be safely punctured up to 25 times. Otherwise, automatic syringe equipment, or a suitable draw-off needle, should be used for the 50 ml vials to avoid excessive puncturing of the closure.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In case of overdose the following symptoms may occur:

Increased heart rate, increased respiratory rate, bronchoconstriction, increased rectal temperature, increased defecation and urination, salivation, nausea and vomiting.

No antidotes are available.

#### **4.11 Withdrawal period(s)**

Cattle, pigs (meat and offal): 2

days Cattle (milk): zero hours

### **5. PHARMACOLOGICAL PROPERTIES**

#### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: prostaglandin-F<sub>2α</sub> - agonist  
ATCvet code: QG02AD90

Cloprostenol belongs to the group of prostaglandin-F<sub>2α</sub>-agonists, which exert a luteolytic effect dependent on species and time of treatment. Furthermore, this group of substances has a contractile effect on the smooth muscles (uterus, gastro-intestinal tract, respiratory tract, vascular system).

Treatment during diestrus or in the case of persistent corpus luteum causes luteolysis. The associated removal of the negative feed-back mechanism induced by progesterone leads in animals with cyclic ovarian function to a premature onset of heat and ovulation.

Cloprostenol produces a 200- to 400-fold luteolytic effect compared to prostaglandin-F<sub>2α</sub>; however, the effect on the smooth muscles appears to be equally strong.

#### **5.2 Pharmacokinetic particulars**

Peak plasma concentrations were observed within 15 minutes to 2 hours after intramuscular injection of Cloprostenol in cattle and pigs. The subsequent rapid elimination phase is characterised by a half-life from 1 to 3 hours, with a subsequent phase of slow elimination over a period of up to 48 hours, with a half-life of 28 hours.

Cloprostenol is distributed evenly throughout the tissues. Excretion via the faeces and urine is virtually equal. Less than 0.4% of the applied dose in cattle is eliminated via the milk. Maximum concentrations are measured approx. 4 hours post application.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Chlorocresol  
Citric acid monohydrate Sodium chloride  
Sodium hydroxide (for pH-  
adjustment) Sodium citrate  
Water for injections

### **6.2 Major incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 4 years  
Shelf life after first opening the immediate packaging: 28 days

### **6.4 Special precautions for storage**

Do not store above 25°C. Protect from light.  
Keep the vial in the outer carton.

### **6.5 Nature and composition of immediate packaging**

Vial of colourless glass, type I, with a fluorinated bromobutyl stopper and an aluminium cap; 1 vial (10 ml) in a cardboard box.  
1 vial (20 ml) in a cardboard box. 1 vial (50 ml) in a cardboard box.  
Not all pack sizes may be marketed.

### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

Veyx-Pharma  
GmbH  
Söhreweg 6  
34639  
Schwarzenborn  
Germany

**8. MARKETING AUTHORISATION NUMBER**

Vm 27569/4002

**9. DATE OF FIRST AUTHORISATION**

9 October 2012

**10. DATE OF REVISION OF THE TEXT**

December 2023

A handwritten signature in black ink, consisting of several vertical strokes followed by a long, sweeping horizontal stroke that curves upwards at the end.

Approved 07 December 2023