

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Euthanimal 20%, 200 mg/ml solution for injection

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each ml contains

#### **Active substance:**

Sodium pentobarbital      200 mg (equivalent to 182 mg pentobarbital)

#### **Excipients:**

Benzyl alcohol (E 1519)    20.0 mg  
Ethanol                        80.0 mg  
Ponceau 4R (E 124)        0.02 mg

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Solution for injection.  
Clear red solution.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Pigs, goats, sheep, cattle, horses, cats and dogs.

#### **4.2 Indications for use, specifying the target species**

For euthanasia.

#### **4.3 Contraindications**

Do not use for anaesthesia.

#### **4.4 Special warnings for each target species**

Intravenous injection of pentobarbital has the ability to cause induction excitement in several species of animal and adequate sedation should be applied if deemed necessary by the veterinary surgeon. Measures must be taken to avoid perivascular administration (e.g. by using intravenous catheter).

Check regularly, up to about 10 minutes post-administration, if live signs return (respiration, heartbeat, corneal reflex). In clinical trials it has been established that this might occur. If such live signs return, it is advised to repeat the administration using between 0.5 and 1 times the prescribed dose.

Avoid use in animals weighing more than 120 kg due to the large injection volume required and difficulty achieving rapid administration.

To reduce the risk of induction excitement, euthanasia should be performed in a quiet area.

In pigs, it was shown that there is a direct correlation between restraint and level of excitation and agitation. Therefore, injection in swine should be done with the least amount of restraint necessary.

Particularly in horses and cattle, the veterinary surgeon should consider premedication with an appropriate sedative to produce profound sedation before euthanasia and an alternative method of euthanasia is recommended to be available should it become necessary.

#### **4.5 Special precautions for use**

##### Special precautions for use in animals

Carcasses and edible products of animals injected with this product may never enter the food chain (see section 4.11) and should be disposed of in accordance with national legislation.

Carcasses or parts of the carcass of animals euthanised with this product should not be fed to other animals due to the risk of secondary intoxication (see section 4.11).

##### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Pentobarbital is a potent hypnotic and a sedative, and thus potentially toxic in man. It can be adsorbed systemically through the skin and if swallowed. Particular care should be taken to avoid accidental ingestion and self-injection.

Systemic uptake (including absorption via skin or eye) of pentobarbital causes sedation, sleep induction and respiratory depression. Moreover, this product may be irritating to the eye and can cause irritation to the skin as well as hypersensitivity reactions (due to the presence of pentobarbital and benzyl alcohol). Embryotoxic effects cannot be excluded.

Avoid direct contact with the skin and eyes, including hand-to-eye contact.  
Do not eat or drink while handling the product.

Avoid accidental self-injection or accidental injection of a second professional when administering the product. Only carry this veterinary medicinal product in an unarmed syringe to avoid accidental self-injection.

People with known hypersensitivity to pentobarbital should avoid contact with the veterinary medicinal product.

Handle the product with utmost care, especially women of childbearing potential. Wear protective gloves. This medicine should only be administered by veterinarians and should only be used in the presence of another professional that can assist in case of

accidental exposure. Instruct the professional if not a medical professional about the risks of the product.

Accidental spillage on the skin or in the eye must be washed off immediately with plenty of water. In case of accidental ingestion, wash out mouth immediately. If there has been serious skin or eye contact or in case of accidental ingestion or self-injection, seek medical attention immediately, indicate poisoning with barbiturates and show the package leaflet or the label to the physician. DO NOT DRIVE as sedation may occur.

After administration of this product, collapse will occur within 10 seconds. In case the animal is standing at time of administration, care should be taken by the person administering the veterinary medicinal product and any other persons present to keep a certain distance from the animal to avoid injury.

This product is flammable, keep away from sources of ignition. Do not smoke.

**Information for the health professional in case of exposure:**

Emergency measures should be directed toward maintenance of respiration and cardiac function. In severe intoxication measures to enhance elimination of absorbed barbiturate may be necessary. Do not leave the patient unattended.

The concentration of pentobarbital in the product is such that the accidental injection or ingestion of quantities as small as 1 ml in human adults can have serious CNS effects. A dose of pentobarbital sodium of 1 g (equivalent to 5 ml of product) has been reported to be fatal in humans. Treatment should be supportive with appropriate intensive therapy and maintenance of respiration.

**4.6 Adverse reactions (frequency and seriousness)**

Death may be delayed if the injection is administered perivascularly or into organs/tissues with low capacity for absorption. Barbiturates can be irritating when administered perivascularly.

Common (more than 1 but less than 10 animals in 100 animals treated):

- Vocalisation
- Muscle twitching

Rare (more than 1 but less than 10 animals in 10,000 animals treated):

- Excitation
- Leg movements
- Defecation and urine loss
- Gasping (in cattle), mostly due to underdosing

Very rare (less than 1 animal in 10,000 animals treated, including isolated reports):

- Convulsions
- Contraction of the diaphragm
- Vomiting
- One or few gasping respirations occur after cardiac arrest

#### **4.7 Use during pregnancy, lactation or lay**

No specific information is available.

The use of this product in pregnant animals is left to the judgement of the veterinarian.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

CNS depressants (narcotics, phenothiazines, antihistamines, etc.) may increase the effect of pentobarbital.

#### **4.9 Amounts to be administered and administration route**

Cattle, horses, pigs, goats, sheep, cats and dogs: 100 mg/kg (corresponding to 0.5 ml/kg) through quick intravenous injection. For larger animals, the use of a pre-inserted intravenous catheter is advocated.

If cardiac arrest is not established after 2 minutes, a second dose needs to be administered, preferably via fast intravenous injection or if this is not feasible, via intra-cardiac injection; intra-cardiac injection is only acceptable after prior deep sedation or anaesthesia.

As the vial cannot be punctured more than 20 times, the user should choose the most appropriate vial size.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In the event of accidental administration to an animal not presented for euthanasia, measures such as artificial respiration, administration of oxygen and the use of analeptics are appropriate.

In view of the activity of this product, double dosage is discouraged, as this will not result in a faster or better euthanasia.

#### **4.11 Withdrawal period(s)**

Not applicable.

Appropriate measures need to be taken to assure that carcasses and edible products of animals injected with this product do not enter the food chain, and are not used for human consumption. Other animals may never eat (parts of) the carcass, as they might be exposed to a lethal dose of pentobarbital.

### **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic group: Barbiturates intended for euthanasia.  
ATCvet code: QN51AA01

## **5.1 Pharmacodynamic properties**

Pentobarbital is a short-acting sedative and hypnotic. It causes depression of the central nervous system by GABA receptor modulation, imitating the action of Gamma-aminobutyric acid.

Barbiturates suppress in particular the reticular activating system (RAS) in the brain, which normally ensures alertness. The immediate effect is the loss of consciousness followed by deep anaesthesia followed by, at high rates, rapid depression of the respiratory centre. Breathing stops and is quickly followed by cardiac arrest and rapid death.

## **5.2 Pharmacokinetic particulars**

After intravenous administration fast distribution over the tissues will occur. Pentobarbital is mainly eliminated through the liver by biotransformation, particularly by the Cytochrome P<sub>450</sub> system, as well as by excretion in the kidneys and redistribution. In pigs redistribution in fatty tissue might cause reduced plasma concentrations and prolonged action.

Barbiturates may diffuse through the placenta in foetal tissue, and traces of barbiturates may be present in the breast milk.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Ethanol 96%  
Propylene glycol  
Benzyl alcohol (E 1519)  
Ponceau 4R (E 124)  
Water for injections

### **6.2 Major incompatibilities**

In the absence of incompatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.  
Shelf life after first opening the immediate packaging: 28 days.

### **6.4 Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

### **6.5 Nature and composition of immediate packaging**

1 vial of 100 ml or 1 vial of 250 ml, type II glass injection vial with a bromobutylrubber stopper and aluminium cap in carton box.

12 vials of 100 ml or 6 vials of 250 ml, type II glass injection vial with a bromobutylrubber stopper and aluminium cap in polystyrene box.

Not all pack sizes may be marketed.

#### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

This medicinal product is dangerous to humans and animals. Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

#### **7. MARKETING AUTHORISATION HOLDER**

Alfasan Nederland BV  
Kuipersweg 9  
3449 JA Woerden  
The Netherlands

#### **8. MARKETING AUTHORISATION NUMBER**

Vm 36408/4001

#### **9. DATE OF FIRST AUTHORISATION**

02 October 2013

#### **10. DATE OF REVISION OF THE TEXT**

December 2017

#### **PROHIBITION OF SALE, SUPPLY AND/OR USE**

Medicine for the exclusive use of the veterinary surgeon. Not for sale to the public. The administration and custody of the medicine should only be carried out by the veterinary surgeon.

Approved: 21 December 2017

