

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Amoxy Active, 697 mg/g, oral powder for pigs and chickens

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each gram contains:

#### **Active substance:**

Amoxicillin	697 mg
as amoxicilline trihydrate	800 mg

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Oral powder.

White to off-white oral powder.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Pigs and chickens (broilers, pullets, chickens for reproduction).

#### **4.2 Indications for use, specifying the target species**

Pigs: Treatment of respiratory tract infections, gastro-intestinal tract infections, urogenital infections, secondary infections following viral infections and septicaemia caused by micro-organisms susceptible to amoxicillin.

Chickens: Treatment of respiratory tract infections and gastro-intestinal tract infections caused by micro-organisms susceptible to amoxicillin.

#### **4.3 Contraindications**

Do not use in cases of hypersensitivity to penicillin or other substances of the beta-lactam group or to any of the excipients.

Do not use in the presence of  $\beta$ -lactamase-producing bacteria.

Do not use in lagomorphs and rodents such as guinea pigs, hamsters or gerbils.

Do not use in animals with serious kidney malfunction including anuria and oliguria.

Do not use in ruminants or horses.

#### 4.4 Special warnings for each target species

Sick animals have an altered drinking behaviour and should be medicated parenterally where applicable.

#### 4.5 Special precautions for use

##### Special precautions for use in animals

Official, national and regional antimicrobial policies should be taken into account when the veterinary medicinal product is used. Use of the veterinary medicinal product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

Use of the veterinary medicinal product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to the amoxicillin and may decrease the effectiveness of the treatment.

##### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Penicillins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may cause cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious. People with known hypersensitivity to beta-lactam antibiotics should avoid contact with the veterinary medicinal product. Handle this veterinary medicinal product with great care to avoid exposure, taking all recommended precautions. Personal protective equipment consisting of gloves and either a disposable half mask respirator conforming to European Standard EN149 or a non-disposable respirator conforming to European Standard EN140 with a filter to EN 143 should be worn when mixing and handling the veterinary medicinal product. Wash hands after use. In case of contact with eyes or skin, wash immediately with water. If you develop symptoms following exposure, such as a skin rash, you should seek medical advice and show the package leaflet or the label to the physician. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

##### Special precautions for the protection of the environment

Not applicable.

#### 4.6 Adverse reactions (frequency and seriousness)

Pigs and chickens:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Hypersensitivity reactions*  Digestive tract disorders (vomiting, diarrhoea)
---	--

\*the severity varying from skin rash to anaphylactic shock.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or the national competent authority via the national reporting system. See section 'Contact details' of the package leaflet.

#### **4.7 Use during pregnancy, lactation or lay**

##### Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy and lactation. Laboratory studies in rats and rabbits have not produced any evidence of a teratogenic, foetotoxic or maternotoxic effects. Use only accordingly to the benefit/risk assessment by the responsible veterinarian.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

Do not combine with bacteriostatic antibiotics.

Not to be used simultaneously with neomycin since it blocks the absorption of oral penicillins. Synergism occurs with  $\beta$ -lactam antibiotics and aminoglycosides.

#### **4.9 Amount(s) to be administered and administration route**

In drinking water use and in-feed use in pigs.

In drinking water use in chickens.

##### Pigs:

The recommended dose is 11.2 mg amoxicillin per kg of body weight daily (corresponding to 16.1 mg of the veterinary medicinal product per 1 kg of body weight per day) given for 3 - 5 consecutive days.

##### Chickens:

The recommended dose is 20 mg amoxicillin per kg of body weight daily (corresponding to 28.7 mg of the veterinary medicinal product per 1 kg of body weight per day) given for 3 - 5 consecutive days.

To ensure a correct dosage, body weight should be determined as accurately as possible to avoid underdosing. The use of suitably calibrated measuring equipment is recommended.

##### *In drinking water use:*

For the preparation of medicated water, the body weight of the animals to be treated and their actual daily water consumption should be taken into account. Consumption may vary depending on factors like species, age, state of health, breed and husbandry system (e.g. different temperature, different light regimes). The intake of medicated water depends on the clinical condition of the animals. In order to obtain the correct dosage, the concentration of amoxicillin may need to be adjusted accordingly.

Preparation of medicated water should provide an amount to be consumed within the next 12 hours. Any unused medicated water should be discarded after 12 hours, and freshly medicated water for the next 12 hours should be prepared.

Based on the recommended dose and the number and weight of animals to be treated, the exact daily concentration of the veterinary medicinal product should be calculated according to the following formula:

$$\frac{\dots \text{mg veterinary medicinal product/}}{\text{kg body weight/day}} \times \frac{\text{average body weight (kg)}}{\text{of animals to be treated}} = \frac{\dots \text{ mg veterinary medicinal product per litre of drinking water}}{\text{average daily water intake (l/animal)}}$$

The veterinary medicinal product should be added to the drinking water by thorough stirring until the veterinary medicinal product is completely dissolved. Maximum solubility of the veterinary medicinal product in water is approximately 6 g/litre. Sufficient access to the system of water supply should be available for the animals to be treated to ensure adequate water consumption. No other source of drinking water should be available during the medication period. In free range husbandry systems animals should be kept in the stable during treatment.

Where applicable the water supply system should be cleaned appropriately after the end of the medication period to avoid intake of sub-therapeutic amounts of the active substance.

#### *In-feed use:*

The veterinary medicinal product may also be offered via the feed at the recommended daily dose. This way of administration is only intended for the treatment of individual pigs on farms where only a small number of pigs are to receive the treatment. Only the pack size of 100 g is suitable for the in-feed use.

Larger groups should be treated with medicated drinking water.

Before each administration the powder should be thoroughly mixed into a small amount of food and should be given directly to the animal before the main ration. Care should be taken that the intended dose will be completely ingested.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In case of overdosing no other effects are known than mentioned in section 4.6 Adverse reactions.

#### **4.11 Withdrawal period(s)**

Pigs: meat and offal: 2 days.

Chickens: meat and offal: 1 day.

Not for use in birds producing eggs for human consumption.

Do not use within 4 weeks of the start of the laying period.

### **5. PHARMACOLOGICAL PROPERTIES**

**Pharmacotherapeutic group:** Beta-lactam antibacterials, Penicillins

**ATCvet code:** QJ01CA04

#### **5.1 Pharmacodynamic properties**

Amoxicillin, is a broad-spectrum penicillin with bactericidal action against many Gram-positive and Gram negative bacteria.

It owes its activity to the inhibition of the development of the peptidoglycan network structure in the bacterial cell wall.

Amoxicillin is acid resistant, but is not resistant to the action of beta-lactamases.

## **5.2 Pharmacokinetic particulars**

Amoxicillin is rapidly and almost completely absorbed from the gastrointestinal tract and is stable in the presence of gastric acids. Maximum amoxicillin concentrations are reached within 1-2 hours. Serum protein binding is low. Amoxicillin is widely distributed throughout the body.

Amoxicillin is mainly eliminated via the kidneys in the active form to give high concentrations in renal tissue and urine. A smaller part of the administered dose of amoxicillin is excreted in the bile.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Sodium carbonate  
Sodium citrate

### **6.2 Major incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.

Shelf life after first opening of the immediate packaging: 28 days.

Shelf life after dissolution according to directions: 12 hours.

Shelf life after incorporation into meal: use immediately.

### **6.4 Special precautions for storage**

Store below 25 °C.

Store in the original container.

### **6.5 Nature and composition of immediate packaging**

- Securitainer: white polypropylene container, covered with a low-density polyethylene lid. The securitainer contains 100 g, 250 g, 500 g or 1 kg of veterinary medicinal product.
- Bucket: white polypropylene bucket provided with a polypropylene lid. The bucket contains 1 kg, 2.5 kg or 5 kg of veterinary medicinal product.

Not all pack sizes may be marketed.

## **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Medicines should not be disposed of via wastewater.  
Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

## **7 MARKETING AUTHORISATION HOLDER**

Dopharma Research B.V.  
Zalmweg 24  
4941 VX Raamsdonksveer  
The Netherlands

## **8. MARKETING AUTHORISATION NUMBER**

Vm 28365/5004

## **9. DATE OF FIRST AUTHORISATION**

27 June 2014

## **10. DATE OF REVISION OF THE TEXT**

October 2023

## **PROHIBITION OF SALE, SUPPLY AND/OR USE**

## **11. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCT**

Veterinary medicinal product subject to prescription.

Approved 10 January 2024

