

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Milaxyn 230/20mg Flavoured Film Coated Tablets for Cats (UK)

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each film coated tablet contains:

#### **Active substances:**

Pyrantel embonate            230 mg

Praziquantel                 20 mg

#### **Excipients:**

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Film-coated tablets.

A white to off white round, biconvex coated tablet with a breakline on one side and plain on the other side.

The tablets can be divided into 2 equal parts

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Cats.

#### **4.2 Indications for use, specifying the target species**

For the treatment of mixed infections caused by the following gastrointestinal roundworms and tapeworms:

**Roundworms:** *Toxocara cati*, *Toxascaris leonina*,

**Tapeworms:** *Dipylidium caninum*, *Taenia taeniaeformis*, *Echinococcus multilocularis*.

#### **4.3 Contraindications**

Do not use simultaneously with piperazine compounds.

Do not use in kittens less than 6 weeks of age.

Do not use in cases of hypersensitivity to the active substances or to any of the

excipients.

#### **4.4 Special warnings for each target species**

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*. Tapeworm infestation is certain to reoccur unless control of intermediate hosts such as fleas, mice, etc. is undertaken.

Parasitic resistance to a certain class of anthelmintics can occur after frequent and repeated use of an anthelmintic from this class.

#### **4.5 Special precautions for use**

##### **Special precautions for use in animals**

Not applicable.

##### **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

In the interests of good hygiene, persons administering the tablets directly to the cat, or by adding them to the cat's food, should wash their hands afterwards.

In case of accidental ingestion, seek medical advice and show the package leaflet to the physician.

##### **Other precautions**

Echinococcosis represents a hazard for humans. As Echinococcosis is a notifiable disease to the World Organisation for Animal Health (OIE), specific guidelines on the treatment and follow-up, and on the safeguard of persons, need to be obtained from the relevant competent authority.

#### **4.6 Adverse reactions (frequency and seriousness)**

Mild and transient digestive tract disorders such as hypersalivation and/or vomiting and mild and transient neurological disorders such as ataxia may occur very rarely.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)

- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

#### **4.7 Use during pregnancy, lactation or lay**

Do not use during pregnancy but may be used during lactation.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

Do not use simultaneously with piperazine compounds.

#### **4.9 Amounts to be administered and administration route**

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

##### *Dosage*

The recommended dose is: 20 mg/kg pyrantel (57.5 mg/kg pyrantel embonate) and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 4 kg bodyweight.

Body weight	tablets
1.0 - 2.0 kg	½
2.1 - 4.0 kg	1
4.1 - 6.0 kg	1 ½
6.1 - 8.0 kg	2

##### *Administration and duration of treatment*

Single oral administration. The tablet should be given directly to the cat, but if necessary can be disguised in food.

In ascarid infestation, especially in kittens, complete elimination cannot be expected, so a risk of infection for humans can persist. Repeat treatments should, therefore, be carried out with a suitable roundworm product at 14 day intervals until 2-3 weeks after weaning.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

After doses higher than 5 times the recommended dose, signs of intolerance such as vomiting have been observed.

#### **4.11 Withdrawal period(s)**

Not applicable.

## 5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Anthelmintics, praziquantel combinations.

ATC vet code: QP52AA51

### 5.1 Pharmacodynamic properties

This product contains anthelmintics active against gastrointestinal roundworms and tapeworms. The product contains two active substances, as follows:

1. Pyrantel embonate (pamoate), a tetrahydropyrimidine derivative and
2. Praziquantel, a partially hydrogenated pyrazinoisoquinoline derivative.

Pyrantel acts as a cholinergic agonist. Its mode of action is to stimulate nicotinic cholinergic receptors of the parasite, induce spastic paralysis and thereby allow expulsion from the gastro-intestinal (GI) system by peristalsis.

Praziquantel is very rapidly absorbed and distributed throughout the parasite. Both in vitro and in vivo studies have shown that praziquantel causes severe damage to the parasite integument, resulting in contraction and paralysis. There is an almost instantaneous tetanic contraction of the parasite musculature and a rapid vacuolization of the syncytial tegument. This rapid contraction has been explained by changes in divalent cation fluxes, especially calcium.

In this fixed combination, pyrantel is active against the following ascarids: *Toxocara cati*, and *Toxascaris leonina*. Praziquantel is effective against tapeworms in particular *Dipylidium caninum* and *Taenia taeniaeformis*.

Since it contains praziquantel, the product is effective against *Echinococcus multilocularis*.

### 5.2 Pharmacokinetic particulars

Praziquantel is rapidly absorbed, metabolised and distributed in the body. It is also believed to be excreted back into the intestinal lumen by the mucous membrane.

Following administration of the product to cats, peak plasma concentrations of praziquantel were achieved by approximately 2 hours.

Pyrantel is poorly absorbed so it is expected that a large proportion of the administered dose remains in the GIT where it exerts its therapeutic effect and it is excreted largely unchanged in the faeces.

Following administration of the product to cats, peak plasma concentrations of pyrantel were achieved by approximately 3 hours.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Core tablet:

Maize starch

Microcrystalline cellulose (E460)

Crospovidone

Magnesium stearate (E572)

Colloidal anhydrous silica,

#### Film coat

Grilled meat flavour

Opadry Complete Film Coating System 03F28415 White consisting of :

HPMC 2910 /Hypromellose (E464),

Macrogol/PEG 4000 (E1521),

Titanium Dioxide (E171).

### **6.2 Major Incompatibilities**

Not Applicable

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 5 years

Discard unused half tablets.

#### **6.4 Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions

#### **6.5 Nature and composition of immediate packaging**

The product is presented in either:

Individual blisters made up of a PVC/PE/PCTFE white opaque copolymer and a 20µm heatseal lacquer/aluminium containing 2, 4, 6, 8, 10, 12, 14, 16, 18 or 20 tablets.

or

Individual blisters made up of 45µm PVC/aluminium/orientated polyamide and a 20µm heatseal lacquer/aluminium containing 2 or 8 tablets.

The blisters are packed into cartons containing either: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 24, 28, 30, 32, 36, 40, 42, 44, 48, 50, 52, 56, 60, 64, 68, 70, 72, 76, 80, 84, 88, 92, 96, 98, 100, 104, 106, 108, 112, 116, 120, 128, 136, 140, 144, 150, 152, 160, 168, 176, 180, 184, 192, 200, 204, 206, 208, 216, 224, 232, 240, 248, 250, 280, 300, 500 or 1000 tablets

Not all pack sizes may be marketed.

#### **6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements

### **7. MARKETING AUTHORISATION HOLDER**

C&H Generics Ltd  
c/o Michael McEvoy and Co  
Seville House  
New Dock Street  
Galway  
Ireland

### **8. MARKETING AUTHORISATION NUMBER**

Vm 40162/4009

**9. DATE OF FIRST AUTHORISATION**


19 March 2015

**10. DATE OF REVISION OF THE TEXT**

March 2020

**PROHIBITION OF SALE, SUPPLY AND/OR USE**

Veterinary medicinal product subject to prescription.  
To be administered by a veterinary surgeon or under their direct responsibility.

A handwritten signature in black ink, consisting of several loops and a final horizontal stroke.

Approved: 11 March 2020