

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Cefaseptin 300 mg Tablets for Dogs

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

One tablet contains:

Active substance:

cefalexin (as cefalexin monohydrate) .....300 mg

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Beige oblong tablet.

The tablet can be divided into 2 or 4 equal parts.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Dogs

#### **4.2 Indications for use, specifying the target species**

For the treatment of bacterial skin infections (including deep and superficial pyoderma) caused by organisms, including *Staphylococcus* spp., susceptible to cefalexin.

For the treatment of urinary-tract infections (including nephritis and cystitis) caused by organisms, including *Escherichia coli*, susceptible to cefalexin.

#### **4.3 Contraindications**

Do not use in cases of known hypersensitivity to the active substance, to other cephalosporins, to other substances of the  $\beta$ -lactam group or to any of the excipients. Do not use in known cases of resistance to cephalosporins or penicillins.

Do not use in rabbits, guinea pigs, hamsters and gerbils.

#### **4.4 Special warnings for each target species**

None

## 4.5 Special precautions for use

### Special precautions for use in animals

The need for systemic antibiotics compared with non-antibiotic alternatives for the treatment of superficial pyoderma should be carefully considered by the responsible veterinarian.

As with other antibiotics which are excreted mainly by the kidneys, systemic accumulation may occur in the body when renal function is impaired. In case of known renal insufficiency, the dose should be reduced and antimicrobials known to be nephrotoxic should not be administered concurrently.

This product should not be used to treat puppies of less than 1 kg of bodyweight.

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about the susceptibility of the target bacteria.

Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to cefalexin and may decrease the effectiveness of treatment with other cephalosporins and penicillins, due to the potential for cross-resistance.

Official, national and regional antimicrobial policies should be taken into account when the product is used.

*Pseudomonas aeruginosa* is known for intrinsic (or natural) resistance to cefalexin.

The tablets are flavoured (presence of porcine liver powder). In order to avoid accidental ingestion, store tablets out of reach of the animals.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross-reactions to cephalosporin and vice versa. Allergic reactions to these substances may occasionally be serious.

1. Do not handle this product if you know you are sensitised or if you have been advised not to work with such preparations.
2. Handle this product with great care to avoid exposure, taking all recommended precautions. Wash hands after use.
3. If you develop symptoms following exposure such as skin rash, you should seek medical advice and show the package leaflet or the label to the physician. Swelling of the face, lips or eyes or difficulty in breathing are more-serious symptoms and require urgent medical attention.

## 4.6 Adverse reactions (frequency and seriousness)

In rare cases hypersensitivity can occur.

In cases of hypersensitivity reactions the treatment should be stopped.

In very rare cases, nausea, vomiting and/or diarrhoea have been observed in some dogs after administration.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals displaying adverse reaction(s) during the course of one treatment)
- common (more than 1 but less than 10 animals in 100 animals)
- uncommon (more than 1 but less than 10 animals in 1,000 animals )
- rare (more than 1 but less than 10 animals in 10,000 animals)
- very rare (less than 1 animal in 10,000 animals, including isolated reports).

#### **4.7 Use during pregnancy, lactation or lay**

The safety of the veterinary medicinal product has not been established in bitches during pregnancy and lactation.

Laboratory studies have not produced any evidence of teratogenic effects in mice (up to 400 mg cefalexin/kg bw/day) and rats (up to 1200 mg cefalexin/kg bw/day). In mice, maternal effects and foetotoxicity were observed from the lowest dose tested (100 mg cefalexin/kg bw/day). In rats, there is evidence of foetotoxicity at 500 mg cefalexin/kg bw/day and maternal effects from the lowest dose tested (300 mg cefalexin/kg bw/day).

Use only in accordance with the benefit/risk assessment by the responsible veterinarian.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

In order to ensure efficacy, the veterinary medicinal product should not be used in combination with bacteriostatic antibiotics (macrolides, sulfonamides and tetracyclines). Concurrent use of first generation cephalosporins with aminoglycoside antibiotics or some diuretics such as furosemide can enhance nephrotoxicity risks. Concomitant use with such active substances should be avoided.

#### **4.9 Amounts to be administered and administration route**

Oral use.

15 mg of cefalexin per kg of bodyweight twice daily (equivalent to 30 mg per kg of bodyweight per day) corresponding to one tablet per 20 kg bodyweight twice daily.

Dogs:

Urinary-tract infection: 14 days

Superficial bacterial infection of the skin: At least 15 days

Deep bacterial infection of the skin: At least 28 days

To ensure correct dosage, bodyweight should be determined as accurately as possible to avoid underdosing.

The product may be crushed or added to food if necessary.

In severe or acute conditions, except in cases of known renal insufficiency (see section 4.5), the dose may be doubled.

#### 4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Studies on animals with up to 5 times the recommended twice daily dosage of 15 mg cefalexin/kg have been performed.

Adverse reactions that may occur at the recommended dose (nausea, vomiting, and/or diarrhea) are expected in the case of overdose. In the event of overdose, treatment should be symptomatic.

#### 4.11 Withdrawal period(s)

Not applicable.

### 5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Antibacterials for systemic use, first generation cephalosporins  
ATCvet code: QJ01DB01

#### 5.1 Pharmacodynamic properties

Cefalexin is a time-dependent bactericidal antibiotic that acts by inhibiting the nucleopeptide synthesis of the bacterial wall. Cephalosporins interfere with the enzymes of transpeptidation making it unable to cross-link the peptidoglycans of the bacterial cell wall. The glycan cross-linking is essential for the cell to build its cell wall. Inhibition of the biosynthesis results to a weakened cell wall, which eventually ruptures to osmotic pressure. The combined action results in cell lysis and filament formation.

Cefalexin is active against a wide range of Gram-positive (e.g. *Staphylococcus* spp.) and Gram-negative (e.g. *Escherichia coli*) aerobic bacteria.

The following breakpoints are recommended by the CLSI (VET08, 4<sup>th</sup> edition, August 2019) in dogs:

In dogs for skin and soft tissue infections:

Bacterial species	Susceptible	Resistant
<i>Staphylococcus aureus</i>	≤ 2	≥ 4
<i>Staphylococcus pseudintermedius</i>		
<i>Streptococcus</i> spp and <i>E. coli</i>	≤ 2	≥ 8

In dogs for urinary tract infections:

Bacterial species	Susceptible	Resistant
<i>E. coli</i>	≤ 16	≥ 32
<i>Klebsiella pneumoniae</i>		
<i>Proteus mirabilis</i>		

MIC data for the use of Cefaseptin in dogs with skin and soft tissues infections and with urinary tract infections.

Data were collected between 2011 and 2017.

<b>Bacterial species</b>	<b>Range of MIC (mg/L)</b>	<b>MIC<sub>50</sub> (mg/L)</b>	<b>MIC<sub>90</sub> (mg/L)</b>
<b>Dermatological infections</b>			
<b><i>Staphylococcus</i> spp.<sup>a</sup></b>	0.25-512	0.993	12.435
<b><i>Staphylococcus aureus</i><sup>b</sup></b>	1-512	2.160	153.987
<b><i>Coagulase negative staphylococcus</i><sup>c</sup></b>	0.25-64	0.989	14.123
<b><i>Staphylococcus pseudintermedius</i><sup>b</sup></b>	0.5-512	0.768	5.959
<b><i>Streptococcus</i> spp.<sup>d</sup></b>	0.06-0.5	0.155	0.234
<b><i>Streptococcus canis</i><sup>d</sup></b>	0.06-0.5	0.146	0.226
<b><i>Streptococcus dysgalactiae</i><sup>d</sup></b>	0.25-0.5	0.185	0.354
<b><i>Escherichia coli</i><sup>b</sup></b>	4-512	5.481	11.314
<b><i>Pasteurella multocida</i><sup>b</sup></b>	0.12-4	1.373	1.877
<b>Urinary tract infections</b>			
<b><i>Proteus mirabilis</i><sup>b</sup></b>	8-512	6.498 - 12.491	12.553 - 207.937
<b><i>Klebsiella pneumoniae</i><sup>b</sup></b>	2-512	3.564	362.039
<b><i>E. coli</i><sup>b</sup></b>	4-512	5.022-5.82	7.671-13.929

a: period 2011-2017; b: period 2011-2015; c: period 2016-2017; d: period 2012-2015.

Resistance to cefalexin can be due to one of the following mechanisms of resistance. Firstly, the production of cephalosporinases, that inactivate the antibiotic by hydrolysis of the  $\beta$ -lactam ring, is the most prevalent mechanism among Gram-negative bacteria. This resistance is transmitted by plasmid or chromosomally. Secondly, a decreased affinity of the PBPs (penicillin-binding proteins) for beta-lactam drugs is frequently involved for beta-lactam resistant Gram-positive bacteria. Lastly, efflux pumps, extruding the antibiotic from the bacterial cell, and structural changes in porins, reducing passive diffusion of the drug through the cell wall, may contribute to improve the resistant phenotype of a bacterium.

Well-known cross-resistance (involving the same resistance mechanism) exists between antibiotics belonging to the beta-lactam group due to structural similarities. It occurs with beta-lactamases enzymes, structural changes in porins or variations in efflux pumps. Co-resistance (different resistance mechanisms involved) has been described in *E. coli* due to a plasmid harbouring various resistance genes.

*Pseudomonas aeruginosa* is known for resistance to cefalexin.

## **5.2 Pharmacokinetic particulars**

After single oral administration of the recommended dosage of 15 mg of cefalexin per kg of bodyweight to Beagle dogs, plasma concentrations were observed within 30 minutes. The plasma peak was observed at 1.3 hour with a plasma concentration of 18.2 µg/ml.

The bioavailability of the active was over 90 %. Cefalexin was detected until 24 hours after the administration. The first urine specimen was collected within 2 to 12 hours with peak concentrations of cefalexin measured at 430 to 2758 µg/ml within 12 hours.

After repeated oral administration of the same dosage, twice a day for 7 days, plasma peaks occurred 2 hours later with a concentration of 20 µg/ml. Over the treatment period, concentrations were maintained above 1 µg/ml. The mean elimination half-life is 2 hours. Skin levels were around 5.8 to 6.6 µg/g, 2 hours after treatment.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose monohydrate  
Povidone K30  
Croscarmellose sodium  
Microcrystalline cellulose  
Porcine liver powder  
Yeast  
Crospovidone  
Sodium stearyl fumarate

### **6.2 Major incompatibilities**

Not applicable

### **6.3 Shelf life**

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.  
Shelf life after first opening the immediate packaging: 48 hours.

### **6.4 Special precautions for storage**

Store in the original package.  
Return any part used tablet to the opened blister-pack.

### **6.5 Nature and composition of immediate packaging**

PVC/aluminium/OPA – PVC blister  
Cardboard box of 1 blister of 10 tablets  
Cardboard box of 10 blisters of 10 tablets  
Cardboard box of 25 blisters of 10 tablets

Not all pack sizes may be marketed.

**6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

Vetoquinol UK Limited  
Steadings Barn  
Pury Hill Business Park  
Nr Alderton  
Towcester  
Northamptonshire  
NN12 7LS

**8. MARKETING AUTHORISATION NUMBER**


Vm 08007/4142

**9. DATE OF FIRST AUTHORISATION**

12 January 2016

**10. DATE OF REVISION OF THE TEXT**

November 2020

A handwritten signature in black ink, consisting of several loops and a long horizontal stroke at the end.

Approved: 24 November 2020