

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

ITCH WORMER for Cats & Kittens 230/20mg Flavoured Tablets

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each film-coated tablet contains Pyrantel embonate 230 mg and Praziquantel 20 mg

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Film-coated tablets.

A white to off white round, biconvex coated tablet with a breakline on one side and plain on the other side.

The tablet can be divided in half.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Cats.

#### **4.2 Indications for use, specifying the target species**

For the treatment of mixed infections caused by the following gastrointestinal roundworms and tapeworms:

**Roundworms:** *Toxocara cati*, *Toxascaris leonina*,

**Tapeworms:** *Dipylidium caninum*, *Taenia taeniaeformis*, *Echinococcus multilocularis*.

#### **4.3 Contraindications**

Do not use simultaneously with piperazine compounds as piperazine may block the action of pyrantel embonate contained in the product.

Other worming products may contain piperazine.

Do not use simultaneously with other deworming products without veterinary advice.

Do not use in kittens less than 6 weeks of age.

Do not use in animals with known hypersensitivity to the active substances or to any of the excipients.

Do not use during pregnancy.

#### **4.4 Special warnings for each target species**

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*. Tapeworm infestation is certain to reoccur unless control of intermediate hosts such as fleas, mice, etc. is undertaken.

Parasitic resistance to a certain class of anthelmintics can occur after frequent and repeated use of an anthelmintic from this class.

#### **4.5 Special precautions for use**

##### **Special precautions for use in animals**

Not intended for use in cats weighing less than 1 kg bodyweight.

##### **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

In the interests of good hygiene, persons administering the tablets directly to the cat, or by adding them to the cat's food, should wash their hands afterwards.

In case of accidental ingestion, seek medical advice and show the package leaflet to the physician.

Other precautions

Echinococcosis represents a hazard for humans. As echinococcosis is a notifiable disease to the World Organisation for Animal Health (OIE), specific guidelines on the treatment and follow-up, and on the safeguard of persons, need to be obtained from the relevant competent authority.

#### **4.6 Adverse reactions (frequency and seriousness)**

Mild and short-lived digestive tract disorders such as excessive salivation and/or vomiting and mild and short-lived disorders of the nervous system such as loss of balance may be reported very rarely in spontaneous reports.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

#### **4.7 Use during pregnancy, lactation or lay**

Do not use during pregnancy but may be used during lactation.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

Do not use simultaneously with piperazine compounds (see section 4.3 above).

#### 4.9 Amounts to be administered and administration route

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

##### *Dosage*

The recommended dose is: 20 mg/kg pyrantel (57.5 mg/kg pyrantel embonate) and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 4 kg bodyweight.

Body weight	tablets
1.0 - 2.0 kg	½
Greater than 2.0 up to 4.0 kg	1
Greater than 4.0 up to 6.0 kg	1 ½
Greater than 6.0 kg	2

##### *Administration and duration of treatment*

Single oral administration.

The tablet should be given directly to the cat, but if necessary can be disguised in food.

In roundworm infestation, especially in kittens, complete elimination cannot be expected, so a risk of infection for humans can persist. Repeat treatments should, therefore, be carried out with a suitable roundworm product at 14 day intervals until 2-3 weeks after weaning.

If signs of disease persist or appear, consult a veterinary surgeon.

#### 4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

After doses higher than 5 times the recommended dose, signs of intolerance such as vomiting have been observed.

#### 4.11 Withdrawal period(s)

Not applicable.

### 5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Anthelmintics, praziquantel combinations.  
ATC vet code: QP52AA51

#### 5.1 Pharmacodynamic properties

This product contains anthelmintics active against gastrointestinal roundworms and tapeworms. The product contains two active substances, as follows:

1. Pyrantel embonate (pamoate), a tetrahydropyrimidine derivative and
2. Praziquantel, a partially hydrogenated pyrazinoisoquinoline derivative.

Pyrantel acts as a cholinergic agonist. Its mode of action is to stimulate nicotinic cholinergic receptors of the parasite, induce spastic paralysis and thereby allow expulsion from the gastro-intestinal (GI) system by peristalsis.

Praziquantel is very rapidly absorbed and distributed throughout the parasite.

Both in vitro and in vivo studies have shown that praziquantel causes severe damage to the parasite integument, resulting in contraction and paralysis. There is an almost instantaneous tetanic contraction of the parasite musculature and a rapid vacuolization of the syncytial tegument. This rapid contraction has been explained by changes in divalent cation fluxes, especially calcium.

In this fixed combination, pyrantel is active against the following ascarids: *Toxocara cati*, and *Toxascaris leonina*. Praziquantel is effective against tapeworms in particular *Dipylidium caninum* and *Taenia taeniaeformis*.

Since it contains praziquantel, the product is effective against *Echinococcus multilocularis*.

## 5.2 Pharmacokinetic particulars

Praziquantel is rapidly absorbed, metabolised and distributed in the body. It is also believed to be excreted back into the intestinal lumen by the mucous membrane.

Following administration of the product to cats, peak plasma concentrations of praziquantel were achieved by approximately 2 hours.

Pyrantel is poorly absorbed so it is expected that a large proportion of the administered dose remains in the GIT where it exerts its therapeutic effect and it is excreted largely unchanged in the faeces.

Following administration of the product to cats, peak plasma concentrations of pyrantel were achieved by approximately 3 hours.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

#### Core tablet:

Maize starch  
Microcrystalline cellulose  
Crospovidone  
Magnesium stearate  
Colloidal anhydrous silica

#### Film coat

Grilled meat flavour  
Opadry Complete Film Coating System 03F28415 White consisting of HPMC 2910/Hypromellose (E464), Macrogol/PEG 4000 (E1521), Titanium Dioxide (E171).

### 6.2 Major incompatibilities

Not Applicable

### 6.3 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 5 years

#### **6.4 Special precautions for storage**

This veterinary medicinal product does not require any special temperature storage restrictions.

Unused halved tablets should be discarded.

Do not remove tablets from strip packaging until required for use.

Keep blister in outer carton.

#### **6.5 Nature and composition of immediate packaging**

The product is presented in either:

Individual blisters made up of a PVC/PE/PCTFE white opaque copolymer and a 20µm heat seal lacquer/aluminium containing 2tablets.

or

Individual blisters made up of 45µm PVC/aluminium/orientated polyamide and a 20µm heat seal lacquer/aluminium containing 2.

The blisters are packed into cartons containing 2tablets

Not all pack sizes may be marketed.

#### **6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

### **7. MARKETING AUTHORISATION HOLDER**

Chanelle Pharmaceuticals Manufacturing Ltd.  
Loughrea  
Co. Galway  
Ireland

### **8. MARKETING AUTHORISATION NUMBER**

Vm 08749/4068

### **9. DATE OF FIRST AUTHORISATION**

13 May 2016

### **10. DATE OF REVISION OF THE TEXT**

June 2021

Approved: 18/06/21

