

PARTICULARS TO APPEAR ON THE OUTER PACKAGE
Cardboard or plastic box (PET tray)

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Nobivac DHPPi lyophilisate for suspension for injection for dogs

2. STATEMENT OF ACTIVE SUBSTANCES

Each 1 ml dose of reconstituted vaccine contains:

Live canine distemper virus (CDV), strain Onderstepoort	$\geq 10^{4.0}$ TCID ₅₀
Live canine adenovirus type 2 (CAV2), strain Manhattan LPV3	$\geq 10^{4.0}$ TCID ₅₀
Live canine parvovirus (CPV), strain 154	$\geq 10^{7.0}$ TCID ₅₀
Live canine parainfluenza virus (CPI), strain Cornell	$\geq 10^{5.5}$ TCID ₅₀

3. PACKAGE SIZE

10 x 1 dose

50 x 1 dose

4. TARGET SPECIES

Dogs.

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Subcutaneous use.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {mm/yyyy}

Once reconstituted use within 30 minutes.

9. SPECIAL STORAGE PRECAUTIONS

Store in a refrigerator.

Do not freeze.
Protect from light.

Avoid prolonged and repetitive exposure to high ambient temperature.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Intervet International B.V.

14. MARKETING AUTHORISATION NUMBERS

Vm 06376/4104

15. BATCH NUMBER

Lot {number}

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING
UNITS Vial label

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Nobivac DHPPi



2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

1 dose

CDV $\geq 10^{4.0}$ TCID₅₀
CAV2 $\geq 10^{4.0}$ TCID₅₀
CPV $\geq 10^{7.0}$ TCID₅₀
CPi $\geq 10^{5.5}$ TCID₅₀.

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

Once reconstituted use within 30 minutes.

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Nobivac DHPPi lyophilisate for suspension for injection for dogs

2. Composition

Each 1 ml dose of reconstituted vaccine contains:

Active substances:

Live canine distemper virus (CDV), strain Onderstepoort $\geq 10^{4.0}$ TCID₅₀*

Live canine adenovirus 2 (CAV2), strain Manhattan LPV3 $\geq 10^{4.0}$ TCID₅₀*

Live canine parvovirus (CPV), strain 154 $\geq 10^{7.0}$ TCID₅₀*

Live canine parainfluenza virus (CPI), strain Cornell $\geq 10^{5.5}$ TCID₅₀*

* TCID₅₀ = median Tissue Culture Infective Dose

Lyophilisate: off white or cream-coloured pellet.

3. Target species

Dogs.

4. Indications for use

For active immunisation of dogs to prevent mortality and clinical signs caused by canine distemper virus infection. To reduce clinical signs of infectious hepatitis and viral excretion due to canine adenovirus type 1 infection. To prevent mortality, clinical signs and viral excretion following canine parvovirus infection. To reduce clinical signs and viral excretion caused by canine parainfluenza virus infection and to reduce clinical signs of respiratory disease and viral excretion following adenovirus type 2 infection.

Onset of immunity:

Canine distemper virus, canine adenovirus and canine parvovirus vaccine components: 1 week.

Canine parainfluenza virus vaccine component: 4 weeks.

Duration of immunity:

Canine distemper virus, canine adenovirus and canine parvovirus vaccine components: 3 years.

The duration of immunity for the canine parainfluenza virus component has not been demonstrated, but an anamnestic response is produced in dogs revaccinated one year after basic vaccination. Annual revaccination with the canine parainfluenza virus vaccine component is recommended.

5. Contraindications

None.

6. Special warnings

Special warnings:

Vaccinate healthy animals only.

The vaccine may not be effective in dogs incubating the disease at the time of vaccination.

A good immune response is reliant on the reaction of an immunogenic agent and a fully competent immune system. Immunocompetence of the animal may be compromised by a variety of factors including poor health, nutritional status, genetic factors, concurrent drug therapy and stress.

The immunogenicity of the vaccine antigen will be reduced by poor storage or inappropriate administration.

The vaccine has been proved to be of benefit against virulent challenge in the presence of maternal antibody levels to CDV, CAV2, CPV and CPi that are likely to be encountered under field conditions.

Experience has shown that the maternal antibody status of pups within a litter varies greatly and reliance should not be placed on serological examination of the bitch alone.

Special precautions for safe use in the target species:

Animals that have received a corresponding anti-serum or immunosuppressive drugs should not be vaccinated until an interval of at least 4 weeks has elapsed.

Vaccinated dogs may excrete the parvovirus vaccine strain at very low levels for up to 8 days following vaccination. However, there is no evidence of any reversion to virulence of the vaccine strain and therefore no need to separate unvaccinated dogs from contact with recently vaccinated dogs.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

In the case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Pregnancy:

Can be used in pregnant bitches which have previously been vaccinated with the CDV (strain Onderstepoort), CAV2 (strain Manhattan LPV3), CPV (strain 154) and CPi (strain Cornell) antigens included in the Nobivac vaccine range.

Interaction with other medicinal products and other forms of interaction:

Safety and efficacy data (viral excretion) are available which demonstrate that this vaccine can be mixed and administered with the inactivated vaccines in the Nobivac range against canine leptospirosis caused by all or some of the following serovars: *L. interrogans* serogroup Canicola serovar Canicola, *L. interrogans* serogroup Icterohaemorrhagiae serovar Copenhageni, *L. interrogans* serogroup Australis serovar Bratislava, and *L. kirschneri* serogroup Grippotyphosa serovar Bananal/Liangguang.

The product information of the relevant Nobivac vaccines should be consulted before administration of the mixed product. When mixed with Nobivac leptospirosis vaccines at annual revaccination, it has been established that there is no interference with the anamnestic response induced by the injectable canine parainfluenza virus component.

After administration with one of the leptospirosis vaccines, a mild and transient increase in body temperature (≤ 1 °C) may occur for a few days after vaccination, with some pups showing less activity and/or a reduced appetite. A small transient swelling (≤ 4 cm), which can occasionally be firm and painful on palpation, may be observed at the site of injection. Any such swelling will either have disappeared or be clearly diminished by 14 days post-vaccination.

After mixed administration of an overdose of this vaccine and an overdose of the leptospirosis vaccines in the Nobivac range, transient local reactions such as diffuse to firm swellings from 1 to 5 cm in diameter may be observed, usually these will persist no longer than 5 weeks, however some may take a little longer to completely disappear.

Safety data and efficacy data for the canine distemper virus, canine adenovirus and canine parvovirus components of this vaccine are available which demonstrate that this vaccine can be administered at the same time but not mixed with the inactivated vaccine in the Nobivac range against *Bordetella bronchiseptica*.

When this vaccine is administered in association with the inactivated vaccine in the Nobivac range against *Bordetella bronchiseptica*, the demonstrated antibody response data for the live canine parainfluenza component of this vaccine are the same as when this vaccine is administered alone.

When this vaccine is used with any of the other Nobivac vaccines referred to above, the minimum vaccination age for each vaccine must be taken into account such that at the time of vaccination, the dogs are at or older than the oldest minimum vaccination age for the individual vaccines.

Safety and efficacy data are available which demonstrate that this vaccine can also be administered on the same day with Nobivac Solvent or Nobivac Rabies. These can be used to reconstitute this freeze-dried vaccine.

No information is available on the compatibility of this vaccine when used with any other veterinary medicinal product except the products mentioned above. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Overdose:

No clinical signs other than those indicated in section 7.
In some dogs the swelling may be more painful or may be observed for a longer period.

Major incompatibilities:

Do not mix with any other veterinary medicinal products except solvent supplied for use with the veterinary medicinal product or other Nobivac dog vaccines mentioned in section 3.8.

7. Adverse events

Dogs:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Injection site swelling. ¹ Elevated temperature. ² Hypersensitivity reaction (e.g. lethargy, facial oedema, pruritus, vomiting or diarrhoea). ³
--	--

¹ Small and transient (≤ 5 cm), which can occasionally be firm and painful on palpation. Any such swelling will either have disappeared or be clearly diminished by 14 days post-vaccination.

² Transient.

³ Such reaction may evolve to a more severe condition (anaphylaxis), which may be life-threatening with additional signs like ataxia, dyspnoea, tremor and collapse. If such reactions occur, appropriate treatment is recommended.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or the local representative of the marketing authorisation holder using the contact details at the end of this leaflet, or via your national reporting system at:

Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

e-mail: adverse.events@vmd.gov.uk

8. Dosage for each species, routes and method of administration

Reconstitute the vaccine with 1 ml solvent or 1 ml (1 dose) of the inactivated vaccines mentioned in section 12.

Subcutaneous use.

Maternal antibodies can negatively interfere with the efficacy of a vaccine. Strict adherence to the vaccination programme is therefore recommended.

Vaccination programme:

Primary vaccinations:

A single injection should establish active immunity to canine distemper, infectious canine hepatitis and disease caused by canine parvovirus infection in dogs of 10 weeks of age or older. Where earlier protection is required a first dose may be given to puppies from 6 weeks of age, but because maternally derived passive antibody can interfere with the response to vaccination a final dose at 10 weeks of age or older is generally recommended. For an optimal response to the parainfluenza component, animals should be vaccinated twice, 2 - 4 weeks apart with the final vaccination at 10 weeks of age or more.

If the initial primary course dose of Nobivac DHPPi is delayed to 10 weeks of age or older, a single dose of Nobivac Pi at 12 weeks of age or older should suffice to establish immunity for this component.

Revaccination:

It is recommended that dogs be revaccinated with canine distemper virus, canine adenovirus and canine parvovirus every 3 years and against canine parainfluenza virus every year.

It was not possible to produce clinical signs of kennel cough by parainfluenza challenge in adult dogs and duration of immunity could not therefore be demonstrated, but an anamnestic response was seen in dogs given a booster one year after primary vaccination. Revaccination against parainfluenza is recommended prior to exposure to high risk environments (such as kennelling, showing or mixing with dogs of unknown vaccination history).

Reconstituted product: off-pink or pink coloured suspension.

9. Advice on correct administration

Avoid contamination of vaccine with traces of chemical sterilising agents. Do not use chemicals such as disinfectant or spirit to disinfect the skin prior to inoculation.

10. Withdrawal periods

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.

Store in a refrigerator (2 °C – 8 °C).

Do not freeze.

Protect from light.

Avoid prolonged or repetitive exposure to high ambient temperatures following withdrawal from the refrigerator prior to use. In hot summer conditions vaccine potency can be severely reduced within a few hours.

Do not use this veterinary medicinal product after the expiry date which is stated on the label and outer packaging after Exp. The expiry date refers to the last day of that month.

Shelf life after reconstitution according to directions: 30 minutes.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. Marketing authorisation numbers and pack sizes

Vm 06376/4104

Cardboard or plastic box with 10 x 1 dose or 50 x 1 dose vials.
Not all pack sizes may be marketed.

15. PID link (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. Contact details

Marketing authorisation holder:
Intervet International B.V.
Wim de Körverstraat 35
5831 AN Boxmeer
Netherlands

Manufacturer responsible for batch release:

Intervet International B.V.
Wim de Körverstraat 35
5831 AN Boxmeer
Netherlands

Local representative:

MSD Animal Health UK Limited
Walton Manor, Walton
Milton Keynes
MK7 7AJ, United Kingdom

Contact details to report suspected adverse reactions:

UK(GB)

MSD Animal Health UK
Ltd.
Tel.: +44(0)1908 685685

UK(NI)

Intervet Ireland Ltd.
Tel.: +353(0)1 297 0220

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

17. Other information

POM-V

Gavin Hall

Approved: 02 December 2025