PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING WHERE THERE IS NO PACKAGE LEAFLET, i.e. Combined label and package leaflet {Bottle labels}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Dysect Sheep Pour-On solution 12.5 g/l

2. COMPOSITION

Each litre contains:

Active substance:
Alphacypermethrin 12.5 g

Excipient:
Copper Chlorophyll Complex (E141) 0.6 g

Clear dark green liquid free from solids.

3. PACKAGE SIZE

2.5 litres.
5 litres.

4. TARGET SPECIES

Sheep.

5. INDICATIONS FOR USE

Indications for use

For the treatment and control of blowfly strike in sheep and lambs, and for the treatment of lice and ticks on sheep.

6. CONTRAINDICATIONS

Contraindications

Do not treat lambs under one week old.
Do not treat animals in very hot weather.

7. SPECIAL WARNINGS

Special warnings

Special precautions for safe use in the target species:
For external use only.

Because alphacypermethrin is retained in the wool grease of the fleece, it is recommended that the product be applied to sheep with a minimum wool length of 1.0 cm.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:
Personal protective equipment consisting of waterproof apron, boots, and impervious gloved must be worn when handling the veterinary medicinal product and recently treated sheep.
Use in a well-ventilated area.
Wash hands and exposed skin before eating, drinking, or smoking, and after work.
Wash splashes from skin and eyes immediately.
Remove heavily contaminated clothing immediately and wash before re-use.

Pregnancy and lactation:
The safety of the veterinary medicinal product has not been established during pregnancy and lactation.

Overdose:
No adverse reactions were recorded in double dose exposure trials.

8. ADVERSE EVENTS

Adverse events

Sheep:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Fleece discolouration[1]

[1] Observed from approximately 6 to 8 weeks after treatment.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed on this label, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder using the contact details on this label, or via your national reporting system at:

Website: https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine

e-mail: adverse.events@vmd.gov.uk

9. DOSAGE FOR EACH TARGET SPECIES, ROUTES AND METHOD OF ADMINISTRATION

Dosage for each species, routes and method of administration

READ THE INSTRUCTIONS CAREFULLY BEFORE USE

	Dosage, ml	
Pest	**Sheep (> 25kg)**	**Lambs (< 25kg)**
Blowfly	40	25
Lice	40	
Ticks	40	
Headfly	5	5

For dosages of 40 ml and 25 ml, apply as two half-doses. Apply the first half-dose evenly along the back line from the neck to the saddle, and the second half-dose evenly from the saddle to the tail head and around the rump.

For the control of blowfly, it is important that there is good coverage around the tail area. The T-bar must be held against the fleece so that the dose is applied directly into it. For the treatment of existing strike, apply the dose directly to the struck area. Most strikes will require the application of 5 ml (saucer size) or 10 ml (dinner plate size).

For the reduction of headfly strike, 5 ml per animal is applied between the horns. Because alphacypermethrin is retained in the wool grease of the fleece, it is recommended that the product be applied to sheep with a minimum wool length of 1.0 cm.

10. ADVICE ON CORRECT ADMINISTRATION

Advice on correct administration

Remove the transit cap from the bottle and pierce the seal. Fit the spouted cap to the bottle. Fit the tube from the applicator gun to the spouted cap and fit the metal spring over the tube on the cap spout to ensure it is fixed securely. Check the gun is fitted with the T-bar and that it is working correctly. Before use the gun must be primed and set to the required volume. Clean the gun after use and maintain the gun to ensure accurate dosing.

The SimcroTech 25ml Pour-On Applicator gun is recommended as the product may have a detrimental effect on certain parts of conventional dosing guns.

Ready to use topical insecticidal solution. Treats a maximum of 124 ewes or a maximum of 200 lambs.

Contains the pyrethroid alphacypermethrin, which has both a repellent and insecticidal action to protect ewes and lambs from blowfly strike. The product has been formulated as a ready to use pour-on. Treated animals are easily identified by the natural green colour of the product which will fade after a few days without staining the fleece.

For the best results, apply before the spring or autumn tick rise and before the first appearance of headfly or blowfly; treat for lice when they are present. To ensure the best results the whole flock should be treated.

Use:
Blowfly – For the control of blowfly a single application will normally provide 8 to 10 weeks protection. Re-treatment may be necessary after this period.
For the treatment of blowfly strike, a single treatment applied to the infected area will ensure blowfly larvae are killed.
Ticks – A single treatment will normally provide control of ticks for 8 to 12 weeks.
Headfly – A single treatment will reduce the incidence for up to 6 weeks.
Lice – A single treatment will normally kill all lice.

11. WITHDRAWAL PERIODS

Withdrawal periods
Meat and offal: 49 days.
Milk: Not permitted for use in animals producing milk for human consumption.

12. SPECIAL STORAGE PRECAUTIONS

Special storage precautions

Keep out of the sight and reach of children.

Do not store above 25 °C.
Do not freeze.
If contents freeze, they must be thawed and thoroughly mixed before use.
Store in the original container.
Keep the container tightly closed.
Store in a dry place.
Protect from direct sunlight.
Store away from food, drink, and animal feeding stuffs.

Do not use this veterinary medicinal product after the expiry date which is stated on the label. The expiry date refers to the last day of that month.

13. SPECIAL PRECAUTIONS FOR DISPOSAL

Special precautions for disposal

Medicines should not be disposed of via wastewater.

DANGEROUS to fish and aquatic life. Do not contaminate ponds, waterways or ditches with the product or empty container.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

14. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

15. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 42058/5160

Pack sizes: 2.5 litre and 5 litres.

Not all pack sizes may be marketed.

16. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

17. CONTACT DETAILS

Marketing authorisation holder and contact details to report suspected adverse reactions:
Zoetis UK Limited
1st Floor, Birchwood Building
Springfield Drive
Leatherhead
Surrey
KT22 7LP
United Kingdom
Tel: +44 (0) 345 300 8034

Manufacturer responsible for batch release:
Zoetis Manufacturing Italia S.R.L.
Catania Branch
Via Franco Gorgone – Zona Industriale
1-95100 Catania CT
Italy'

18. OTHER INFORMATION

POM-VPS

19. THE WORDS "FOR ANIMAL TREATMENT ONLY"

For animal treatment only.

20. EXPIRY DATE

Exp. {mm/yyyy}

21. BATCH NUMBER

Lot {number}

Gavin Hall

Approved: 06 January 2026