

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

10 kg BAG

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Pulmotil G200 Premix for medicated feeding stuff

2. STATEMENT OF ACTIVE SUBSTANCES

Active substance: Tilmicosin (as phosphate) 200 g/kg

3. PHARMACEUTICAL FORM

Premix for medicated feeding stuff
Pharmaceutical description: A yellowish tan to reddish tan free-flowing granular material for in-feed use.

4. PACKAGE SIZE

Net Weight: 10 kg

5. TARGET SPECIES

For use in pigs and rabbits.

6. INDICATION(S)

Pigs: Prevention and treatment of respiratory disease caused by *Actinobacillus pleuropneumoniae*, *Mycoplasma hyopneumoniae*, *Pasteurella multocida* and other organisms sensitive to tilmicosin.

Rabbits: Prevention and treatment of respiratory disease caused by *Pasteurella multocida* and *Bordetella bronchiseptica*, susceptible to tilmicosin.

7. METHOD AND ROUTE(S) OF ADMINISTRATION

The uptake of medicated feed depends on the clinical condition of the animals. In order to obtain a correct dosage the concentration of tilmicosin has to be adjusted accordingly.

Use the following formula:

$$\text{Kg Premix/tonne feed} = \frac{\text{Dose rate (mg/kg bodyweight)} \times \text{bodyweight (kg)}}{\text{Daily feed intake (kg)} \times \text{premix strength (g/kg)}}$$

Pigs

Administer in the feed at a dose of 8 to 16 mg/kg body weight/day of tilmicosin activity (equivalent to 200 to 400 ppm in the feed) for a period of 15 to 21 days.

Indication	Dose of tilmicosin	Duration of treatment	Inclusion rate in feed
Prevention and treatment of respiratory disease	8-16 mg/kg bodyweight /day	15 to 21 days	1-2 kg Pulmotil G200 Premix/tonne

Rabbits

Administer in the feed at 12.5 mg/kg body weight/day of tilmicosin (equivalent to 200 ppm in the feed) for 7 days.

Indication	Dose of tilmicosin	Duration of treatment	Inclusion rate in feed
Prevention and treatment of respiratory disease	12.5 mg/kg bodyweight /day	7 days	1 kg Pulmotil G200 Premix/tonne

Not to be incorporated into feeds containing Bentonite.

To ensure thorough dispersion of the product, it should first be mixed with a suitable quantity of feed ingredients (20 to 50 kg) before incorporation into the finished feed. This product can be incorporated into pelleted feed, preconditioned for the minimum time-period at a temperature not exceeding 75°C.

Official, national and regional antimicrobial policies should be taken into account when the product is used.

Due to the likely variability (time, geographical) in the occurrence of the resistance of bacteria for tilmicosin, bacteriological sampling and susceptibility testing are recommended.

8. WITHDRAWAL PERIOD(S)

Withdrawal period(s):

Pigs: 21 days

Rabbits: 4 days

9. SPECIAL WARNING(S), IF NECESSARY

Special warnings

The safety of tilmicosin has not been established in boars used for breeding purposes.

No symptoms of overdose have been seen in pigs fed a ration containing levels of tilmicosin up to 10 times the recommended dose for 15 days.

For the user:

When mixing the veterinary medicinal product and handling the medicated feed, direct contact with eyes, skin and mucous membranes should be avoided.

Personal protective equipment should be worn. In case of accidental ingestion seek medical advice immediately and show the label to the physician.

- Tilmicosin may induce irritation. Macrolides, such as tilmicosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tilmicosin may lead to cross reactions to other macrolides and vice versa.

Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

- To avoid exposure during preparation of the medicated feed, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN140 with a filter to EN143. Do not eat, drink or smoke when handling this product. Wash hands after use.
 - In the case of accidental ingestion, wash out mouth immediately with water and seek medical advice. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.
 - Do not handle the product if you are allergic to ingredients in the product.
 - If you develop symptoms after exposure, such as skin rash, you should seek medical advice and show the physician this warning.
- Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Adverse reactions

In very rare cases, feed intake may decrease (including feed refusal) in animals receiving medicated feed. This effect is transient.

If you notice any serious effects or other effects not mentioned, please inform your veterinary surgeon.

Contraindications

Do not allow horses and other equines access to medicated feed containing tilmicosin.

Do not use in case of hypersensitivity to tilmicosin or to any of the excipients.

10. EXPIRY DATE

EXP {month/year}

11. SPECIAL STORAGE CONDITIONS

Once opened, use within 3 months. Once incorporated into meal or pelleted feed, use within 3 months.

Store in a dry place.

Do not store above 25°C.

Protect from direct sunlight.

Do not use after the expiry date which is stated on the bag after 'EXP'.

12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

Dispose of any unused product and empty containers in accordance with guidance from your national waste regulation authority.

Pig manure should not be spread onto grass land and when spread onto arable land should be ploughed to a depth of 30 cm.

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE

For animal treatment only. To be supplied only on veterinary prescription.

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Elanco Europe Ltd.
Form 2, Bartley Way
Bartley Wood Business Park
Hook
RG27 9XA
United Kingdom

16. MARKETING AUTHORISATION NUMBER

Vm 00879/4170

17. MANUFACTURER’S BATCH NUMBER

Lot {number}

OTHER INFORMATION


Pulmotil G200 Premix is available in bags containing 1 kg and 10 kg. Not all pack sizes may be marketed.

Manufacturer for the batch release:

TriRX Speke Limited
Fleming Road
Speke
Liverpool
L24 9LN, UK

Elanco France S.A.S
26 rue de la Chapelle
68330 Huingue
France

Date on which the information was last approved: April 2022



Approved: 12 April 2022